## 1. POPULATION

| Population | < 1 | | Total < 1 | 1 to 4 | | Total 1 to 4 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | ≥ 60 | | Total ≥ 60 | Grand total |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | M | F | | M | F | | M | F | | M | F | | M | F | | |
| Refugee population | 749 | 708 | 1,457 | 2,862 | 2,964 | 5,826 | 8,815 | 8,366 | 17,180 | 8,140 | 8,234 | 16,374 | 325 | 471 | 796 | 41,634 |
| % by age group | | | 4% | | | 14% | | | 41% | | | 39% | | | 2% | 100% |

## 2. MORTALITY

### 2.1 Mortality

| | Record ONE cause of mortality per case | < 1 | | Total < 1 | 1 to 4 | | Total 1 to 4 | U5 % mort | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | ≥ 60 | | Total ≥ 60 | Grand total | Crude % mort |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | M | F | | M | F | | | M | F | | M | F | | M | F | | | |
| 1 | Malaria (suspected) | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 2 | Malaria (confirmed) | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3 | URTI | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4 | LRTI | 0 | 1 | 1 | 0 | 0 | 0 | 17% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 8% |
| 5 | Neonatal death | 1 | 1 | 2 | 0 | 0 | 0 | 33% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 15% |
| 6 | Sexually transmitted infections | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 7 | Watery diarrhoea | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 8 | Bloody diarrhoea | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 9 | Tuberculosis | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 10 | Meningitis (Non-meningococcal) | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 11 | HIV/AIDS | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 12 | Acute malnutrition | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 13 | Chronic malnutrition | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 14 | Anaemia | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 15 | Acute Jaundice Syndrome | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 16 | Asthma | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 17 | Cancer | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 18 | Cardiovascular disease (Non-IHD) | 0 | 1 | 1 | 0 | 0 | 0 | 17% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 2 | 15% |
| 19 | Ischaemic heart disease | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 20 | Chronic obstructive pulmonary disease (non-asthma) | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 21 | Diabetes | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 22 | Hypertension | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 23 | Liver disease | 0 | 1 | 1 | 0 | 0 | 0 | 17% | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 2 | 15% |
| 24 | Musculoskeletal disorders | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 25 | Mental health | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 26 | Maternal Death - Hemorrhage | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 8% |
| 27 | Maternal Death - Hypertensive disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 28 | Maternal Death - Infections | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 29 | Maternal Death - Abortions | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 30 | Maternal Death - Embolism | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 31 | Maternal Death - Other direct causes | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 32 | Maternal Death - Indirect causes | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 33 | COVID-19 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 34 | Acute Flaccid Paralysis | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 35 | Cholera | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 36 | Yellow Fever | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 37 | Viral Hemorrhagic Fever | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 38 | Plague | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 39 | Measles | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 40 | Meningococcal Meningitis | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 41 | Salmonelosis | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 42 | Shigella | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 43 | Tetanus | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 44 | Varicella | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 45 | Whooping cough | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 46 | Monkeypox | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 47 | Assault | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 48 | Bites | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 49 | Accidents | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 50 | Self-harm | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 51 | Suicidal attempt | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 52 | Other | 0 | 1 | 1 | 0 | 0 | 0 | 17% | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 2 | 3 | 5 | 38% |
| | **Total** | 1 | 5 | 6 | 0 | 0 | 0 | 100% | 0 | 1 | 1 | 1 | 1 | 2 | 1 | 3 | 4 | 13 | 100% |
| | % by age group | | | 46% | | | 0% | | | | 8% | | | 15% | | | 31% | 100% | |

| Mortality indicators | Male | Female | Total |
|---|---|---|---|
| Crude Mortality Rate (CMR) (/1000/month) | 0.0 | 0.2 | 0.10 |
| Under 5 Mortality Rate (U5MR) (/1000/month) | 0.1 | 0.5 | 0.3 |
| Infant Mortality Rate (IMR) (/1000/month) | 0.4 | 2.4 | 1.4 |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | | | 10.0 |

### 2.2 Maternal death

| | |
|---|---|
| Number of maternal deaths reported | 1 |
| Number of maternal deaths investigated | 1 |
| Proportion of maternal deaths investigated | 100% |

## 3. MORBIDITY

### 3.1 Staffing and service availability

| | |
|---|---|
| Number of full-time clinicians | 21.4 |
| Average number of full days on which each health facility was functioning | 5.4 |
| Total number of outpatient consultations | 59,786 |
| Health facility utilisation rate | 4.3 |
| Consultations per clinician per day | 40 |

| OUTPATIENT CONSULTATIONS | Male | Female | TOTAL |
|---|---|---|---|
| New Consultations | 20,438 | 24,488 | 44,926 |
| Re-visit Consultations | 6,520 | 8,340 | 14,860 |
| TOTAL Consultations | 26,958 | 32,828 | 59,786 |

### 3.2 Acute health conditions

| | Record NEW cases of disease only | 0 to 4 | | Total <5 | U5 Incid | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | ≥ 60 | | Total ≥ 60 | Grand total | % by cause | Crude Incide |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | M | F | | | M | F | | M | F | | M | F | | | | |
| 1 | URTI | 2,795 | 2,407 | 5,202 | 714.2 | 3,077 | 3,026 | 6,103 | 2,209 | 2,770 | 4,979 | 183 | 220 | 403 | 16,687 | 38% | 400.8 |
| 2 | ILI | 177 | 163 | 340 | 46.7 | 91 | 127 | 218 | 60 | 190 | 250 | 8 | 16 | 24 | 832 | 2% | 20.0 |
| 3 | LRTI | 322 | 209 | 531 | 72.9 | 139 | 102 | 241 | 82 | 101 | 183 | 12 | 22 | 34 | 989 | 2% | 23.8 |
| 4 | Watery diarrhoea | 187 | 139 | 326 | 44.8 | 54 | 34 | 88 | 19 | 27 | 46 | 0 | 3 | 3 | 463 | 1% | 11.1 |
| 5 | Bloody diarrhoea | 2 | 0 | 2 | 0.3 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 3 | 0% | 0.1 |
| 6 | Skin disease | 213 | 234 | 447 | 61.4 | 257 | 397 | 654 | 194 | 396 | 590 | 12 | 31 | 43 | 1,734 | 4% | 41.6 |
| 7 | Eye disease | 95 | 84 | 179 | 24.6 | 117 | 98 | 215 | 204 | 130 | 334 | 12 | 20 | 32 | 760 | 2% | 18.3 |
| 8 | Ear disease | 195 | 165 | 360 | 49.4 | 152 | 186 | 338 | 171 | 370 | 541 | 18 | 36 | 54 | 1,293 | 3% | 31.1 |
| 9 | Dental conditions | 31 | 46 | 77 | 10.6 | 989 | 1,116 | 2,105 | 1,379 | 2,738 | 4,117 | 64 | 85 | 149 | 6,448 | 15% | 154.9 |
| 10 | Intestinal worms | 73 | 72 | 145 | 19.9 | 210 | 204 | 414 | 54 | 77 | 131 | 0 | 1 | 1 | 691 | 2% | 16.6 |
| 11 | Acute Jaundice Syndrome | 2 | 0 | 2 | 0.3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% | 0.0 |
| 12 | Urinary Tract Infection | 62 | 143 | 205 | 28.1 | 116 | 370 | 486 | 283 | 584 | 867 | 20 | 52 | 72 | 1,630 | 4% | 39.2 |
| 13 | Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 14 | Measles (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 15 | Meningitis (suspected) | 5 | 0 | 5 | 0.7 | 5 | 2 | 7 | 0 | 0 | 0 | 0 | 0 | 0 | 12 | 0% | 0.3 |
| 16 | Acute malnutrition- Moderate | 0 | 0 | 0 | 0.0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% | 0.0 |
| 17 | Acute malnutrition- Severe | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 18 | Anaemia | 114 | 95 | 209 | 28.7 | 84 | 108 | 192 | 22 | 185 | 207 | 0 | 6 | 6 | 614 | 1% | 14.7 |
| 19 | Abortion | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 20 | HIV/AIDS | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 21 | Other | 1,030 | 903 | 1,933 | 265.4 | 1,486 | 1,605 | 3,091 | 2,490 | 3,935 | 6,425 | 279 | 447 | 726 | 12,175 | 27% | 292.4 |
| | **Total** | 5,303 | 4,660 | 9,963 | 1367.9 | 6,778 | 7,375 | 14,153 | 7,168 | 11,503 | 18,671 | 608 | 939 | 1,547 | 44,334 | 100% | 1064.9 |
| | % by age group | | | 22% | | | | 32% | | | 42% | | | 3% | 100% | | |

### 3.3 Notifiable diseases

| | Record NEW cases of disease only | 0 to 4 | | Total <5 | U5 Incid | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | ≥ 60 | | Total ≥ 60 | Grand total | % by cause | Crude Incide |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | M | F | | | M | F | | M | F | | M | F | | | | |
| 1 | COVID-19 (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 2 | COVID-19 (confirmed) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 3 | Acute Flaccid Paralysis | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 4 | Cholera | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 5 | Yellow Fever | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 6 | Viral Hemorrhagic Fever | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 7 | Plague | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 8 | Measles | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 9 | Meningococcal Meningitis | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 10 | Salmonelosis | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 11 | Shigella | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 12 | Tetanus | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 13 | Varicella | 7 | 7 | 14 | 1.9 | 53 | 47 | 100 | 0 | 2 | 2 | 0 | 0 | 0 | 116 | 100% | 2.8 |
| 14 | Whooping cough | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| 15 | Monkeypox | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| | **Total** | 7 | 7 | 14 | 1.9 | 53 | 47 | 100 | 0 | 2 | 2 | 0 | 0 | 0 | 116 | 100% | 2.8 |
| | % by age group | | | 12% | | | | 86% | | | 2% | | | 0% | 100% | | |

### 3.4 Chronic health conditions

| | Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | ≥ 60 | | Total ≥ 60 | Grand total | % by cause |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | M | F | | M | F | | M | F | | M | F | | | |
| 1 | Diabetes | 2 | 1 | 3 | 4 | 17 | 21 | 138 | 251 | 389 | 89 | 160 | 249 | 662 | 20% |
| 2 | Hypertension | 0 | 0 | 0 | 1 | 0 | 1 | 281 | 377 | 658 | 153 | 247 | 400 | 1,059 | 33% |
| 3 | Ischaemic heart disease | 0 | 0 | 0 | 0 | 0 | 0 | 52 | 13 | 65 | 26 | 26 | 52 | 117 | 4% |
| 4 | Cardiovascular disease (other) | 0 | 0 | 0 | 7 | 1 | 8 | 9 | 18 | 27 | 4 | 2 | 6 | 41 | 1% |
| 5 | Asthma | 37 | 14 | 51 | 56 | 38 | 94 | 38 | 62 | 100 | 4 | 12 | 16 | 261 | 8% |
| 6 | Chronic obstructive pulmonary disease | 0 | 0 | 0 | 0 | 0 | 0 | 22 | 4 | 26 | 11 | 16 | 27 | 53 | 2% |
| 7 | Thyroid disorders | 15 | 10 | 25 | 20 | 30 | 50 | 25 | 160 | 185 | 4 | 14 | 18 | 278 | 9% |
| 8 | Cancer | 0 | 0 | 0 | 0 | 0 | 0 | 7 | 2 | 9 | 3 | 0 | 3 | 12 | 0% |
| 9 | Liver disease | 0 | 3 | 3 | 0 | 0 | 0 | 1 | 2 | 3 | 3 | 0 | 3 | 9 | 0% |
| 10 | Musculoskeletal disorders | 4 | 3 | 7 | 4 | 10 | 14 | 34 | 76 | 110 | 6 | 10 | 16 | 147 | 5% |
| 11 | Other | 29 | 21 | 50 | 65 | 70 | 135 | 155 | 219 | 374 | 24 | 16 | 40 | 599 | 18% |
| | **Total** | 87 | 52 | 139 | 157 | 166 | 323 | 762 | 1,184 | 1,946 | 327 | 503 | 830 | 3,238 | 100% |
| | % by age group | | | 4% | | | 10% | | | 60% | | | 26% | 100% | |

### 3.5 Mental health

| | | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | ≥ 60 | | Total ≥ 60 | Grand total | % by cause |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | M | F | | M | F | | M | F | | M | F | | | |
| 1 | Epilepsy/Seizures | 43 | 49 | 92 | 127 | 78 | 205 | 94 | 72 | 166 | 5 | 8 | 13 | 476 | 30% |
| 2 | Alcohol/Substance abuse | 0 | 0 | 0 | 0 | 0 | 0 | 10 | 0 | 10 | 0 | 0 | 0 | 10 | 1% |
| 3 | Intellectual disability and development disorders | 10 | 3 | 13 | 49 | 17 | 66 | 17 | 11 | 28 | 0 | 1 | 1 | 108 | 7% |
| 4 | Psychosis (including bipolar disorder) | 0 | 0 | 0 | 0 | 4 | 4 | 75 | 68 | 143 | 10 | 21 | 31 | 178 | 11% |
| 5 | Moderate-severe emotional disorder | 1 | 0 | 1 | 96 | 65 | 161 | 130 | 299 | 429 | 30 | 8 | 38 | 629 | 40% |
| 6 | Medically unexplained somatic complaint | 0 | 0 | 0 | 1 | 2 | 3 | 11 | 7 | 18 | 0 | 0 | 0 | 21 | 1% |
| 7 | Dementia or delirium | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 14 | 17 | 17 | 1% |
| 8 | Other psychological complaint | 3 | 1 | 4 | 66 | 21 | 87 | 19 | 17 | 36 | 0 | 1 | 1 | 128 | 8% |
| | **Total** | 57 | 53 | 110 | 339 | 187 | 526 | 356 | 474 | 830 | 48 | 53 | 101 | 1,567 | 100% |
| | % by age group | | | 7% | | | 34% | | | 53% | | | 6% | 100% | |

### 3.6 Injuries

| | | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | ≥ 60 | | Total ≥ 60 | Grand total | % by cause |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | M | F | | M | F | | M | F | | M | F | | | |
| 1 | Burns | 5 | 7 | 12 | 6 | 8 | 14 | 5 | 21 | 26 | 5 | 2 | 7 | 59 | 12% |
| 2 | Bites | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 3 | 4 | 0 | 0 | 0 | 5 | 1% |
| 3 | Assault | 1 | 2 | 3 | 12 | 1 | 13 | 2 | 1 | 3 | 0 | 1 | 1 | 20 | 4% |
| 4 | Accidents | 26 | 11 | 37 | 107 | 40 | 147 | 56 | 26 | 82 | 1 | 1 | 2 | 268 | 53% |
| 5 | Self-harm | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 6 | Suicidal attempt | 0 | 1 | 1 | 2 | 1 | 3 | 0 | 1 | 1 | 0 | 0 | 0 | 5 | 1% |
| 7 | Other | 12 | 14 | 26 | 60 | 18 | 78 | 28 | 19 | 47 | 0 | 1 | 1 | 152 | 30% |
| | **Total** | 44 | 35 | 79 | 188 | 68 | 256 | 92 | 71 | 163 | 6 | 5 | 11 | 509 | 100% |
| | % by age group | | | 16% | | | 50% | | | 32% | | | 2% | 100% | |

### 3.7 Reproductive Tract Infections

| | | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | ≥ 60 | | Total ≥ 60 | Grand total | % by cause |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | M | F | | M | F | | M | F | | M | F | | | |
| 1 | Urethral discharge syndrome | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 2 | Vaginal discharge syndrome | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 340 | 340 | 0 | 0 | 0 | 340 | 57% |
| 3 | Genital ulcer disease | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4 | Pelvic inflammatory disease | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 5 | Congenital syphilis | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 6 | Ophthalmia Neonatorum | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 7 | Other | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 253 | 253 | 0 | 0 | 0 | 254 | 43% |
| | **Total** | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 593 | 593 | 0 | 0 | 0 | 594 | 100% |
| | % by age group | | | 0% | | | 0% | | | 100% | | | 0% | 100% | |

## 4. INPATIENT DEPARTMENT (In-Camp)

### 4.1 In-patient Department Activities

| | | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | ≥ 60 | | Total ≥ 60 | Grand total |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | M | F | | M | F | | M | F | | M | F | | |
| Number at beginning of reporting period | | 5 | 4 | 9 | 0 | 2 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 13 |
| **Total admissions** | | 132 | 79 | 211 | 40 | 52 | 92 | 18 | 187 | 205 | 0 | 0 | 0 | 508 |
| Number of exits | authorized | 125 | 72 | 197 | 35 | 47 | 82 | 18 | 182 | 200 | 0 | 0 | 0 | 479 |
| | unauthorized | 0 | 2 | 2 | 1 | 0 | 1 | 0 | 4 | 4 | 0 | 0 | 0 | 7 |
| | death | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | referral | 10 | 4 | 14 | 2 | 7 | 9 | 0 | 0 | 0 | 0 | 0 | 0 | 23 |
| **Total exits** | | 135 | 78 | 213 | 38 | 54 | 92 | 18 | 186 | 204 | 0 | 0 | 0 | 509 |
| Deaths within 24 hours | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| | |
|---|---|
| Number of days in reporting period | 91 |
| Number of beds | 18 |
| Sum number of days stay for authorised discharges | 987 |
| Average length of stay (days) | **2.1** |
| Bed occupancy | **64%** |
| Hospitalisation rate (/1000/month) | **4.1** |

### 4.2 In-patient Admissions and Deaths

| | | 0 to 4 | | U5 CFR | 5 to 17 | | 5-17 CFR | 18 to 59 | | 18-59 CFR | ≥ 60 | | Total adm | Total deaths | Crude CFR |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Adm | Deaths | | Adm | Deaths | | Adm | Deaths | | Adm | Deaths | | | |
| 1 | Paediatric medicine | 164 | 0 | | 61 | 0 | | 0 | 0 | | 0 | 0 | 225 | 0 | |
| 2 | General Surgery | 47 | 0 | | 24 | 0 | | 45 | 0 | | 0 | 0 | 116 | 0 | |
| 3 | Internal medicine | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | 0 | 0 | |
| 4 | Obstetrics | 0 | 0 | | 7 | 0 | | 155 | 0 | | 0 | 0 | 162 | 0 | |
| 5 | Gynaecology | 0 | 0 | | 0 | 0 | | 5 | 0 | | 0 | 0 | 5 | 0 | |
| 6 | Other | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | 0 | 0 | |
| | **Total** | 211 | 0 | | 92 | 0 | | 205 | 0 | | 0 | 0 | 508 | 0 | |

*Adm. = Admission; CFR = Case Fatality Rate*

## 5. REFERRALS

### 5.1 Referrals by location

| | | 0 to 4 | 5 to 17 | 18 to 59 | ≥ 60 | Grand total |
|---|---|---|---|---|---|---|
| 1 | MOH Hospital | 143 | 173 | 189 | 29 | 534 |
| 2 | Jordan University Hospital | 0 | 0 | 2 | 0 | 2 |
| 3 | Private affiliated Hospital | 1 | 47 | 512 | 0 | 560 |
| 4 | Royal Medical Services | 0 | 0 | 0 | 0 | 0 |
| 5 | Other | 0 | 0 | 0 | 0 | 0 |

| | |
|---|---|
| Total number of reffered cases out side the camp | 1,096 |
| Referral Rate (/1000/month) | 8.8 |

### 5.2 Referrals by service

| | | 0 to 4 | 5 to 17 | 18 to 59 | ≥ 60 | Grand total |
|---|---|---|---|---|---|---|
| 1 | Paediatrics | 71 | 9 | 0 | 0 | 80 |
| 2 | Internal medicine | 50 | 148 | 611 | 26 | 835 |
| 3 | Obstetric | 0 | 2 | 75 | 0 | 77 |
| 4 | Gynaecological | 0 | 3 | 0 | 0 | 3 |
| 5 | Surgical | 23 | 58 | 17 | 3 | 101 |
| 6 | Other | 0 | 0 | 0 | 0 | 0 |

## 6. LABORATORY

| | Total |
|---|---|
| Number of laboratory tests conducted | 15,847 |

# 7. REPRODUCTIVE HEALTH

## 7.1 Antenatal care

| | | < 18 | ≥ 18 | Total |
|---|---|---|---|---|
| 1 | First antenatal visit (during the first trimester) | 5 | 222 | 227 |
| 2 | First antenatal visit (after the first trimester) | 3 | 38 | 41 |
| 3 | Repeat antenatal visit | 58 | 1,961 | 2,019 |
| 4 | High risk pregnancies detected | 1 | 292 | 293 |
| 5 | Tested for anaemia | 25 | 652 | 677 |
| 6 | Confirmed severe and moderate anaemia | 5 | 85 | 90 |
| 7 | Abortions | 3 | 47 | 50 |

| ANC Indicators | |
|---|---|
| Proportion of first time antenatal visits made < 1st trimester | 85% |

## 7.2 Antenatal care in facility

| | | | | |
|---|---|---|---|---|
| Number of pregnant women at time of delivery who received: | ≥ 4 ANC visits | 7 | 171 | 178 |
| | 2 TT doses | 4 | 59 | 63 |
| | anaemia screening | 8 | 167 | 175 |

| | |
|---|---|
| Coverage of complete antenatal care | 89% |
| Coverage of antenatal tetanus immunisation | 32% |
| Coverage of anaemia screening in pregnancy | 88% |

## 7.3 Delivery care

| | | Health Facility | | Home Delivery | | TOTAL |
|---|---|---|---|---|---|---|
| | | < 18 | ≥ 18 | < 18 | ≥ 18 | |
| 1 | Livebirths | 8 | 191 | 0 | 1 | 200 |
| 2 | Stillbirths | 0 | 1 | 0 | 0 | 1 |
| 3 | Low birth weight (< 2500g) | 0 | 13 | 0 | 0 | 13 |
| 4 | Attended by a skilled health worker | 8 | 191 | 0 | 0 | 199 |
| 5 | Obstetric complications | 0 | 3 | 0 | 0 | 3 |
| 6 | Caesarean sections | 1 | 57 | 0 | 0 | 58 |

| Delivery Indicators | |
|---|---|
| Crude birth rate (/1000 population/ month) | 1.6 |
| Proportion of births attended by skilled health worker (excl. TBA) | 100% |
| Proportion of low birth weight deliveries | 7% |
| Proportion of all births performed by caesarean section | 29% |

## 7.4 Postnatal care

| | < 18 | ≥ 18 | Total |
|---|---|---|---|
| At least three PNC visits within 6 weeks | 0 | 85 | 85 |

| Postnatal Indicators | |
|---|---|
| Coverage of complete postnatal care | 43% |

### 7.5 Family planning

| | New | | Repeat | | Total |
|---|---|---|---|---|---|
| | < 18 | ≥ 18 | < 18 | ≥ 18 | |
| Combined pill (low dose) | 0 | 159 | 0 | 217 | 376 |
| Combined pill (highdose) | 0 | 2 | 0 | 10 | 12 |
| Progesterone pill (Micro-val; Micro-lut) | 1 | 62 | 0 | 85 | 148 |
| Emergency contraceptive pill | 0 | 0 | 0 | 0 | 0 |
| Injectable (Depo-Provera) | 0 | 31 | 0 | 16 | 47 |
| Implantable (Norplant) | 0 | 9 | 0 | 0 | 9 |
| Intra-uterine device (IUD) | 0 | 11 | 0 | 1 | 12 |
| Condom (male) | 0 | 165 | 0 | 112 | 277 |
| Condom (female) | 0 | 8 | 0 | 5 | 13 |
| Sterilisation (male) | 0 | 0 | 0 | 0 | 0 |
| Sterilisation (female) | 0 | 0 | 0 | 0 | 0 |
| Counseling (only) | 5 | 132 | 0 | 214 | 351 |
| Other | 0 | 3 | 0 | 10 | 13 |

### 7.6 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|---|---|---|---|---|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 0 | 0 |

*ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection*

| SGBV Indicators | |
|---|---|
| Incidence of reported rape (/10,000 population/year) | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

## 8. NUTRITION

### 8.1 Outpatient Therapeutic Program

| | | 6 to 23 months | | 24 to 59 months | | ≥ 60 months | |
|---|---|---|---|---|---|---|---|
| | | M | F | M | F | M | F |
| 1 | New admissions | 2 | 4 | 0 | 0 | 0 | 0 |
| 2 | Re-visits | 0 | 0 | 0 | 0 | 0 | 0 |
| 3 | Exits- Recovered (moved to TSFP) | 1 | 0 | 0 | 3 | 0 | 0 |
| 4 | Exits - None response | 2 | 0 | 0 | 0 | 0 | 0 |
| 5 | Exits - Moved to SC | 0 | 0 | 0 | 0 | 0 | 0 |
| 6 | Exits - Moved to other OTP | 3 | 0 | 0 | 0 | 0 | 0 |
| 7 | Exists - Medical Referral | 0 | 0 | 1 | 0 | 0 | 0 |
| 8 | Exits- Death | 0 | 0 | 0 | 0 | 0 | 0 |
| 9 | Exists- Defaulter | 0 | 0 | 0 | 0 | 0 | 0 |
| 10 | Sum number of days stay for discharged children (< 5) | | | 450 | | | |
| 11 | Sum average weight gain for discharged children (< 5) | | | 8 | | | |

## 8.2 Supplementary Feeding Program - U5

| | | 6 to 23 months | | 24 to 59 months | | ≥ 60 months | |
|---|---|---|---|---|---|---|---|
| | | M | F | M | F | M | F |
| 1 | New admissions | 6 | 5 | 0 | 2 | 0 | 0 |
| 2 | Re-visits | 1 | 0 | 1 | 0 | 0 | 0 |
| 3 | Exits- Recovered | 3 | 10 | 4 | 0 | 0 | 0 |
| 4 | Exits - None response | 1 | 1 | 1 | 0 | 0 | 0 |
| 5 | Exits - Moved to SC | 0 | 0 | 0 | 0 | 0 | 0 |
| 6 | Exits - Moved to OTP | 0 | 0 | 1 | 1 | 0 | 0 |
| 7 | Exits - Moved to other TSFP | 0 | 0 | 0 | 0 | 0 | 0 |
| 8 | Exists - Medical Referral | 0 | 0 | 0 | 0 | 0 | 0 |
| 9 | Exits- Death | 0 | 0 | 0 | 0 | 0 | 0 |
| 10 | Exists- Defaulter | 0 | 2 | 1 | 0 | 0 | 0 |
| 11 | Sum number of days stay for discharged children (< 5) | | | 2,203 | | | |
| 12 | Sum average weight gain for discharged children (< 5) | | | 13 | | | |

## 8.3 Supplementary Feeding Program - PLW

| | | < 18 | ≥ 18 |
|---|---|---|---|
| 1 | New admissions | 0 | 7 |
| 2 | Re-visits | 0 | 1 |
| 3 | Exits- Recovered | 0 | 6 |
| 4 | Exits - None response | 0 | 1 |
| 5 | Exits - Moved to other TSFP | 0 | 0 |
| 6 | Exists - Medical Referral | 0 | 0 |
| 7 | Exits- Death | 0 | 0 |
| 8 | Exists- Defaulter | 0 | 11 |

## 8.4 MUAC (Health Facility)

| | | 6 to 23 months | | 24 to 59 months | |
|---|---|---|---|---|---|
| | | M | F | M | F |
| 1 | Children screened - Green (normal) | 335 | 359 | 36 | 19 |
| 2 | Children screened - Yellow (borderline) | 5 | 5 | 0 | 0 |
| 3 | Children screened - Red (danger) | 2 | 5 | 0 | 0 |
| 4 | Children screened - Oedema | 0 | 0 | 0 | 0 |

## 8.5 GMP (Community)

| | | 6 to 23 months | | 24 to 59 months | |
|---|---|---|---|---|---|
| | | M | F | M | F |
| 1 | Children screened - Green (normal) | 159 | 157 | 166 | 162 |
| 2 | Children screened - Yellow (borderline) | 0 | 0 | 0 | 0 |
| 3 | Children screened - Red (danger) | 0 | 0 | 0 | 0 |
| 4 | Children screened - Oedema | 0 | 0 | 0 | 0 |

**Data Source** | Health Information System

UNHCR gratefully acknowledges the support of the following partners: