

Camps reporting this week: Kawergosk, Qushtapa

1. Population

| Population | 0 to < 1 | | Total <1 | 1 to 4 | | Total 1 to 4 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total |
|---------------------------------------|----------|-----|----------|--------|-------|--------------|---------|-------|---------------|----------|-------|----------------|-------|-------|------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | M | F | | |
| Refugee population of reporting camps | 263 | 265 | 528 | 1,052 | 1,063 | 2,115 | 2,322 | 1,922 | 4,244 | 4,012 | 3,516 | 7,528 | 1,121 | 1,074 | 2,195 | 16,610 |
| % by age group | | | 3% | | | 13% | | | 26% | | | 45% | | | 13% | 100% |

* estimates from registered population in Kawergosk and Gawilan camps

2. Mortality

2.1 Mortality

| Record ONE cause of mortality per case | 0 to 4 | | Total <5 | U5 % mort | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | Crude % mort |
|---|----------|----------|----------|-------------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| 1. Lower Respiratory Tract Infection (LRTI) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 2. Watery diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 3. Bloody diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 4. Cardiovascular disorders | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 5. Ischaemic heart disease | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 6. Injury | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 7. Cancer | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 8. Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 9. Tuberculosis (confirmed) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 10. Neonatal death | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Other | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Total | 0 | 0 | 0 | 100% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100% |
| % by age group | | | | | | | | | | | | | | 100% | |

| Mortality indicators | Male | Female | Total |
|--------------------------------|------|--------|-------|
| Crude Mortality Rate (CMR) | 0.0 | 0.0 | 0.0 |
| Under 5 Mortality Rate (U5MR) | 0.0 | 0.0 | 0.0 |
| Neonatal Mortality Rate (NNMR) | | | 0.0 |

2.2 Maternal death

| | |
|--|---|
| Number of maternal deaths reported | 0 |
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3. Morbidity

3.1 Staffing and service availability

| | |
|---|-----|
| Number of full-time clinicians | 6 |
| Average number of full days on which each health facility was | 5.7 |
| Total number of outpatient consultations | 999 |
| Health facility utilisation rate | 3.1 |
| Consultations per clinician per day | 29 |

3.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total <5 | U5 Incid** | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause | Crude Incid** |
|--|------------|------------|------------|--------------|------------|------------|---------------|------------|------------|----------------|-----------|----------|------------|-------------|-------------|---------------|
| | M | F | | | M | F | | M | F | | M | F | | | | |
| Upper Respiratory Tract Infection (URTI) | 72 | 83 | 155 | 58.6 | 65 | 44 | 109 | 57 | 67 | 124 | 0 | 0 | 0 | 388 | 41% | 23.4 |
| Lower Respiratory Tract Infection (LRTI) | 12 | 13 | 25 | 9.5 | 1 | 2 | 3 | 7 | 6 | 13 | 1 | 0 | 1 | 42 | 4% | 2.5 |
| Watery diarrhoea* | 17 | 4 | 21 | 7.9 | 4 | 2 | 6 | 1 | 3 | 4 | 1 | 0 | 1 | 32 | 3% | 1.9 |
| Bloody diarrhoea* | 3 | 1 | 4 | 1.5 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 5 | 1% | 0.3 |
| Malaria (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Skin infection | 4 | 5 | 9 | 3.4 | 3 | 2 | 5 | 12 | 9 | 21 | 0 | 0 | 0 | 35 | 4% | 2.1 |
| Eye infection | 1 | 7 | 8 | 3.0 | 3 | 3 | 6 | 4 | 2 | 6 | 3 | 0 | 3 | 23 | 2% | 1.4 |
| Ear infection | 12 | 15 | 27 | 10.2 | 4 | 6 | 10 | 9 | 9 | 18 | 0 | 0 | 0 | 55 | 6% | 3.3 |
| Dental Conditions | 0 | 1 | 1 | 0.4 | 7 | 7 | 14 | 7 | 9 | 16 | 0 | 0 | 0 | 31 | 3% | 1.9 |
| Intestinal Worms | 0 | 1 | 1 | 0.4 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% | 0.1 |
| Acute Jaundice Syndrome* | 0 | 0 | 0 | 0.0 | 4 | 5 | 9 | 1 | 0 | 1 | 0 | 0 | 0 | 10 | 1% | 0.6 |
| Measles (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Meningitis (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Fever of unknown origin | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Sexually Transmitted Infection (STI) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 0% | 0.1 |
| Urinary tract infection (UTI) | 4 | 2 | 6 | 2.3 | 7 | 8 | 15 | 19 | 16 | 35 | 8 | 1 | 9 | 65 | 7% | 3.9 |
| HIV/AIDS | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute malnutrition | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Leishmaniasis | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Other | 6 | 18 | 24 | 9.1 | 32 | 23 | 55 | 51 | 98 | 149 | 17 | 4 | 21 | 249 | 27% | 15.0 |
| Total | 131 | 150 | 281 | 106.3 | 130 | 103 | 233 | 168 | 221 | 389 | 30 | 5 | 35 | 938 | 100% | 56.5 |
| % by age group | | | 30% | | | | 25% | | | 41% | | | 4% | 100% | | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

**incidence is cases per 1000 per week

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|--|----------|----------|-----------|----------|----------|---------------|----------|-----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| Diabetes | 0 | 0 | 0 | 1 | 1 | 2 | 1 | 1 | 2 | 2 | 0 | 2 | 6 | 17% |
| Hypertension | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 5 | 0 | 1 | 1 | 6 | 17% |
| Asthma | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 2 | 6% |
| Ischaemic heart disease | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 1 | 0 | 1 | 3 | 9% |
| Cardiovascular disease (other) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 0 | 1 | 2 | 6% |
| Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 3% |
| Liver disease | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 2 | 6% |
| Thyroid problems | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 3% |
| Cancer | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 2 | 6% |
| Other | 0 | 0 | 0 | 1 | 0 | 1 | 3 | 5 | 8 | 1 | 0 | 1 | 10 | 29% |
| Total | 1 | 0 | 1 | 3 | 1 | 4 | 7 | 17 | 24 | 5 | 1 | 6 | 35 | 100% |
| % by age group | | | 3% | | | 11% | | | 69% | | | 17% | 100% | |

3.4 Mental health

| | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|---|----------|----------|-----------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Epilepsy / seizures | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 2. Alcohol and other substance abuse disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Psychotic disorder | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 4 | 0 | 0 | 0 | 4 | 22% |
| 4. Mental retardation / intellectual disability | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 5. Severe emotional disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 6. Medically unexplained somatic complaint | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 0 | 0 | 2 | 11% |
| Other | 0 | 0 | 0 | 2 | 5 | 7 | 2 | 2 | 4 | 0 | 1 | 1 | 12 | 67% |
| Total | 0 | 0 | 0 | 2 | 5 | 7 | 4 | 6 | 10 | 0 | 1 | 1 | 18 | 100% |
| % by age group | | | 0% | | | 39% | | | 56% | | | 6% | 100% | |

3.5 Injuries

| | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|---------------------------|----------|----------|-----------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Injury (intentional) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 2. Injury (unintentional) | 1 | 0 | 1 | 1 | 1 | 2 | 1 | 0 | 1 | 0 | 0 | 0 | 4 | 50% |
| 3. Burns | 1 | 2 | 3 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 50% |
| 4. Accident | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Total | 2 | 2 | 4 | 2 | 1 | 3 | 1 | 0 | 1 | 0 | 0 | 0 | 8 | 100% |
| % by age group | | | 50% | | | 38% | | | 13% | | | 0% | 100% | |

4. Referrals

4.1 Referrals - general

| | 0 to 4 | 5 to 14 | 15 to 49 | ≥ 50 | Grand total |
|----------------|----------|----------|-----------|----------|-------------|
| Emergency | 3 | 0 | 18 | 0 | 21 |
| Elective | 5 | 0 | 9 | 0 | 14 |
| Other | 0 | 0 | 0 | 0 | 0 |
| Total | 8 | 0 | 27 | 0 | 35 |
| % by age group | 23% | 0% | 77% | 0% | 100% |

4.2 Referrals - by specialty

| | 0 to 4 | 5 to 14 | 15 to 59 | ≥ 50 | Grand total |
|-------------------|---------|---------|----------|------|-------------|
| Internal medicine | NO DATA | | | | |
| General Surgery | | | | | |
| Gynaecology | | | | | |
| Other | | | | | |
| Total | | | | | |
| % by age group | | | | | |

5. Laboratory & other investigations

| | Total |
|-------------------------|---------|
| Haematology | NO DATA |
| Imaging (X-ray, MRI/CT) | |
| Other | |
| Total | |

6. Reproductive Health

6.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|--|---------|------|-------|
| First antenatal visit < 1st trimester | NO DATA | | |
| First antenatal visit > 1st trimester | | | |
| Number of pregnant women at time of delivery who received: | | | |
| ≥ 4 ANC visits | | | |
| 2 TT doses | | | |
| anaemia screening | | | |

ANC Indicators

Proportion of first time antenatal visits made < 1st trimester

6.2 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|---------|------|-------|
| Women who attended 3 postnatal visits within 6 wks of delivery | NO DATA | | |

6.3 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|---|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | | | | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 0 | 0 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

| SGBV Indicators | |
|--|-----|
| Incidence of reported rape | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

Data Sources

1. Health Information System

Publication date:

20 December 2013

Contact Information

Erbil

Dr Chimmen Akawi, DOH Erbil,
Email: CHIMMENTYA@yahoo.com

Erbil

M. M. Taimur Hasan, UNHCR
Email: HASANMMT@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

