

1. Population

| Population | 0 to < 1 | | Total <1 | 1 to 4 | | Total 1 to 4 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total |
|--------------------|----------|----|----------|--------|-----|--------------|---------|-----|---------------|----------|-------|----------------|------|-----|------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | M | F | | |
| Refugee population | 95 | 95 | 190 | 378 | 379 | 757 | 838 | 678 | 1,516 | 1,505 | 1,257 | 2,762 | 427 | 388 | 815 | 6,040 |
| % by age group | | | 3% | | | 13% | | | 25% | | | 46% | | | 13% | 100% |

* estimates from registered population

2. Mortality

2.1 Mortality

| Record ONE cause of mortality per case | 0 to 4 | | Total <5 | U5 % mort | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | Crude % mort |
|---|--------|---|----------|-----------|---------|---|---------------|----------|---|----------------|------|---|------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| 1. Lower Respiratory Tract Infection (LRTI) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 2. Watery diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 3. Bloody diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 4. Cardiovascular disorders | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 5. Ischaemic heart disease | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 6. Injury | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 7. Cancer | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 8. Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 9. Tuberculosis (confirmed) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 10. Neonatal death | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Other | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Total | 0 | 0 | 0 | 100% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100% |
| % by age group | | | | | | | | | | | | | | 100% | |

| Mortality indicators | Male | Female | Total |
|--------------------------------|------|--------|-------|
| Crude Mortality Rate (CMR) | 0.0 | 0.0 | 0.0 |
| Under 5 Mortality Rate (U5MR) | 0.0 | 0.0 | 0.0 |
| Neonatal Mortality Rate (NNMR) | | | 0.0 |

2.2 Maternal death

| | |
|--|---|
| Number of maternal deaths reported | 0 |
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3. Morbidity

3.1 Staffing and service availability

| | |
|---|-------|
| Number of full-time clinicians | 2 |
| Average number of full days on which each health facility was | 7.0 |
| Total number of outpatient consultations | 1,744 |
| Health facility utilisation rate | 15.0 |
| Consultations per clinician per day | 125 |

3.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total <5 | U5 Incid** | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause | Crude Incid** |
|--|--------|-----|----------|------------|---------|-----|---------------|----------|-----|----------------|------|----|------------|-------------|------------|---------------|
| | M | F | | | M | F | | M | F | | M | F | | | | |
| Upper Respiratory Tract Infection (URTI) | 92 | 97 | 189 | 199.6 | 135 | 133 | 268 | 191 | 212 | 403 | 23 | 37 | 60 | 920 | 55% | 152.3 |
| Lower Respiratory Tract Infection (LRTI) | 9 | 10 | 19 | 20.1 | 10 | 1 | 11 | 8 | 7 | 15 | 6 | 7 | 13 | 58 | 3% | 9.6 |
| Watery diarrhoea* | 8 | 9 | 17 | 18.0 | 8 | 4 | 12 | 3 | 2 | 5 | 0 | 1 | 1 | 35 | 2% | 5.8 |
| Bloody diarrhoea* | 1 | 0 | 1 | 1.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% | 0.2 |
| Malaria (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Skin infection | 0 | 0 | 0 | 0.0 | 1 | 1 | 2 | 2 | 3 | 5 | 0 | 0 | 0 | 7 | 0% | 1.2 |
| Eye infection | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Ear infection | 1 | 1 | 2 | 2.1 | 4 | 0 | 4 | 0 | 8 | 8 | 0 | 0 | 0 | 14 | 1% | 2.3 |
| Dental Conditions | 1 | 1 | 2 | 2.1 | 2 | 3 | 5 | 19 | 28 | 47 | 2 | 1 | 3 | 57 | 3% | 9.4 |
| Intestinal Worms | 1 | 1 | 2 | 2.1 | 1 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 0% | 0.7 |
| Acute Jaundice Syndrome* | 0 | 0 | 0 | 0.0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% | 0.2 |
| Measles (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Meningitis (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Fever of unknown origin | 0 | 2 | 2 | 2.1 | 3 | 1 | 4 | 4 | 3 | 7 | 0 | 0 | 0 | 13 | 1% | 2.2 |
| Sexually Transmitted Infection (STI) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Urinary tract infection (UTI) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 9 | 9 | 0 | 0 | 0 | 9 | 1% | 1.5 |
| HIV/AIDS | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute malnutrition | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Leishmaniasis | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Other | 24 | 37 | 61 | 64.4 | 40 | 43 | 83 | 125 | 258 | 383 | 15 | 16 | 31 | 558 | 33% | 92.4 |
| Total | 137 | 158 | 295 | 311.5 | 205 | 187 | 392 | 352 | 530 | 882 | 46 | 62 | 108 | 1677 | 100% | 277.6 |
| % by age group | | | 18% | | | | 23% | | | 53% | | | 6% | 100% | | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **Incidence is cases per 1000 per week

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|--|--------|---|-----------|---------|---|---------------|----------|----|----------------|------|---|------------|-------------|------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| Diabetes | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 12 | 15 | 3 | 2 | 5 | 20 | 40% |
| Hypertension | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 7 | 9 | 0 | 2 | 2 | 11 | 22% |
| Asthma | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 0 | 4 | 3 | 0 | 3 | 7 | 14% |
| Ischaemic heart disease | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Cardiovascular disease (other) | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 4 | 6 | 2 | 2 | 4 | 10 | 20% |
| Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Liver disease | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Thyroid problems | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 1 | 1 | 2 | 4% |
| Cancer | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Total | 0 | 0 | 0 | 0 | 0 | 0 | 12 | 23 | 35 | 8 | 7 | 15 | 50 | 100% |
| % by age group | | | 0% | | | 0% | | | 70% | | | 30% | 100% | |

3.4 Mental health

| | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|---|----------|----------|-----------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Epilepsy / seizures | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 100% |
| 2. Alcohol and other substance abuse disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Psychotic disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4. Mental retardation / intellectual disability | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 5. Severe emotional disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 6. Medically unexplained somatic complaint | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Total | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 100% |
| % by age group | | | 0% | | | 0% | | | 100% | | | 0% | 100% | |

3.5 Injuries

| | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|---------------------------|----------|----------|------------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Injury (intentional) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 6% |
| 2. Injury (unintentional) | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 6% |
| 3. Burns | 3 | 4 | 7 | 1 | 0 | 1 | 3 | 3 | 6 | 0 | 0 | 0 | 14 | 88% |
| 4. Accident | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Total | 3 | 4 | 7 | 1 | 0 | 1 | 4 | 4 | 8 | 0 | 0 | 0 | 16 | 100% |
| % by age group | | | 44% | | | 6% | | | 50% | | | 0% | 100% | |

4. Referrals

4.1 Referrals - general

| | 0 to 4 | 5 to 14 | 15 to 49 | ≥ 50 | Grand total |
|-----------------------|----------|----------|----------|----------|-------------|
| Emergency | 0 | 0 | 0 | 0 | 0 |
| Elective | 0 | 0 | 0 | 0 | 0 |
| Other | 0 | 0 | 0 | 0 | 0 |
| Total | 0 | 0 | 0 | 0 | 0 |
| % by age group | | | | | 100% |

ND: no data submitted

4.2 Referrals - by specialty

| | 0 to 4 | 5 to 14 | 15 to 59 | ≥ 50 | Grand total |
|-----------------------|---------|---------|----------|------|-------------|
| Internal medicine | NO DATA | | | | |
| General Surgery | | | | | |
| Gynaecology | | | | | |
| Other | | | | | |
| Total | | | | | |
| % by age group | | | | | |

ND: no data submitted

5. Laboratory & other investigations

| | Total |
|-------------------------|---------|
| Haematology | NO DATA |
| Imaging (X-ray, MRI/CT) | |
| Other | |
| Total | |

6. Reproductive Health

6.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|--|---------|------|-------|
| First antenatal visit < 1st trimester | NO DATA | | |
| First antenatal visit > 1st trimester | | | |
| Number of pregnant women at time of delivery who received: | | | |
| ≥ 4 ANC visits | | | |
| 2 TT doses | | | |
| anaemia screening | | | |

ND: no data submitted

| | |
|--|--|
| ANC Indicators | |
| Proportion of first time antenatal visits made < 1st trimester | |

6.2 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|---------|------|-------|
| Women who attended 3 postnatal visits within 6 wks of delivery | NO DATA | | |

ND: no data submitted

6.3 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|---|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | | | | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 0 | 0 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

| | |
|--|-----|
| SGBV Indicators | |
| Incidence of reported rape | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

Data Sources

1. Health Information System

Publication date:

08 January 2014

Contact Information

Erbil
Dr Chimmen Akrawi, DOH Erbil

Email: chimmentya@yahoo.com

Erbil
M. M. Taimur Hasan, UNHCR

Email: HASANMMT@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

