




Refugee population

Total refugee population: **99,850** *

Demographic breakdown:



Key facts

Camp opened:
July 2012

* Source of population data:
WFP Distribution Figures

Indicators at a glance

| | | |
|---|------|---|
| 1 Health utilisation rate | 7.3 | ✓ |
| 2 Number of consultations per clinician per day | 28 | ✓ |
| 3 Hospitalisation rate | 3.4 | ⚠ |
| 4 Referral rate | | ? |
| 5 % deliveries attended by skilled personnel | 100% | ⚠ |
| 6 Crude Mortality Rate (CMR) (/1000/month) | 0.1 | ✓ |
| 7 Under-five Mortality Rate (U5MR) (/1000/month) | 0.2 | ✓ |
| 8 Neonatal Mortality Rate (NNMR) (/1000 livebirths) | 18.2 | ✗ |

Notes


This public health profile for Za'atri refugee camp is compiled from health facility data from the following partners:

- IMC Clinic
- JHAS Clinic
- Jordan Italian Field Hospital
- MDM Clinics 1 and 2
- Moroccan Field Hospital
- Saudi Clinic

Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Morbidity

Acute health conditions (%)




Top 5 acute health conditions


| Condition | % morbidity | # of cases |
|--|-------------|------------|
| 1 Upper Respiratory Tract Infection (URTI) | 29% | 2,283 |
| 2 Skin infection | 14% | 1,144 |
| 3 Influenza-like illness (ILI) | 13% | 1,023 |
| 4 Lower Respiratory Tract Infection (LRTI) | 7% | 347 |
| 5 Dental Conditions | 4% | 532 |
| - Other | 33% | 3,497 |

Breakdown of acute health conditions


by age



by gender



Chronic health conditions (%)




Top 5 chronic health conditions


| Condition | % morbidity | # of cases |
|--|-------------|------------|
| 1 Hypertension | 11% | 543 |
| 2 Diabetes | 9% | 446 |
| 3 Asthma | 8% | 388 |
| 4 Ischaemic heart disease | 4% | 174 |
| 5 Chronic obstructive pulmonary disease (COPD) | 3% | 135 |
| - Other | 66% | 3,224 |

Breakdown of chronic health conditions

by age



by gender



UNHCR gratefully acknowledges the support of the following partners:



Public Health

Health Impact

| | Indicator | Standard | |
|---|-----------|----------|---|
| Crude Mortality Rate (CMR) (/1000/month) | 0.1 | < 1.5 | ✓ |
| Under-five Mortality Rate (U5MR) (/1000/month) | 0.2 | < 3.0 | ✓ |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | 18.2 | < 40 | ✓ |

Access and utilization

| | | | | |
|--|--------|-----------|--------------|---|
| No. of health facilities | 7 | 1 : 16642 | 1 : < 10,000 | ✗ |
| Total number of consultations | 13,947 | | | |
| Health utilisation rate | 7.3 | 1 - 4 | | ✗ |
| No. of consultations per trained clinician per day | 28 | < 50 | | ✓ |

Outbreak Alert and Response

| | Indicator | Standard |
|---|-----------|----------|
| Number of outbreaks reported | 0 | |
| % of outbreaks investigated within 48 hours | | 100% |

Laboratory


| | |
|------------------------------|-----|
| Number of laboratory samples | 637 |
|------------------------------|-----|

Immunization


| | | |
|--------------------------|-------|---|
| Measles vaccination rate | > 95% | ? |
| Full vaccination rate | > 95% | ? |
| Program drop-out rate | < 10% | ? |
| Vaccine wastage rate | < 8% | ? |

Breakdown of mental health conditions


By cause



By age




By gender




Breakdown of injuries


By cause



By age



By gender



In-patient and referral

In-patient department

| | Indicator | Standard | |
|--|-----------|----------|---|
| Hospitalisation rate (/1000/month) | 3.4 | 50 - 150 | ✗ |
| Bed occupancy rate | 44% | > 75% | ✗ |
| Mean length of stay for authorised discharges (days) | 1.7 | | |

| | Indicator | Standard |
|---|-----------|----------|
| Overall case fatality rate | 0% | |
| Proportion of deaths within 24 hours of admission | 0% | |

Referral

| | |
|---------------------|---|
| Total referral rate | ? |
|---------------------|---|

Reproductive Health

Antenatal care

| | Indicator | Standard | |
|---|-----------|----------|---|
| Coverage of antenatal care (4 or more visits) | 15% | ≥ 90% | ✓ |
| Coverage of TT vaccination | 11% | ≥ 90% | ✗ |
| Coverage of anaemia screening in pregnancy | 56% | ≥ 90% | ✓ |

Delivery care

| | | | |
|---|------|---------|---|
| Proportion of deliveries attended by skilled personnel | 100% | ≥ 90% | ⚠ |
| Proportion of deliveries performed by caesarean section | 9% | 5 - 15% | ✓ |
| Proportion of low birth weight deliveries (<2500g) | 0% | < 15% | ✓ |

Maternal mortality

| | | | |
|---|---|------|--|
| Number of maternal deaths reported | 0 | | |
| % of maternal deaths investigated within 48 hours | | 100% | |

Postnatal care

| | | | |
|----------------------------|------|-------|---|
| Coverage of postnatal care | 202% | ≥ 90% | ⚠ |
|----------------------------|------|-------|---|

Family Planning

| | | | |
|-------------------------------|----|--|---|
| Contraceptive prevalence rate | 2% | | ✗ |
|-------------------------------|----|--|---|

Sexual and Gender-based Violence

| | | | |
|---|---|------|---|
| Incidence of rape (reported cases/10,000/year)* | 0 | | ✗ |
| % of rape survivors who received PEP** < 72 hours | | 100% | ✗ |
| % of rape survivors who received ECP*** < 120 hours | | 100% | ✗ |
| % of rape survivors who received ST**** prophylaxis < 2 wks | | 100% | ✗ |

*Only includes rapes reported at health facilities **Post-exposure prophylaxis ***Emergency contraceptive pill ****Sexually transmitted infection

