


Refugee population

Total refugee population:

3,184 *

Demographic breakdown:



Basirma camp

Key facts

Camp opened:
August 2013Planned capacity:
4,580Area:
0.15 Km sqrPopulation:
Syrian refugees, majority
from Qamishli and Hasak* Source of population data:
Estimates from UNHCR
registration

Indicators at a glance

1	Health utilisation rate	12.6	1 - 4
2	Number of consultations per clinician per day	55	< 50
3	Proportion of first time antenatal visits made < 1st trimester	N/A	≥ 90%
4	Measles immunisation coverage	26%	> 90%
5	Polio III immunisation coverage	71%	> 90%
6	Crude Mortality Rate (CMR) (/1000/month)	0.0	< 1.5
7	Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 3.0
8	Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0.0	< 40

Notes

HIGHLIGHTS

Health utilisation rate and consultations per clinician per day elevated.
Upper respiratory tract infections remains the major cause of morbidity.

This public health profile for the refugee camp is compiled from health data obtained from reporting agencies. Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Agencies reporting:


DOH Erbil

Alerts


Incidence of watery diarrhoea above threshold

Morbidity

Acute health conditions (%)



Chronic health conditions (%)



Top 5 acute health conditions


	% morbidity	# of cases
1 Upper Respiratory Tract Infection (URTI)	49%	351
2 Dental Conditions	7%	47
3 Skin infection	5%	38
4 Eye infection	3%	25
5 Watery diarrhoea*	3%	24
- Other	33%	235

Top 5 chronic health conditions


	% morbidity	# of cases
1 Hypertension	52%	16
2 Diabetes	39%	12
3 Asthma	3%	1
4 Cardiovascular disease (other)	3%	1
5 Ischaemic heart disease	0%	0
- Other	0.032258	1

Breakdown of acute health conditions

by age




by gender




Breakdown of chronic health conditions

by age




by gender




Trends since the January 2014


Number of consultations




Weekly health utilisation rates




Watery diarrhoea




Bloody diarrhoea




Acute Jaundice Syndrome




Suspected measles




Suspected Meningitis




Acute flaccid paralysis



Upper respiratory tract infections



Lower respiratory tract infections



Calculation of disease alert thresholds:

Watery diarrhoea = 1.5 x baseline, Bloody diarrhoea = 1.5 x baseline, Acute Jaundice Syndrome = 1.5 x baseline,
Meningitis = 5 cases, Acute Flaccid Paralysis = 1 case, Measles = 1 case,
baseline = average incidence or number of cases over the previous 3 weeks

Upper respiratory tract infections = 1.5 x baseline,

Public Health

Health Impact

Indicator	Standard
Crude Mortality Rate (CMR) (/1000/month)	0.0 < 1.5
Under-five Mortality Rate (U5MR) (/1000/month)	0.0 < 3.0
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0.0 < 40

Access and utilization

No. of health facilities	1.0	1 : < 10,000
Total number of consultations	774	
Health utilisation rate	12.6	1 - 4
No. of consultations per trained clinician per day	55	< 50

Reproductive health

Proportion of first time antenatal visits made < 1st trimester	N/A	≥ 90%
--	-----	-------

Outbreak Alert and Response

Indicator	Standard
Number of outbreaks reported	0
% of outbreaks investigated within 48 hours	N/A 100%

Routine immunization

BCG immunisation coverage	52%	> 90%
DPT 1 immunisation coverage	41%	> 90%
Measles immunisation coverage	26%	> 90%
Polio III immunisation coverage	71%	> 90%
MMR immunisation coverage	78%	> 90%
Vitamin A coverage	58%	> 90%

Sexual and gender-based violence

Incidence of reported rape	0.0
----------------------------	-----

Top 3 Mental health conditions


	% morbidity	# of cases
1 Epilepsy / seizures	100%	2
2 Alcohol and other substance abuse disorder	0%	0
3 Psychotic disorder	0%	0
- Other	0%	0

Top 3 injuries


	% morbidity	# of cases
1 Injury (unintentional)	71%	15
2 Burns	29%	6
3 Injury (intentional)	0%	0
- Other	0%	0

Breakdown of mental health conditions


By cause



By age




By gender




Breakdown of injuries


By cause



By age



By gender



We gratefully acknowledge the support of the following agencies:



View interactive maps and statistics online
<http://twine.unhcr.org>

