



### 3.4 Mental health

	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
1. Epilepsy / seizures	0	0	0	0	0	0	2	2	4	0	0	0	4	6%
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
3. Psychotic disorder	0	0	0	0	0	0	5	3	8	0	0	0	8	11%
4. Mental retardation / intellectual disability	0	0	0	1	0	1	1	0	1	0	0	0	2	3%
5. Severe emotional disorder	0	0	0	1	0	1	5	4	9	0	0	0	10	14%
6. Medically unexplained somatic complaint	0	0	0	1	0	1	1	1	2	0	0	0	3	4%
Other	3	2	5	0	3	3	7	25	32	0	3	3	43	61%
<b>Total</b>	<b>3</b>	<b>2</b>	<b>5</b>	<b>3</b>	<b>3</b>	<b>6</b>	<b>21</b>	<b>35</b>	<b>56</b>	<b>0</b>	<b>3</b>	<b>3</b>	<b>70</b>	<b>100%</b>
% by age group			7%			9%			80%			4%	100%	

### 3.5 Injuries

	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
1. Injury (intentional)	1	0	1	3	2	5	2	1	3	0	0	0	9	6%
2. Injury (unintentional)	9	12	21	13	2	15	43	6	49	3	1	4	89	63%
3. Burns	5	4	9	1	0	1	0	4	4	0	0	0	14	10%
4. Accident	4	0	4	1	0	1	15	0	15	0	0	0	20	14%
Other	2	1	3	1	1	2	4	1	5	0	0	0	10	7%
<b>Total</b>	<b>21</b>	<b>17</b>	<b>38</b>	<b>19</b>	<b>5</b>	<b>24</b>	<b>64</b>	<b>12</b>	<b>76</b>	<b>3</b>	<b>1</b>	<b>4</b>	<b>142</b>	<b>100%</b>
% by age group			27%			17%			54%			3%	100%	

## 4. Referrals

### 4.1 Referrals - general

	0 to 4	5 to 14	15 to 49	≥ 50	Grand total
Emergency	8	6	15	4	33
Elective	18	24	51	17	110
Other	0	0	0	0	0
<b>Total</b>	<b>26</b>	<b>30</b>	<b>66</b>	<b>21</b>	<b>143</b>
% by age group	18%	21%	46%	15%	100%

ND: no data submitted

## 5. Laboratory & other investigations

	Total
Haematology	0
Imaging (X-ray, MRI/CT)	0
Other	0
<b>Total</b>	<b>6</b>

## 6. EPI and Vitamin A

	<1	1 to <5	Total
BCG	0	0	0
Hepatitis B birth dose	0	0	0
DPT I/HiB I/Hep B I* (Pentavalent I)	0	0	0
DPT II/HiB II* (Tetravalent)	0	0	0
DPT III/HiB III/Hep B II* (Pentavalent II)	0	0	0
Polio 0	0	0	0
Polio I	0	0	0
Polio II	0	0	0
Polio III	0	0	0
Measles	0	0	0
MMR	0	0	0
Rotavirus 1	0	0	0
Rotavirus 2	0	0	0
Rotavirus 3	0	0	0
Vitamin A	0	0	0
Additional vaccines	0	0	0

EPI Indicators (since December 2013)	
Total vaccinated	0
a. Program dropout rate*	-22%
b. Immunisation coverage	
BCG	52%
Hepatitis B birth dose	52%
DPT I	86%
DPT III	105%
Polio I	86%
Polio III	105%
Measles	107%
MMR**	129%
Rotavirus I	86%
Rotavirus III	105%
c. Vitamin A coverage	100%

\*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines  
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%  
\*\*For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

## 7. Reproductive Health

### 7.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester			
First antenatal visit > 1st trimester			
Number of pregnant women at time of delivery who received:	<b>NO DATA</b>		
≥ 4 ANC visits			
2 TT doses			
anaemia screening			

ND: no data submitted

ANC Indicators	
Proportion of first time antenatal visits made < 1st trimester	

### 7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery			<b>NO DATA</b>

ND: no data submitted

### 7.3 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No of rape survivors given PEP within 72 hours	0	0	0	0	0
No of female rape survivors given ECP within 120 hours					0
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators	
Incidence of reported rape	<b>0.0</b>
Proportion of rape survivors who receive PEP < 72 hrs	
Proportion of female rape survivors who receive ECP < 120 hrs	
Proportion of rape survivors who receive STI presumptive treatment < 2 wks	

Data Sources: 1. Health Information System

Publication date: 29 April 2014

Contact Information: Erbil, Dr Raad Najmaddin, DOH Erbil

Email: RAAD976@yahoo.com

Erbil, Sandra Harlass, UNHCR

Email: HARLASS@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

