

1. Population

| Population | 0 to < 1 | | Total <1 | 1 to 4 | | Total 1 to 4 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total |
|--------------------|----------|-------|----------|--------|-------|--------------|---------|--------|---------------|----------|--------|----------------|------|-------|------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | M | F | | |
| Refugee population | 1,674 | 1,594 | 3,267 | 6,121 | 5,665 | 11,787 | 14,985 | 14,265 | 29,250 | 16,319 | 16,878 | 33,197 | 909 | 1,283 | 2,192 | 79,693 |
| % by age group | | | 4% | | | 15% | | | 37% | | | 42% | | | 3% | 100% |

2. Mortality

2.1 Mortality

| Record ONE cause of mortality per case | 0 to 4 | | Total <5 | US % mort | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | Crude % mort |
|--|--------|----|----------|-----------|---------|---|---------------|----------|---|----------------|------|----|------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| 1. Lower Respiratory Tract Infection (LRTI) | 0 | 1 | 1 | 3% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1% |
| 2. Watery diarrhoea | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Bloody diarrhoea | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4. Ischaemic heart disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 3 | 0 | 3 | 7 | 2 | 9 | 12 | 12% |
| 5. Cardiovascular disorders (other) | 1 | 0 | 1 | 3% | 0 | 0 | 0 | 3 | 2 | 5 | 3 | 6 | 9 | 15 | 15% |
| 6. Cerebrovascular disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 3 | 3 | 4 | 4 | 8 | 11 | 11% |
| 7. Chronic kidney disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 1 | 1 | 2 | 1 | 3 | 4 | 4% |
| 8. Injury | 0 | 0 | 0 | 0% | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 | 2 | 2% |
| 9. Cancer | 2 | 0 | 2 | 7% | 1 | 0 | 1 | 5 | 1 | 6 | 6 | 4 | 10 | 19 | 19% |
| 10. Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0 | 3 | 3 | 3% |
| 11. Tuberculosis (confirmed) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 12. Neonatal death | 13 | 10 | 23 | 77% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 23 | 23% |
| 13. Other | 1 | 2 | 3 | 10% | 1 | 2 | 3 | 0 | 2 | 2 | 1 | 2 | 3 | 11 | 11% |
| Total | 17 | 13 | 30 | 100% | 3 | 2 | 5 | 11 | 9 | 20 | 27 | 19 | 46 | 101 | 100% |
| % by age group | | | 30% | | | | 5% | | | 20% | | | 46% | 100% | |

| Mortality indicators | Male | Female | Total |
|---|------|--------|-------|
| Crude Mortality Rate (CMR) (/1000/month) | 0.2 | 0.2 | 0.2 |
| Under 5 Mortality Rate (USMR) (/1000/month) | 0.4 | 0.3 | 0.3 |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | | | 13.3 |

2.2 Maternal death

| | |
|--|------|
| Number of maternal deaths reported | 1 |
| Number of maternal deaths investigated | 1 |
| Proportion of maternal deaths investigated | 100% |

3. Morbidity

3.1 Staffing and service availability

| | |
|---|---------|
| Number of full-time clinicians | 57.5 |
| Average number of full days on which each health facility was functioning | 5.5 |
| Total number of outpatient consultations | 301,692 |
| Health facility utilisation rate | 7.6 |
| Consultations per clinician per day | 37 |

3.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total <5 | US Incid | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause | Crude Incide |
|--|--------|--------|----------|----------|---------|--------|---------------|----------|--------|----------------|-------|-------|------------|-------------|------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | | |
| Upper Respiratory Tract Infection (URTI) | 10,296 | 9,556 | 19,852 | 1318.7 | 7,670 | 8,571 | 16,241 | 8,631 | 9,185 | 17,816 | 1,452 | 1,227 | 2,679 | 56,588 | 31% | 710.1 |
| Influenza-like illness (ILI) | 2,590 | 2,528 | 5,118 | 340.0 | 2,699 | 2,903 | 5,602 | 3,365 | 3,758 | 7,123 | 633 | 561 | 1,194 | 19,037 | 10% | 238.9 |
| Lower Respiratory Tract Infection (LRTI) | 945 | 838 | 1,783 | 118.4 | 1,092 | 986 | 2,078 | 1,488 | 1,417 | 2,905 | 490 | 430 | 920 | 7,686 | 4% | 96.4 |
| Watery diarrhoea* | 442 | 373 | 815 | 54.1 | 181 | 168 | 349 | 134 | 127 | 261 | 37 | 31 | 68 | 1,493 | 1% | 18.7 |
| Bloody diarrhoea* | 4 | 3 | 7 | 0.5 | 2 | 2 | 4 | 2 | 2 | 4 | 0 | 0 | 0 | 15 | 0% | 0.2 |
| Skin infection | 1,321 | 1,345 | 2,666 | 177.1 | 1,420 | 1,705 | 3,125 | 1,811 | 1,864 | 3,675 | 657 | 567 | 1,224 | 10,690 | 6% | 134.1 |
| Eye infection | 649 | 547 | 1,196 | 79.4 | 510 | 615 | 1,125 | 831 | 989 | 1,820 | 616 | 569 | 1,185 | 5,326 | 3% | 66.8 |
| Ear infection | 877 | 1,043 | 1,920 | 127.5 | 1,061 | 1,171 | 2,232 | 1,537 | 1,552 | 3,089 | 691 | 575 | 1,266 | 8,507 | 5% | 106.7 |
| Dental Conditions | 420 | 549 | 969 | 64.4 | 2,275 | 2,756 | 5,031 | 5,853 | 6,587 | 12,440 | 823 | 661 | 1,484 | 19,924 | 11% | 250.0 |
| Intestinal Worms | 413 | 462 | 875 | 58.1 | 689 | 626 | 1,315 | 158 | 164 | 322 | 6 | 2 | 8 | 2,520 | 1% | 31.6 |
| Acute Jaundice Syndrome* | 1 | 0 | 1 | 0.1 | 0 | 0 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 3 | 0% | 0.0 |
| Measles (suspected)* | 0 | 2 | 2 | 0.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% | 0.0 |
| Meningitis (suspected)* | 3 | 2 | 5 | 0.3 | 3 | 3 | 6 | 1 | 0 | 1 | 0 | 0 | 0 | 12 | 0% | 0.2 |
| Acute Flaccid Paralysis* | 1 | 1 | 2 | 0.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% | 0.0 |
| Leishmaniasis | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Fever of unknown origin | 7 | 14 | 21 | 1.4 | 1 | 0 | 1 | 1 | 1 | 2 | 0 | 0 | 0 | 24 | 0% | 0.3 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 2 | 2 | 2 | 3 | 5 | 1 | 2 | 3 | 10 | 0% | 0.1 |
| Other | 4,776 | 4,528 | 9,304 | 618.0 | 5,986 | 6,462 | 12,448 | 12,614 | 13,170 | 25,784 | 2,822 | 2,764 | 5,586 | 53,122 | 29% | 666.6 |
| Total | 22,745 | 21,791 | 44,536 | 2958.4 | 23,589 | 25,970 | 49,559 | 36,430 | 38,819 | 75,249 | 8,228 | 7,389 | 15,617 | 184,961 | 100% | 2320.9 |
| % by age group | 24% | | 24% | | 27% | | 27% | | 41% | | 41% | | 8% | | 100% | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|--|--------|-------|-----------|---------|-------|---------------|----------|--------|----------------|-------|-------|------------|-------------|------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| Diabetes | 2 | 9 | 11 | 113 | 120 | 233 | 1,960 | 2,693 | 4,653 | 1,482 | 1,564 | 3,046 | 7,943 | 16% |
| Hypertension | 1 | 0 | 1 | 7 | 67 | 74 | 2,719 | 3,450 | 6,169 | 1,869 | 2,036 | 3,905 | 10,149 | 20% |
| Asthma | 1,072 | 899 | 1,971 | 527 | 544 | 1,071 | 1,228 | 1,208 | 2,436 | 776 | 642 | 1,418 | 6,896 | 14% |
| Ischaemic heart disease | 0 | 3 | 3 | 4 | 5 | 9 | 456 | 500 | 956 | 656 | 534 | 1,190 | 2,158 | 4% |
| Cardiovascular disease (other) | 8 | 5 | 13 | 34 | 22 | 56 | 731 | 792 | 1,523 | 594 | 504 | 1,098 | 2,690 | 5% |
| Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0 | 0 | 0 | 571 | 534 | 1,105 | 735 | 605 | 1,340 | 2,445 | 5% |
| Liver disease | 0 | 0 | 0 | 5 | 3 | 8 | 64 | 62 | 126 | 25 | 10 | 35 | 169 | 0% |
| Thyroid problems | 4 | 0 | 4 | 45 | 152 | 197 | 869 | 1,635 | 2,504 | 665 | 812 | 1,477 | 4,182 | 8% |
| Cancer | 0 | 0 | 0 | 0 | 0 | 0 | 9 | 13 | 22 | 6 | 1 | 7 | 29 | 0% |
| Other | 445 | 452 | 897 | 872 | 930 | 1,802 | 3,593 | 4,098 | 7,691 | 1,554 | 1,359 | 2,913 | 13,303 | 27% |
| Total | 1,532 | 1,368 | 2,900 | 1,607 | 1,843 | 3,450 | 12,200 | 14,985 | 27,185 | 8,362 | 8,067 | 16,429 | 49,964 | 100% |
| % by age group | | | 6% | | | 7% | | | 54% | | | 33% | 100% | |

3.4 Mental health

| | | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|--------------------------------|---|--------|----|-----------|---------|-----|---------------|----------|-----|----------------|------|----|------------|-------------|------------|
| | | M | F | | M | F | | M | F | | M | F | | | |
| 1. | Epilepsy / seizures | 52 | 34 | 86 | 219 | 119 | 338 | 295 | 194 | 489 | 13 | 9 | 22 | 935 | 25% |
| 2. | Alcohol and other substance use disorder | 0 | 0 | 0 | 0 | 0 | 0 | 13 | 3 | 16 | 2 | 1 | 3 | 19 | 1% |
| 3. | Psychotic disorder (including mania) | 0 | 1 | 1 | 6 | 5 | 11 | 452 | 184 | 636 | 10 | 3 | 13 | 661 | 18% |
| 4. | Intellectual disability | 6 | 15 | 21 | 129 | 128 | 257 | 92 | 38 | 130 | 0 | 1 | 1 | 409 | 11% |
| 5. | Severe emotional disorder, including moderate-severe depression | 0 | 0 | 0 | 12 | 31 | 43 | 582 | 373 | 955 | 19 | 20 | 39 | 1,037 | 28% |
| 6. | Medically unexplained somatic complaint | 0 | 0 | 0 | 23 | 24 | 47 | 31 | 18 | 49 | 0 | 0 | 0 | 96 | 3% |
| Other psychological complaints | | 2 | 12 | 14 | 112 | 171 | 283 | 166 | 72 | 238 | 29 | 12 | 41 | 576 | 15% |
| Total | | 60 | 62 | 122 | 501 | 478 | 979 | 1,631 | 882 | 2,513 | 73 | 46 | 119 | 3,733 | 100% |
| % by age group | | | | 3% | | | 26% | | | 67% | | | 3% | 100% | |

3.5 Injuries

| | | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|----------------|--|--------|-----|-----------|---------|-------|---------------|----------|-------|----------------|------|-----|------------|-------------|------------|
| | | M | F | | M | F | | M | F | | M | F | | | |
| 1. | Bites (all causes) | 4 | 4 | 8 | 3 | 5 | 8 | 8 | 1 | 9 | 0 | 0 | 0 | 25 | 0% |
| 2. | Injury (war related) | 1 | 6 | 7 | 29 | 28 | 57 | 18 | 24 | 42 | 0 | 0 | 0 | 106 | 1% |
| 3. | Injury (Burns) | 320 | 348 | 668 | 209 | 366 | 575 | 132 | 273 | 405 | 13 | 16 | 29 | 1,677 | 15% |
| 4. | Assault (with weapon non-war related) | 0 | 0 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% |
| Other | | 668 | 629 | 1,297 | 1,354 | 1,078 | 2,432 | 1,919 | 1,663 | 3,582 | 974 | 897 | 1,871 | 9,182 | 84% |
| Total | | 993 | 987 | 1,980 | 1,597 | 1,477 | 3,074 | 2,077 | 1,961 | 4,038 | 987 | 913 | 1,900 | 10,992 | 100% |
| % by age group | | | | 18% | | | 28% | | | 37% | | | 17% | 100% | |

3.6 Reproductive Tract Infections

| | | < 18 | | Total < 18 | ≥ 18 | | Total ≥ 18 | Grand total | % by cause |
|----------------|----------------------|------|-----|------------|------|-------|------------|-------------|------------|
| | | M | F | | M | F | | | |
| 1. | Vaginal discharge | 0 | 309 | 309 | 0 | 2,229 | 2,229 | 2,538 | 75% |
| 2. | Urethral discharge | 0 | 0 | 0 | 2 | 0 | 2 | 2 | 0% |
| 3. | Genital ulcer | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. | Lower abdominal pain | 0 | 112 | 112 | 0 | 740 | 740 | 852 | 25% |
| Other | | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 0% |
| Total | | 0 | 421 | 421 | 2 | 2,970 | 2,972 | 3,393 | 100% |
| % by age group | | | | 12% | | | 88% | 100% | |

4. In-patient Department

4.1 In-patient Department Activities

| | | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total |
|---|--------------|--------|---|-----------|---------|-----|---------------|----------|-------|----------------|------|---|------------|-------------|
| | | M | F | | M | F | | M | F | | M | F | | |
| Number at beginning of reporting period | | 0 | 0 | 0 | 2 | 1 | 3 | 2 | 6 | 8 | 0 | 0 | 0 | 11 |
| Total admissions | | 17 | 5 | 22 | 18 | 183 | 201 | 195 | 1,648 | 1,843 | 15 | 6 | 21 | 2,087 |
| Number of exits | authorized | 17 | 5 | 22 | 17 | 126 | 143 | 174 | 1,169 | 1,343 | 15 | 6 | 21 | 1,529 |
| | unauthorized | 0 | 0 | 0 | 0 | 3 | 3 | 0 | 31 | 31 | 0 | 0 | 0 | 34 |
| | death | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | referral | 0 | 0 | 0 | 1 | 53 | 54 | 14 | 450 | 464 | 0 | 0 | 0 | 518 |
| Total exits | | 17 | 5 | 22 | 18 | 182 | 200 | 188 | 1,650 | 1,838 | 15 | 6 | 21 | 2,081 |
| Deaths within 24 hours | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| | |
|---|--------|
| Number of days in reporting period | 182 |
| Number of beds | 38 |
| Sum number of days stay for authorised discharges | 1672.2 |
| Proportion of < 5 IPD deaths within 24 hours | |
| Average length of stay (days) | 1.1 |
| Bed occupancy | 33% |
| Hospitalisation rate (/1000/month) | 4.4 |

4.2 In-patient Admissions and Deaths

| | | 0 to 4 | | US CFR | 5 to 17 | | 5-17 CFR | 18 to 59 | | 18-59 CFR | > 60 | | Total adm | Total deaths | Crude CFR |
|-------|--------------------------|--------|--------|--------|---------|--------|----------|----------|--------|-----------|------|--------|-----------|--------------|-----------|
| | | Adm | Deaths | | Adm | Deaths | | Adm | Deaths | | Adm | Deaths | | | |
| 1. | Paediatric medicine | 21 | 0 | 0% | 5 | 0 | 0% | 1 | 0 | 0% | 1 | 0 | 28 | 0 | 0% |
| 2. | General Surgery | 1 | 0 | 0% | 10 | 0 | 0% | 169 | 0 | 0% | 7 | 0 | 187 | 0 | 0% |
| 3. | Injury (non-war related) | 0 | 0 | | 0 | 0 | | 5 | 0 | 0% | 1 | 0 | 6 | 0 | 0% |
| 4. | Internal medicine | 0 | 0 | | 15 | 0 | 0% | 75 | 0 | 0% | 12 | 0 | 102 | 0 | 0% |
| 5. | Obstetrics | 0 | 0 | | 166 | 0 | 0% | 1511 | 0 | 0% | 0 | 0 | 1,677 | 0 | 0% |
| 6. | Gynaecology | 0 | 0 | | 3 | 0 | 0% | 48 | 0 | 0% | 0 | 0 | 51 | 0 | 0% |
| 7. | Other | 0 | 0 | | 1 | 0 | 0% | 35 | 0 | 0% | 0 | 0 | 36 | 0 | 0% |
| Total | | 22 | 0 | 0% | 200 | 0 | 0% | 1844 | 0 | 0% | 21 | 0 | 2,087 | 0 | 0% |

Adm. = Admission; CFR = Case Fatality Rate

5. Referral

5.1 Referrals by location

| | | 0 to 4 | Total < 5 | 5 to 17 | Total 5 to 17 | 18 to 59 | Total 18 to 59 | > 60 | Total | Grand total |
|--|-----------------------------|--------|-----------|---------|---------------|----------|----------------|------|-------|-------------|
| 1. | MOH Hospital | 41 | 41 | 49 | 49 | 195 | 195 | 60 | 60 | 345 |
| 2. | Jordan University Hospital | 124 | 124 | 95 | 95 | 196 | 196 | 22 | 22 | 437 |
| 3. | Private affiliated Hospital | 578 | 578 | 560 | 560 | 1,836 | 1,836 | 256 | 256 | 3,230 |
| 4. | Royal Medical Services | 5 | 5 | 0 | 0 | 6 | 6 | 0 | 0 | 11 |
| 5. | Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total No of referred cases out side the Camp | | 4,023 | | | | | | | | |

5.2 Referrals by service

| | | 0 to 4 | Total < 5 | 5 to 17 | Total 5 to 17 | 18 to 59 | Total 18 to 59 | > 60 | Total | Grand total |
|----|-------------------|--------|-----------|---------|---------------|----------|----------------|------|-------|-------------|
| 1. | Paediatrics | 426 | 426 | 0 | 0 | 0 | 0 | 0 | 0 | 426 |
| 2. | Internal medicine | 129 | 129 | 278 | 278 | 987 | 987 | 186 | 186 | 1,580 |
| 3. | Obstetric | 0 | 0 | 57 | 57 | 495 | 495 | 0 | 0 | 552 |
| 4. | Gynaecological | 0 | 0 | 128 | 128 | 206 | 206 | 0 | 0 | 334 |
| 5. | Surgical | 179 | 179 | 204 | 204 | 462 | 462 | 120 | 120 | 965 |
| 6 | Other | 14 | 14 | 37 | 37 | 84 | 84 | 31 | 31 | 166 |

6. Laboratory

| | Total |
|--------------------------------------|---------|
| Number of laboratory tests conducted | 180,458 |

8. Reproductive Health

8.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|---------------------------------------|------|-------|-------|
| First antenatal visit < 1st trimester | 555 | 1,853 | 2,408 |
| First antenatal visit > 1st trimester | 116 | 814 | 930 |

| ANC Indicators | |
|--|-----|
| Proportion of first time antenatal visits made < 1st trimester | 72% |

8.2 Antenatal care in facility

| | | | | |
|--|-------------------|-----|------|-------|
| Number of pregnant women at time of delivery who received: | ≥ 4 ANC visits | 149 | 1125 | 1,274 |
| | 2 TT doses | 142 | 1136 | 1,278 |
| | anaemia screening | 170 | 1443 | 1,613 |

| | |
|--|-----|
| Coverage of complete antenatal care | 74% |
| Coverage of antenatal tetanus immunisation | 74% |
| Coverage of anaemia screening in pregnancy | 94% |

8.3 Delivery care

| | < 18 | ≥ 18 | Total |
|---|------|------|-------|
| Live births | 174 | 1549 | 1,723 |
| Still births | 1 | 4 | 5 |
| Low birth weight (<2500 g) | 1 | 16 | 17 |
| Attended by skilled health worker | 174 | 1549 | 1,723 |
| Number of obstetric complications treated | 0 | 14 | 14 |
| Number of caesarean sections performed | 49 | 499 | 548 |

| Delivery Indicators | |
|--|------|
| Crude birth rate (/1000 population/ month) | 3.6 |
| Proportion of births attended by skilled health worker (excl. TBA) | 100% |
| Proportion of low birth weight deliveries | 1% |
| Proportion of all births performed by caesarean section | 32% |
| Proportion of obsetric complications treated among girls < 18 | 0% |

8.4 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|-------|
| At least three PNC visits within 6 weeks | 176 | 884 | 1,060 |

| Postnatal Indicators | |
|-------------------------------------|-----|
| Coverage of complete postnatal care | 62% |

8.5 Family planning

| | New | | Repeat | | Total |
|-----------------------------|------|-------|--------|-------|-------|
| | < 18 | ≥ 18 | < 18 | ≥ 18 | |
| Oral contraceptives | 95 | 1,023 | 135 | 3,198 | 4,451 |
| Emergency Contaceptive Pill | 0 | 0 | 0 | 0 | 0 |
| Injectable (Depo-Provera) | 5 | 78 | 15 | 293 | 391 |
| Implantable (Norplant) | 0 | 0 | 0 | 0 | 0 |
| Intra-Uterine Device (IUD) | 11 | 343 | 0 | 60 | 414 |
| Condom (Male) | 55 | 691 | 97 | 2,303 | 3,146 |
| Other | 36 | 133 | 28 | 192 | 389 |

| | < 18 | ≥ 18 |
|-----------------------------|------|-------|
| Family Planning Counseling* | 901 | 3,256 |

8.6 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|----|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | 0 | | 0 | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 1 | 5 | 0 | 49 | 55 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

| SGBV Indicators | |
|--|-----|
| Incidence of reported rape (/10,000 population/year) | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

| | | | |
|---------------------|------------------------------|--------------------------|------------|
| Data Sources | 1. Health Information System | Publication date: | 26/07/2017 |
|---------------------|------------------------------|--------------------------|------------|

| | | |
|----------------------------|--|---|
| Contact Information | Za'atri Asaad Kadhum, UNHCR Email: kadhum@unhcr.org | Amman Adam Musa Khalifa, UNHCR Email: khalifaa@unhcr.org |
|----------------------------|--|---|

UNHCR gratefully acknowledges the support of the following partners:

