Zaatari Camp | Jordan

Refugee population

Total refugee population: **77,089**

Demographic breakdown:

Age group	Male	Female
> 60	1%	2%
18 - 59	21%	21%
5 - 17	19%	18%
0 - 4	10%	9%

Male | % | Female

Key facts

Camp opened:
July 2012

Indicators at a glance

#	Indicator	Value	
1	Health utilisation rate	3.7	√
2	Number of consultations per clinician per day	25	√
3	Referral rate (/1,000/month)	4.6	
4	Crude Mortality Rate (CMR) (/1000/month)	0.1	√
5	Under-five Mortality Rate (U5MR) (/1000/month)	0.3	√
6	Neonatal Mortality Rate (NNMR) (/1000 livebirths)	13.2	√

Notes

This public health profile for Za'atri refugee camp is compiled from health facility data from the following partners:

- IMC Clinic
- IMC ER Center
- JHAS Clinic
- Moroccan Field Hospital
- Qatari Clinic
- Saudi Clinic
- IRC Clinic
- AMR Clinic 2 (Maternity and Childhood center)
- SAMS Clinic

Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Morbidity

Acute health conditions (%)

- Upper Respiratory Tract Infection (URTI): 21%
- Influenza-like illness (ILI): 11%
- Dental Conditions: 8%
- Skin infection: 6%
- Ear infection: 5%
- Other: 49%

Chronic health conditions (%)

- Hypertension: 26%
- Diabetes: 22%
- Asthma: 15%
- Thyroid problems: 8%
- Ischaemic heart disease: 3%
- Other: 26%

Top 5 acute health conditions

		% morbidity	# of cases
1	Upper Respiratory Tract Infection (URTI)	22%	11,434
2	Influenza-like illness (ILI)	11%	5,612
3	Dental Conditions	8%	4,122
4	Skin infection	6%	3,378
5	Ear infection	5%	2,718
-	Other	49%	25,880

Top 5 chronic health conditions

		% morbidity	# of cases
1	Hypertension	26%	2,875
2	Diabetes	22%	2,421
3	Asthma	14%	1,577
4	Thyroid problems	8%	904
5	Ischaemic heart disease	3%	365
-	Other	26%	2,870

Breakdown of acute health conditions

by age
- 0-4: 27%
- 5 to 17: 29%
- 18 to 59: 33%
- > 60: 11%

by gender
- Male: 50%
- Female: 50%

Breakdown of chronic health conditions

by age
- 0-4: 7%
- 5 to 17: 12%
- 18 to 59: 50%
- > 60: 31%

by gender
- Male: 44%
- Female: 56%

UNHCR gratefully acknowledges the support of the following partners:

Public Health

Health Impact

	Indicator	Standard	
Crude Mortality Rate (CMR) (/1,000/month)	0.1	< 0.51	√
Under-five Mortality Rate (U5MR) (/1,000/month)	0.3	< 1.5	√
Neonatal Mortality Rate (NNMR) (/1,000 livebirths)	13.2	< 14.9	√

Access and utilization

	Indicator	Standard	
No. of health facilities	1 : 8566	1 : < 10,000	√
Total number of consultations	70,416		
Health utilisation rate	3.7	1 - 4	√
No. of consultations per trained clinician per day	25	< 50	√

Breakdown of mental health conditions

By cause

- 36% Severe emotional disorder, including moderate-severe depression
- 27% Epilepsy / seizures
- 24% Psychotic disorder (including mania)
- 10% Intellecutal disability
- 3% Other

By age

- 0%: 0-4
- 17%: 5 to 17
- 83%: 18 to 59
- > 60

By gender

- 56% Male
- 44% Female

Breakdown of injuries

By cause

- 29% Injury (burns)
- 1% Injury (war-related)
- Bites (all causes)
- Assault (with weapon non-war related)
- 70% Other

By age

- 20%: 0-4
- 29%: 5 to 17
- 33%: 18 to 59
- 18%: > 60

By gender

- 55% Male
- 45% Female

In-patient and referral

In-patient department

	Indicator	Standard	
Hospitalisation rate (/1000/year)	33.0	50 - 150	⚑
Bed occupancy rate	54%	> 75%	X
Mean length of stay for authorised discharges (days)	1.6		

	Indicator	Standard
Overall case fatality rate	0%	
Proportion of deaths within 24 hours of admission		

Reproductive Health

Antenatal care

	Indicator	Standard	
Coverage of antenatal care (4 or more visits)	99%	≥ 90%	√
Coverage of TT vaccination	99%	≥ 90%	√
Coverage of anaemia screening in pregnancy	99%	≥ 90%	√

Delivery care

	Indicator	Standard	
Proportion of deliveries attended by skilled personnel	100%	≥ 90%	√
Proportion of deliveries performed by caesarean section	35%	5 - 15%	X
Proportion of low birth weight deliveries (<2500g)	0.3%	< 15%	√

Maternal mortality

	Indicator	Standard
Number of maternal deaths reported	0	
% of maternal deaths investigated		100%

Postnatal care

	Indicator	Standard	
Coverage of postnatal care	68%	≥ 90%	X

Sexual and Gender-based Violence

	Indicator	Standard		
Incidence of rape (reported cases/10,000/year)*	0.5		^	⚑
% of rape survivors who received PEP** < 72 hours	100%	100%	^	⚑
% of rape survivors who received ECP*** < 120 hours	100%	100%	^	⚑
% of rape survivors who received STI**** prophylaxis < 2 wks	100%	100%	^	⚑

*Only includes rapes reported at health facilities **Post-exposure prophylaxis ***Emergency contraceptive pill
****Sexually transmitted infection