

1. Population

| Population | 0 to <1 | | Total <1 | 1 to 4 | | Total 1 to 4 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total |
|--------------------|---------|-------|----------|--------|-------|--------------|---------|--------|---------------|----------|--------|----------------|------|-------|------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | M | F | | |
| Refugee population | 1,610 | 1,534 | 3,144 | 5,828 | 5,546 | 11,374 | 14,227 | 13,486 | 27,713 | 16,105 | 16,301 | 32,406 | 853 | 1,191 | 2,044 | 76,681 |
| % by age group | | | 4% | | | 15% | | | 36% | | | 42% | | | 3% | 100% |

2. Mortality
2.1 Mortality

| Record ONE cause of mortality per case | 0 to 4 | | Total <5 | US % mort | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | Crude % mort |
|--|-----------|-----------|-----------|-------------|----------|----------|---------------|----------|----------|----------------|-----------|-----------|------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| 1. Lower Respiratory Tract Infection (LRTI) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 2 | 2 | 3 | 0 | 3 | 5 | 8% |
| 2. Watery diarrhoea | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Bloody diarrhoea | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4. Ischaemic heart disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 2% |
| 5. Cardiovascular disorders (other) | 2 | 0 | 2 | 7% | 0 | 0 | 0 | 2 | 1 | 3 | 3 | 3 | 6 | 11 | 18% |
| 6. Cerebrovascular disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 5 | 6 | 6 | 10% |
| 7. Chronic kidney disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 2% |
| 8. Injury | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 9. Cancer | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 1 | 1 | 2 | 4 | 0 | 4 | 6 | 10% |
| 10. Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 2% |
| 11. Tuberculosis (confirmed) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 12. Neonatal death | 10 | 7 | 17 | 63% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 17 | 27% |
| 13. COVID-19 | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 14. Other | 5 | 3 | 8 | 30% | 0 | 1 | 1 | 0 | 1 | 1 | 2 | 2 | 4 | 14 | 23% |
| Total | 17 | 10 | 27 | 100% | 0 | 1 | 1 | 4 | 5 | 9 | 13 | 12 | 25 | 62 | 100% |
| % by age group | | | 44% | | | | 2% | | | 15% | | | 40% | 100% | |

| Mortality indicators | Male | Female | Total |
|---|------|--------|-------|
| Crude Mortality Rate (CMR) (/1000/month) | 0.1 | 0.1 | 0.1 |
| Under 5 Mortality Rate (USMR) (/1000/month) | 0.4 | 0.2 | 0.3 |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | | | 13.2 |

2.2 Maternal death

| | |
|--|---|
| Number of maternal deaths reported | 0 |
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3. Morbidity
3.1 Staffing and service availability

| | |
|---|---------|
| Number of full-time clinicians | 42.0 |
| Average number of full days on which each health facility was functioning | 5.4 |
| Total number of outpatient consultations | 196,181 |
| Health facility utilisation rate | 5.0 |
| Consultations per clinician per day | 32 |

3.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total <5 | US Incid. | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause | Crude Incide |
|--|---------------|---------------|---------------|--------------|---------------|---------------|---------------|---------------|---------------|----------------|--------------|--------------|---------------|----------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | | |
| Upper Respiratory Tract Infection (URTI) | 5,729 | 5,447 | 11,176 | 770 | 6,003 | 6,054 | 12,057 | 5,699 | 7,102 | 12,801 | 1,544 | 1,449 | 2,993 | 39,027 | 26% | 509 |
| Influenza-like illness (ILI) | 3,312 | 3,133 | 6,445 | 444 | 3,096 | 3,113 | 6,209 | 2,609 | 2,903 | 5,512 | 1,047 | 1,055 | 2,102 | 20,268 | 14% | 264 |
| Lower Respiratory Tract Infection (LRTI) | 972 | 895 | 1,867 | 129 | 1,030 | 989 | 2,019 | 850 | 786 | 1,636 | 436 | 443 | 879 | 6,401 | 4% | 83 |
| Watery diarrhoea* | 111 | 104 | 215 | 15 | 79 | 64 | 143 | 45 | 47 | 92 | 25 | 17 | 42 | 492 | 0% | 6 |
| Bloody diarrhoea* | 1 | 2 | 3 | 0 | 2 | 1 | 3 | 1 | 1 | 2 | 0 | 0 | 0 | 8 | 0% | 0 |
| Skin infection | 1,223 | 1,380 | 2,603 | 179 | 1,236 | 1,475 | 2,711 | 1,026 | 1,232 | 2,258 | 419 | 412 | 831 | 8,403 | 6% | 110 |
| Eye infection | 790 | 738 | 1,528 | 105 | 854 | 806 | 1,660 | 933 | 1,000 | 1,933 | 385 | 404 | 789 | 5,910 | 4% | 77 |
| Ear infection | 1,358 | 1,277 | 2,635 | 181 | 1,112 | 1,097 | 2,209 | 930 | 985 | 1,915 | 500 | 315 | 815 | 7,574 | 5% | 99 |
| Dental Conditions | 458 | 443 | 901 | 62 | 1,674 | 1,693 | 3,367 | 3,481 | 4,144 | 7,625 | 319 | 374 | 693 | 12,586 | 9% | 164 |
| Intestinal Worms | 532 | 528 | 1,060 | 73 | 585 | 513 | 1,098 | 229 | 224 | 453 | 98 | 75 | 173 | 2,784 | 2% | 36 |
| Acute Jaundice Syndrome* | 5 | 0 | 5 | 0 | 7 | 0 | 7 | 1 | 2 | 3 | 2 | 1 | 3 | 18 | 0% | 0 |
| Measles (suspected)* | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 2 | 0% | 0 |
| Meningitis (suspected)* | 4 | 0 | 4 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 6 | 0% | 0 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 |
| Leishmaniasis | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0 |
| Fever of unknown origin | 376 | 392 | 768 | 53 | 370 | 447 | 817 | 428 | 429 | 857 | 387 | 325 | 712 | 3,154 | 2% | 41 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 0% | 0 |
| COVID-19 (suspected) | 0 | 0 | 0 | 0 | 4 | 0 | 4 | 9 | 1 | 10 | 3 | 3 | 6 | 20 | 0% | 0 |
| Other | 3,068 | 2,878 | 5,946 | 410 | 3,913 | 4,395 | 8,308 | 8,745 | 12,690 | 21,435 | 2,324 | 2,818 | 5,142 | 40,831 | 28% | 532 |
| Total | 17,939 | 17,218 | 35,157 | 2,422 | 19,966 | 20,647 | 40,613 | 24,986 | 31,547 | 56,533 | 7,489 | 7,693 | 15,182 | 147,485 | 100% | 1,923 |
| % by age group | | | 24% | | | | 28% | | | 38% | | | 10% | 100% | | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|--|--------------|------------|--------------|--------------|--------------|---------------|--------------|--------------|----------------|--------------|--------------|--------------|---------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| Diabetes | 24 | 35 | 59 | 63 | 140 | 203 | 1,248 | 1,812 | 3,060 | 889 | 1,324 | 2,213 | 5,535 | 22% |
| Hypertension | 14 | 15 | 29 | 67 | 108 | 175 | 1,571 | 2,283 | 3,854 | 1,206 | 1,727 | 2,933 | 6,991 | 27% |
| Asthma | 472 | 403 | 875 | 596 | 577 | 1,173 | 649 | 622 | 1,271 | 345 | 272 | 617 | 3,936 | 15% |
| Ischaemic heart disease | 4 | 4 | 8 | 4 | 9 | 13 | 241 | 165 | 406 | 178 | 166 | 344 | 771 | 3% |
| Cardiovascular disease (other) | 40 | 29 | 69 | 7 | 8 | 15 | 64 | 55 | 119 | 39 | 52 | 91 | 294 | 1% |
| Chronic obstructive pulmonary disease (COPD) | 1 | 0 | 1 | 1 | 0 | 1 | 28 | 20 | 48 | 56 | 52 | 108 | 158 | 1% |
| Liver disease | 0 | 1 | 1 | 4 | 8 | 12 | 24 | 31 | 55 | 25 | 22 | 47 | 115 | 0% |
| Thyroid problems | 34 | 13 | 47 | 131 | 192 | 323 | 265 | 751 | 1,016 | 274 | 287 | 561 | 1,947 | 8% |
| Cancer | 0 | 0 | 0 | 0 | 5 | 5 | 15 | 25 | 40 | 30 | 9 | 39 | 84 | 0% |
| Other | 430 | 478 | 908 | 571 | 660 | 1,231 | 1,027 | 1,395 | 2,422 | 690 | 602 | 1,292 | 5,853 | 23% |
| Total | 1,019 | 978 | 1,997 | 1,444 | 1,707 | 3,151 | 5,132 | 7,159 | 12,291 | 3,732 | 4,513 | 8,245 | 25,684 | 100% |
| % by age group | | | 8% | | | 12% | | | 48% | | | 32% | 100% | |

3.4 Mental health

| | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|--|----------|----------|-----------|------------|------------|---------------|--------------|--------------|----------------|----------|-----------|------------|--------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Epilepsy / seizures | 7 | 4 | 11 | 181 | 172 | 353 | 630 | 431 | 1,061 | 0 | 0 | 0 | 1,425 | 28% |
| 2. Alcohol and other substance use disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 0% |
| 3. Psychotic disorder (including mania) | 1 | 0 | 1 | 37 | 29 | 66 | 704 | 360 | 1,064 | 0 | 5 | 5 | 1,136 | 22% |
| 4. Intellectual disability | 0 | 0 | 0 | 70 | 121 | 191 | 231 | 86 | 317 | 0 | 0 | 0 | 508 | 10% |
| 5. Severe emotional disorder, including moderate-severe depression | 0 | 0 | 0 | 13 | 26 | 39 | 886 | 974 | 1,860 | 0 | 0 | 0 | 1,899 | 37% |
| 6. Medically unexplained somatic complaint | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0% |
| Other psychological complaints | 0 | 1 | 1 | 49 | 27 | 76 | 21 | 36 | 57 | 2 | 7 | 9 | 143 | 3% |
| Total | 8 | 5 | 13 | 350 | 375 | 725 | 2,473 | 1,888 | 4,361 | 2 | 12 | 14 | 5,113 | 100% |
| % by age group | | | 0% | | | 14% | | | 85% | | | | 0% | 100% |

3.5 Injuries

| | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|---|------------|------------|------------|------------|------------|---------------|------------|------------|----------------|------------|------------|------------|--------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Bites (all causes) | 1 | 1 | 2 | 7 | 2 | 9 | 1 | 0 | 1 | 1 | 0 | 1 | 13 | 0% |
| 2. Injury (war related) | 42 | 41 | 83 | 37 | 31 | 68 | 0 | 0 | 0 | 0 | 0 | 0 | 151 | 3% |
| 3. Injury (Burns) | 162 | 131 | 293 | 227 | 218 | 445 | 164 | 207 | 371 | 129 | 107 | 236 | 1,345 | 27% |
| 4. Assault (with weapon non-war related) | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 2 | 0% |
| Other | 335 | 281 | 616 | 561 | 435 | 996 | 736 | 651 | 1,387 | 266 | 230 | 496 | 3,495 | 70% |
| Total | 540 | 454 | 994 | 832 | 686 | 1,518 | 903 | 858 | 1,761 | 396 | 337 | 733 | 5,006 | 100% |
| % by age group | | | 20% | | | 30% | | | 35% | | | | 15% | 100% |

3.6 Reproductive Tract Infections

| | < 18 | | Total < 18 | ≥ 18 | | Total ≥ 18 | Grand total | % by cause |
|-------------------------|----------|--------------|--------------|-----------|--------------|--------------|--------------|-------------|
| | M | F | | M | F | | | |
| 1. Vaginal discharge | 0 | 484 | 484 | 8 | 2,814 | 2,822 | 3,306 | 39% |
| 2. Urethral discharge | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Genital ulcer | 0 | 0 | 0 | 0 | 3 | 3 | 3 | 0% |
| 3. Lower abdominal pain | 0 | 226 | 226 | 0 | 1,330 | 1,330 | 1,556 | 18% |
| Other | 0 | 332 | 332 | 49 | 3,281 | 3,330 | 3,662 | 43% |
| Total | 0 | 1,042 | 1,042 | 57 | 7,428 | 7,485 | 8,527 | 100% |
| % by age group | | | 12% | | | 88% | 100% | |

4. In-patient Department

4.1 In-patient Department Activities

| | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total |
|--|--------------|-----------|-----------|-----------|------------|---------------|------------|--------------|----------------|-----------|-----------|------------|--------------|
| | M | F | | M | F | | M | F | | M | F | | |
| Number at beginning of reporting period | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total admissions | 19 | 16 | 35 | 4 | 127 | 131 | 88 | 1,265 | 1,353 | 26 | 17 | 43 | 1,562 |
| Number of exits | authorized | 19 | 14 | 33 | 4 | 80 | 84 | 777 | 865 | 26 | 18 | 44 | 1,026 |
| | unauthorized | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 3 | 0 | 0 | 0 | 3 |
| | death | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | referral | 3 | 6 | 9 | 12 | 57 | 69 | 18 | 520 | 538 | 4 | 3 | 7 |
| Total exits | 22 | 20 | 42 | 16 | 137 | 153 | 106 | 1,406 | 1,406 | 30 | 21 | 51 | 1,652 |
| Deaths within 24 hours | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| | |
|---|-------|
| Number of days in reporting period | 189 |
| Number of beds | 22 |
| Sum number of days stay for authorised discharges | 1,416 |
| Proportion of < 5 IPD deaths within 24 hours | |
| Average length of stay (days) | 1.4 |
| Bed occupancy | 52% |
| Hospitalisation rate (/1000/month) | 3.3 |

4.2 In-patient Admissions and Deaths

| | 0 to 4 | | U5 CFR | 5 to 17 | | 5-17 CFR | 18 to 59 | | 18-59 CFR | > 60 | | Total adm | Total deaths | Crude CFR |
|-----------------------------|-----------|----------|-----------|------------|----------|-----------|--------------|----------|-----------|-----------|----------|--------------|--------------|-----------|
| | Adm | Deaths | | Adm | Deaths | | Adm | Deaths | | Adm | Deaths | | | |
| 1. Paediatric medicine | 33 | 0 | 0% | 2 | 0 | 0% | 0 | 0 | 0% | 0 | 0 | 35 | 0 | 0% |
| 2. General Surgery | 2 | 0 | 0% | 8 | 0 | 0% | 155 | 0 | 0% | 41 | 0 | 206 | 0 | 0% |
| 3. Injury (non-war related) | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | 0 | 0 | |
| 4. Internal medicine | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | 0 | 0 | |
| 5. Obstetrics | 0 | 0 | | 113 | 0 | 0% | 1,169 | 0 | 0% | 0 | 0 | 1,282 | 0 | 0% |
| 6. Gynaecology | 0 | 0 | | 8 | 0 | 0% | 34 | 0 | 0% | 0 | 0 | 42 | 0 | 0% |
| 7. Other | 0 | 0 | | 0 | 0 | | 1 | 0 | 0% | 0 | 0 | 1 | 0 | 0% |
| Total | 35 | 0 | 0% | 131 | 0 | 0% | 1,359 | 0 | 0% | 41 | 0 | 1,566 | 0 | 0% |

5. Referral

5.1 Referrals by location

| | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total | Grand total |
|--------------------------------|--------|---|-----------|---------|---|---------------|----------|---|----------------|------|---|-------|-------------|
| | M | F | | M | F | | M | F | | M | F | | |
| 1. MOH Hospital | 106 | | 106 | 55 | | 55 | 110 | | 110 | 61 | | 61 | 332 |
| 2. Jordan University Hospital | 51 | | 51 | 27 | | 27 | 38 | | 38 | 6 | | 6 | 122 |
| 3. Private affiliated Hospital | 268 | | 268 | 302 | | 302 | 1,050 | | 1,050 | 163 | | 163 | 1,783 |
| 4. Royal Medical Services | 1 | | 1 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 1 |
| 5. Other | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 |

| | |
|--|-------|
| Total No of referred cases out side the Camp | 2,238 |
|--|-------|

5.2 Referrals by service

| | | 0 to 4 | Total < 5 | 5 to 17 | Total 5 to 17 | 18 to 59 | Total 18 to 59 | > 60 | Total | Grand total |
|----|-------------------|--------|-----------|---------|---------------|----------|----------------|------|-------|-------------|
| 1. | Paediatrics | 335 | 335 | 67 | 67 | 0 | 0 | 0 | 0 | 402 |
| 2. | Internal medicine | 32 | 32 | 87 | 87 | 301 | 301 | 180 | 180 | 600 |
| 3. | Obstetric | 0 | 0 | 58 | 58 | 596 | 596 | 0 | 0 | 654 |
| 4. | Gynaecological | 0 | 0 | 5 | 5 | 86 | 86 | 0 | 0 | 91 |
| 5. | Surgical | 59 | 59 | 167 | 167 | 215 | 215 | 50 | 50 | 491 |
| 6. | Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

6. Laboratory

| | Total |
|--------------------------------------|--------|
| Number of laboratory tests conducted | 51,021 |

8. Reproductive Health

8.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|---------------------------------------|------|-------|-------|
| First antenatal visit < 1st trimester | 378 | 1,471 | 1,849 |
| First antenatal visit > 1st trimester | 87 | 784 | 871 |

ANC Indicators

| | |
|--|-----|
| Proportion of first time antenatal visits made < 1st trimester | 68% |
|--|-----|

8.2 Antenatal care in facility

| Number of pregnant women at time of delivery who received: | ≥ 4 ANC visits | 2 TT doses | anaemia screening | | | |
|--|----------------|------------|-------------------|--|--|--|
| | 112 | 1,142 | 1,254 | | | |
| | 112 | 1,142 | 1,254 | | | |
| | 112 | 1,142 | 1,254 | | | |

| | |
|--|-----|
| Coverage of complete antenatal care | 97% |
| Coverage of antenatal tetanus immunisation | 97% |
| Coverage of anaemia screening in pregnancy | 97% |

8.3 Delivery care

| | < 18 | ≥ 18 | Total |
|---|------|-------|-------|
| Live births | 114 | 1,177 | 1,291 |
| Still births | 0 | 0 | 0 |
| Low birth weight (<2500 g) | 0 | 6 | 6 |
| Attended by skilled health worker | 114 | 1,174 | 1,288 |
| Number of obstetric complications treated | 1 | 9 | 10 |
| Number of caesarean sections performed | 31 | 430 | 461 |

Delivery Indicators

| | |
|--|-------|
| Crude birth rate (/1000 population/ month) | 2.8 |
| Proportion of births attended by skilled health worker (excl. TBA) | 99.8% |
| Proportion of low birth weight deliveries | 0.5% |
| Proportion of all births performed by caesarean section | 36% |
| Proportion of obstetric complications treated among girls < 18 | 1% |

8.4 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|-------|-------|
| At least three PNC visits within 6 weeks | 120 | 1,043 | 1,163 |

Postnatal Indicators

| | |
|-------------------------------------|-----|
| Coverage of complete postnatal care | 90% |
|-------------------------------------|-----|

8.5 Family planning

| | New | | Repeat | | Total |
|------------------------------|------|------|--------|-------|-------|
| | < 18 | ≥ 18 | < 18 | ≥ 18 | |
| Oral contraceptives | 42 | 751 | 58 | 2,163 | 3,014 |
| Emergency Contraceptive Pill | 0 | 0 | 0 | 0 | 0 |
| Injectable (Depo-Provera) | 3 | 91 | 0 | 169 | 263 |
| Implantable (Norplant) | 0 | 0 | 0 | 0 | 0 |
| Intra-Uterine Device (IUD) | 1 | 178 | 0 | 29 | 208 |
| Condom (Male) | 23 | 285 | 25 | 797 | 1,130 |
| Other | 3 | 102 | 4 | 83 | 192 |

| | < 18 | ≥ 18 |
|-----------------------------|------|-------|
| Family Planning Counseling* | 412 | 6,434 |

8.6 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|---|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 1 | 0 | 0 | 1 |
| No. of rape survivors examined within 72 hours | 0 | 1 | 0 | 0 | 1 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 1 | 0 | 0 | 1 |
| No of female rape survivors given ECP within 120 hours | | 1 | | 0 | 1 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 1 | 0 | 0 | 1 |
| No. cases of trauma in health centre due to domestic violence | 0 | 3 | 0 | 1 | 4 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

| SGBV Indicators | |
|--|------|
| Incidence of reported rape (/10,000 population/year) | 0.3 |
| Proportion of rape survivors who receive PEP < 72 hrs | 100% |
| Proportion of female rape survivors who receive ECP < 120 hrs | 100% |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | 100% |

| | |
|---------------------|------------------------------|
| Data Sources | 1. Health Information System |
|---------------------|------------------------------|

| | |
|----------------------------|---|
| Contact Information | Amman Adam Musa Khalifa, UNHCR Email: khalifaa@unhcr.org |
|----------------------------|---|

UNHCR gratefully acknowledges the support of the following partners:

