

Population	0 to <1			1 to 4		5 to 17		18 to 59		> 60		Total > 60	Grand total			
	M	F	Total <1	M	F	Total 1 to 4	M	F	Total 18 to 59	M	F					
Refugee population	686	648	1,333	3,194	3,124	6,318	7,409	6,776	14,185	7,092	8,320	15,411	336	508	844	38,091
% by age group			4%			17%			37%			40%			2%	100%

2.1 Mortality

Record ONE cause of mortality per case	0 to 4			Total <5	US % mort	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total	Crude % mort
	M	F	Total <5			M	F		M	F		M	F			
1. Lower Respiratory Tract Infection (LRTI)	4	1	5	17%	0	0	0	0	1	0	1	2	2	4	10	11%
2. Watery diarrhoea	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0	0%
3. Bloody diarrhoea	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0	0%
4. Ischaemic heart disease	0	0	0	0%	0	0	0	2	1	3	2	0	2	5	5%	
5. Cardiovascular disorders (other)	2	1	3	10%	0	0	0	1	3	4	5	3	8	15	16%	
6. Cerebrovascular disease	0	1	1	3%	2	0	2	0	1	1	2	2	4	8	9%	
7. Chronic kidney disease	0	1	1	3%	0	0	0	0	0	0	0	1	0	1	2%	
8. Injury	1	0	1	3%	0	0	0	0	0	0	0	0	0	0	1%	
9. Cancer	0	0	0	0%	1	1	2	2	1	3	2	1	3	8	9%	
10. Chronic obstructive pulmonary disease (COPD)	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%	
11. Tuberculosis (confirmed)	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%	
12. Neonatal death	5	6	11	38%	0	0	0	0	0	0	0	0	0	11	12%	
13. COVID-19	0	0	0	0%	0	0	0	4	3	7	5	4	9	16	18%	
14. Other	4	3	7	24%	0	0	0	1	2	3	3	2	5	15	16%	
Total	16	13	29	100%	3	1	4	11	11	22	22	14	36	91	100%	
% by age group			32%				4%			24%			40%		100%	

Mortality indicators	Male	Female	Total
Crude Mortality Rate (CMR) (/1000 population/month)	0.2	0.2	0.2
Under 5 Mortality Rate (USMR) (/1000 population/month)	0.3	0.3	0.3
Neonatal Mortality Rate (NNMR) (/1000 population/month)			8.7

2.2 Maternal death

Number of maternal deaths reported	0
Number of maternal deaths investigated	0
Proportion of maternal deaths investigated	

3.1 Staffing and service availability

Number of full-time clinicians	28.7
Average number of full days on which each health facility was functioning	5.6
Total number of outpatient consultations	273,388
Health facility utilisation rate	7.2
Consultations per clinician per day	32

3.2 Acute health conditions

Record NEW cases of disease only	0 to 4			Total <5	US Incid	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total	% by cause	Crude Incide
	M	F	Total <5			M	F		M	F		M	F				
Upper Respiratory Tract Infection (URTI)	10,458	9,119	19,577	2558.7	6,697	6,515	13,212	5,045	6,609	11,654	246	327	573	45,016	26%	1,182	
Influenza-like illness (ILI)	1,981	1,832	3,813	498.4	837	770	1,607	1,461	1,555	3,016	44	54	98	8,534	5%	224	
Lower Respiratory Tract Infection (LRTI)	2,308	1,638	3,946	515.7	368	237	605	178	224	402	40	45	85	5,038	3%	132	
Watery diarrhoea*	1,608	1,400	3,008	393.1	375	349	724	297	476	773	14	27	41	4,546	3%	119	
Bloody diarrhoea*	3	6	9	1.2	1	2	3	0	2	2	0	0	0	14	0%	0.4	
Skin infection	891	944	1,835	239.8	744	1,263	2,007	766	1,305	2,071	48	63	111	6,024	3%	158	
Eye infection	617	540	1,157	151.2	427	388	815	718	764	1,482	53	33	86	3,540	2%	93	
Ear infection	891	794	1,685	220.2	510	647	1,157	864	1,625	2,489	65	107	172	5,503	3%	144	
Dental Conditions	276	282	558	72.9	3,497	4,151	7,648	7,954	11,894	19,848	158	197	355	28,409	16%	746	
Intestinal Worms	300	342	642	83.9	490	533	1,023	231	234	465	1	9	10	2,140	1%	56	
Acute Jaundice Syndrome*	10	9	19	2.5	9	13	22	4	2	6	1	0	1	48	0%	1.3	
Measles (suspected)*	0	0	0	0.0	0	1	1	0	0	0	0	0	0	1	0%	0.0	
Meningitis (suspected)*	4	2	6	0.8	3	4	7	0	0	0	0	0	0	13	0%	0.3	
Acute Flaccid Paralysis*	1	0	1	0.1	0	0	0	0	0	0	0	0	0	1	0%	0.0	
Leishmaniasis	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0	
Fever of unknown origin	10	5	15	2.0	2	2	4	6	2	8	0	1	1	28	0%	0.7	
Tuberculosis (suspected)	0	0	0	0.0	1	0	1	1	0	1	0	0	0	2	0%	0.1	
COVID-19 (suspected)	48	55	103	13.5	191	223	414	712	824	1,536	51	88	139	2,192	1%	58	
COVID-19 (confirmed)	16	15	31	4.1	89	82	171	287	446	733	25	35	60	995	1%	26	
Other	7,378	7,726	15,104	1974.1	6,496	8,925	15,421	10,802	18,346	29,148	1,247	1,558	2,805	62,478	36%	1,640	
Total	26,800	24,709	51,509	6732.2	20,737	24,105	44,842	29,326	44,308	73,634	1,993	2,544	4,537	174,522	100%	4,582	
% by age group			30%				26%			42%			3%	100%			

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

3.3 Chronic health conditions

Record NEW and REVISITS	0 to 4			Total < 5	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total	% by cause
	M	F	Total < 5		M	F		M	F		M	F			
Diabetes	0	0	0	54	64	118	647	799	1,446	275	422	697	2,261	18%	
Hypertension	1	3	4	2	8	10	1,306	1,586	2,892	531	815	1,346	4,252	34%	
Asthma	189	96	285	164	97	261	186	242	428	55	40	95	1,069	8%	
Ischaemic heart disease	0	0	0	5	5	10	245	212	457	135	128	263	725	6%	
Cardiovascular disease (other)	6	9	15	4	12	16	112	68	180	31	89	120	331	3%	
Chronic obstructive pulmonary disease (COPD)	0	0	0	0	0	0	43	13	56	41	29	70	126	1%	
Liver disease	0	0	0	2	0	2	1	4	5	5	1	6	13	0%	
Thyroid problems	7	0	7	38	58	96	189	668	857	11	56	67	1,027	8%	
Cancer	2	0	2	2	0	2	9	8	17	10	1	11	32	0%	
Other	140	181	321	99	135	234	930	892	1,822	234	204	438	2,815	22%	
Total	345	289	634	365	379	744	3,668	4,492	8,160	1,328	1,785	3,113	12,651	100%	
% by age group			5%			6%			65%			25%	100%		

5.2 Referrals by service

		0 to 4	Total < 5	5 to 17	Total 5 to 17	18 to 59	Total 18 to 59	> 60	Total	Grand total
1.	Paediatrics	386	386	63	63	0	0	0	0	449
2.	Internal medicine	71	71	282	282	1,071	1,071	129	129	1,553
3.	Obstetric	0	0	10	10	279	279	0	0	289
4.	Gynaecological	2	2	8	8	31	31	0	0	41
5.	Surgical	42	42	149	149	140	140	26	26	357
6.	Other	61	61	89	89	162	162	43	43	355

	Total
Number of laboratory tests conducted	110,078

8.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester	51	1,596	1,647
First antenatal visit > 1st trimester	19	452	471

ANC Indicators

Proportion of first time antenatal visits made < 1st trimester	78%
--	-----

8.2 Antenatal care in facility

	≥ 4 ANC visits	2 TT doses	anaemia screening
Number of pregnant women at time of delivery who received:	39	15	50
	1,023	264	1,202
	1,062	279	1,252

Coverage of complete antenatal care	84%
Coverage of antenatal tetanus immunisation	22%
Coverage of anaemia screening in pregnancy	99%

8.3 Delivery care

	< 18	≥ 18	Total
Live births	50	1,209	1,259
Still births	1	15	16
Low birth weight (<2500 g)	4	74	78
Attended by skilled health worker	50	1,182	1,232
Number of obstetric complications treated	0	2	2
Number of caesarean sections performed	11	299	310

Delivery Indicators

Crude birth rate (/1000 population/ month)	2.8
Proportion of births attended by skilled health worker (excl. TBA)	98%
Proportion of low birth weight deliveries	6%
Proportion of all births performed by caesarean section	25%
Proportion of obstetric complications treated among girls < 18	0%

8.4 Postnatal care

	< 18	≥ 18	Total
At least three PNC visits within 6 weeks	18	725	743

Postnatal Indicators

Coverage of complete postnatal care	59%
-------------------------------------	-----

8.5 Family planning

	New		Repeat		Total
	< 18	≥ 18	< 18	≥ 18	
Oral contraceptives	26	943	6	1394	2,369
Emergency Contraceptive Pill	0	0	0	0	0
Injectable (Depo-Provera)	2	154	0	115	271
Implantable (Norplant)	0	2	0	1	3
Intra-Uterine Device (IUD)	4	204	0	15	223
Condom (Male)	21	991	12	853	1,877
Other	0	0	0	0	0

	< 18	≥ 18
Family Planning Counseling*	143	7,344

8.6 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No of rape survivors given PEP within 72 hours	0	0	0	0	0
No of female rape survivors given ECP within 120 hours	0	0	0	0	0
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators	
Incidence of reported rape (/10,000 population/year)	0.0
Proportion of rape survivors who receive PEP < 72 hrs	
Proportion of female rape survivors who receive ECP < 120 hrs	
Proportion of rape survivors who receive STI presumptive treatment < 2 wks	

Note

This report for non-camp Syrian refugees in Jordan is compiled from health data obtained from reporting agencies and clinics. Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Published on

Contact Information

UNHCR
Adam Musa Khalifa
Email: KHALIFAA@unhcr.org



We gratefully acknowledge the contributions of the following agencies:

