

3.4 Mental health

| | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|---|----------|----------|-----------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Epilepsy / seizures | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 2. Alcohol and other substance abuse disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 3. Psychotic disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 4. Mental retardation / intellectual disability | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 5. Severe emotional disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 6. Medically unexplained somatic complaint | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Total | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100% |
| % by age group | | | | | | | | | | | | | 100% | |

3.5 Injuries

| | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|---------------------------|----------|----------|-----------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Injury (intentional) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 2. Injury (unintentional) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 3. Burns | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 4. Accident | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Total | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100% |
| % by age group | | | | | | | | | | | | | 100% | |

4. Referrals

4.1 Referrals - general

| | 0 to 4 | 5 to 14 | 15 to 49 | ≥ 50 | Grand total |
|----------------|---------|---------|----------|------|-------------|
| Emergency | NO DATA | | | | |
| Elective | | | | | |
| Other | | | | | |
| Total | | | | | |
| % by age group | | | | | |

ND: no data submitted

4.2 Referrals - by specialty

| | 0 to 4 | 5 to 14 | 15 to 59 | ≥ 50 | Grand total |
|-------------------|---------|---------|----------|------|-------------|
| Internal medicine | NO DATA | | | | |
| General Surgery | | | | | |
| Gynaecology | | | | | |
| Other | | | | | |
| Total | | | | | |
| % by age group | | | | | |

ND: no data submitted

5. Laboratory & other investigations

| | Total |
|-------------------------|---------|
| Haematology | NO DATA |
| Imaging (X-ray, MRI/CT) | |
| Other | |
| Total | |

6. Reproductive Health

6.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|-------|
| First antenatal visit < 1st trimester | 0 | 0 | 0 |
| First antenatal visit > 1st trimester | 0 | 0 | 0 |
| Number of pregnant women at time of delivery who received: | | | |
| ≥ 4 ANC visits | 0 | 0 | 0 |
| 2 TT doses | 0 | 0 | 0 |
| anaemia screening | 0 | 0 | 0 |

ND: no data submitted

ANC Indicators

Proportion of first time antenatal visits made < 1st trimester

6.2 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|-------|
| Women who attended 3 postnatal visits within 6 wks of delivery | 0 | 0 | 0 |

ND: no data submitted

6.3 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|---|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | | | | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 0 | 0 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators

| | |
|--|-----|
| Incidence of reported rape | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

Data Sources

1. Health Information System

Publication date:

20 January 2014

Contact Information

Erbil
Dr Chimmen Akrawi, DOH Erbil

Email: chimmentya@yahoo.com

Erbil
Mohammed Marzoog, UNHCR

Email: MARZOOG@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

