

1. Population

Population	0 to < 1		Total <1	1 to 4		Total 1 to 4	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total
	M	F		M	F		M	F		M	F		M	F		
Refugee population	712	712	1,424	2,850	2,850	5,700	5,137	4,452	9,589	15,788	9,795	25,583	1,336	1,336	2,672	44,968
% by age group			3%			13%			21%			57%			6%	100%

2. Mortality

2.1 Mortality

Record ONE cause of mortality per case	0 to 4		Total <5	U5 % mort	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	Crude % mort
	M	F			M	F		M	F		M	F			
1. Lower Respiratory Tract Infection (LRTI)	0	0	0		0	0	0	0	0	0	0	0	0	0	
2. Watery diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	
3. Bloody diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	
4. Cardiovascular disorders	0	0	0		0	0	0	0	0	0	0	0	0	0	
5. Ischaemic heart disease	0	0	0		0	0	0	0	0	0	0	0	0	0	
6. Injury	0	0	0		0	0	0	0	0	0	0	0	0	0	
7. Cancer	0	0	0		0	0	0	0	0	0	0	0	0	0	
8. Chronic obstructive pulmonary disease (COPD)	0	0	0		0	0	0	0	0	0	0	0	0	0	
9. Tuberculosis (confirmed)	0	0	0		0	0	0	0	0	0	0	0	0	0	
10. Neonatal death	0	0	0		0	0	0	0	0	0	0	0	0	0	
Other	0	0	0		0	0	0	0	0	0	0	0	0	0	
Total	0	0	0	100%	0	0	0	0	0	0	0	0	0	0	100%
% by age group															

Mortality indicators	Male	Female	Total
Crude Mortality Rate (CMR)	0.0	0.0	0.0
Under 5 Mortality Rate (U5MR)	0.0	0.0	0.0
Neonatal Mortality Rate (NNMR)			0.0

2.2 Maternal death

Number of maternal deaths reported	0
Number of maternal deaths investigated	0
Proportion of maternal deaths investigated	

3. Morbidity

3.1 Staffing and service availability

Number of full-time clinicians	5
Average number of full days on which each health facility was	7.0
Total number of outpatient consultations	1,933
Health facility utilisation rate	2.2
Consultations per clinician per day	55

3.2 Acute health conditions

Record NEW cases of disease only	0 to 4		Total <5	U5 Incid**	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause	Crude Incid**
	M	F			M	F		M	F		M	F				
Upper Respiratory Tract Infection (URTI)	221	180	401	56.3	130	83	213	76	111	187	10	5	15	816	61%	18.1
Lower Respiratory Tract Infection (LRTI)	6	2	8	1.1	0	0	0	1	2	3	0	1	1	12	1%	0.3
Watery diarrhoea*	7	5	12	1.7	1	0	1	0	1	1	0	0	0	14	1%	0.3
Bloody diarrhoea*	1	4	5	0.7	1	0	1	0	0	0	0	0	0	6	0%	0.1
Malaria (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Skin infection	5	1	6	0.8	1	2	3	1	2	3	0	1	1	13	1%	0.3
Eye infection	0	1	1	0.1	0	0	0	2	0	2	1	0	1	4	0%	0.1
Ear infection	8	5	13	1.8	2	3	5	2	2	4	0	1	1	23	2%	0.5
Dental Conditions	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Intestinal Worms	1	0	1	0.1	0	0	0	0	0	0	0	1	1	2	0%	0.0
Acute Jaundice Syndrome*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Measles (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Meningitis (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute Flaccid Paralysis*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Tuberculosis (suspected)	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Fever of unknown origin	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Sexually Transmitted Infection (STI)	0	0	0	0.0	0	7	7	0	21	21	0	0	0	28	2%	0.6
Urinary tract infection (UTI)	0	0	0	0.0	2	4	6	5	34	39	1	1	2	47	3%	1.0
HIV/AIDS	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute malnutrition	0	0	0	0.0	1	0	1	0	0	0	0	0	0	1	0%	0.0
Leishmaniasis	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Other	58	32	90	12.6	41	20	61	69	135	204	11	15	26	381	28%	8.5
Total	307	230	537	75.4	179	119	298	156	308	464	23	25	48	1347	100%	30.0
% by age group			40%				22%			34%			4%	100%		

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **Incidence is cases per 1000 per week

3.3 Chronic health conditions

Record NEW and REVISITS	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
Diabetes	0	0	0	0	0	0	2	2	4	1	8	9	13	28%
Hypertension	0	0	0	0	0	0	2	1	3	3	6	9	12	26%
Asthma	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Ischaemic heart disease	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Cardiovascular disease (other)	0	0	0	0	0	0	4	2	6	3	0	3	9	20%
Chronic obstructive pulmonary disease (COPD)	0	0	0	1	0	1	4	7	11	0	0	0	12	26%
Liver disease	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Thyroid problems	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Cancer	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Total	0	0	0	1	0	1	12	12	24	7	14	21	46	100%
% by age group			0%			2%			52%			46%	100%	

3.4 Mental health

	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
1. Epilepsy / seizures	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0
3. Psychotic disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4. Mental retardation / intellectual disability	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5. Severe emotional disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6. Medically unexplained somatic complaint	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	0	0	0	0	0	0	0	0	0	0	0	0	0	100%
% by age group													100%	

3.5 Injuries

	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
1. Injury (intentional)	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
2. Injury (unintentional)	2	0	2	5	2	7	18	8	26	21	20	41	76	14%
3. Burns	5	7	12	1	1	2	11	16	27	10	14	24	65	12%
4. Accident	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	23	55	78	107	85	192	30	30	60	28	41	69	399	74%
Total	30	62	92	113	88	201	59	54	113	59	75	134	540	100%
% by age group			17%			37%			21%			25%	100%	

4. Referrals

	0 to 4	5 to 14	15 to 49	≥ 50	Grand total
Emergency	2	1	6	2	11
Elective	27	13	30	7	77
Other	0	0	0	0	0
Total	29	14	36	9	88
% by age group	33%	16%	41%	10%	100%

5. Laboratory & other investigations

	Total
Haematology	141
Imaging (X-ray, MRI/CT)	0
Other	132
Total	391

6. EPI and Vitamin A

7.1 Vaccination

	<1	≥1 to <5	Total
BCG			
Hepatitis B birth dose			
DPT I/HiB I/Hep B I* (Pentavalent I)			
DPT II/HiB II* (Tetravalent)			
DPT III/HiB III/Hep B II* (Pentavalent II)			
Polio 0			
Polio I			
Polio II			
Polio III			
Measles			
MMR			
Rotavirus 1			
Rotavirus 2			
Rotavirus 3			
Vitamin A			
Additional vaccines			

NO DATA

EPI Indicators (since December 2013)	
Total vaccinated this week	
a. Program dropout rate*	
b. Immunisation coverage	
BCG	
Hepatitis B birth dose	
DPT I	
DPT II	
Polio I	
Polio II	
Polio III	
Measles	
MMR**	
Rotavirus I	
Rotavirus II	
c. Vitamin A coverage	

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%
**For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

7. Reproductive Health

7.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester	14	65	79
First antenatal visit > 1st trimester	0	0	0
Number of pregnant women at time of delivery who received:			
≥ 4 ANC visits	0	0	0
2 TT doses	0	0	0
anaemia screening	0	0	0

ANC Indicators	
Proportion of first time antenatal visits made < 1st trimester	100%

7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery	0	40	40

7.3 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 - 120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No. of rape survivors given PEP within 72 hours	0	0	0	0	0
No. of female rape survivors given ECP within 120 hours	0	0	0	0	0
No. of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators	
Incidence of reported rape	0.0
Proportion of rape survivors who receive PEP < 72 hrs	
Proportion of female rape survivors who receive ECP < 120 hrs	
Proportion of rape survivors who receive STI presumptive treatment < 2 wks	