

1. Population

| Population | 0 to <1 | | Total <1 | 1 to 4 | | Total 1 to 4 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total |
|--------------------|---------|----|----------|--------|-----|--------------|---------|-----|---------------|----------|-------|----------------|------|-----|------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | M | F | | |
| Refugee population | 95 | 95 | 190 | 378 | 379 | 757 | 838 | 678 | 1,516 | 1,505 | 1,257 | 2,762 | 427 | 388 | 815 | 6,040 |
| % by age group | | | 3% | | | 13% | | | 25% | | | 46% | | | 13% | 100% |

* estimates from registered population

2. Mortality

2.1 Mortality

| Record ONE cause of mortality per case | 0 to 4 | | Total <5 | U5 % mort | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total > 50 | Grand total | Crude % mort |
|---|----------|----------|----------|-------------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| 1. Lower Respiratory Tract Infection (LRTI) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 2. Watery diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 3. Bloody diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 4. Cardiovascular disorders | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 5. Ischaemic heart disease | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 6. Injury | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 7. Cancer | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 8. Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 9. Tuberculosis (confirmed) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 10. Neonatal death | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Other | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Total | 0 | 0 | 0 | 100% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100% |
| % by age group | | | | | | | | | | | | | | | |

| Mortality indicators | Male | Female | Total |
|--------------------------------|------|--------|-------|
| Crude Mortality Rate (CMR) | 0.0 | 0.0 | 0.0 |
| Under 5 Mortality Rate (USMR) | 0.0 | 0.0 | 0.0 |
| Neonatal Mortality Rate (NNMR) | | | 0.0 |

2.2 Maternal death

| | |
|--|---|
| Number of maternal deaths reported | 0 |
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3. Morbidity

3.1 Staffing and service availability

| | |
|---|-------|
| Number of full-time clinicians | 3 |
| Average number of full days on which each health facility was | 7.0 |
| Total number of outpatient consultations | 1,360 |
| Health facility utilisation rate | 11.7 |
| Consultations per clinician per day | 65 |

3.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total <5 | U5 Incid** | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause | Crude Incid** |
|--|------------|------------|------------|--------------|------------|-----------|---------------|------------|------------|----------------|-----------|-----------|------------|-------------|-------------|---------------|
| | M | F | | | M | F | | M | F | | M | F | | | | |
| Upper Respiratory Tract Infection (URTI) | 59 | 33 | 92 | 97.1 | 38 | 48 | 86 | 104 | 175 | 279 | 17 | 19 | 36 | 493 | 44% | 81.6 |
| Lower Respiratory Tract Infection (LRTI) | 81 | 78 | 159 | 167.9 | 36 | 29 | 65 | 21 | 18 | 39 | 1 | 1 | 2 | 265 | 24% | 43.9 |
| Watery diarrhoea* | 2 | 4 | 6 | 6.3 | 0 | 0 | 0 | 3 | 3 | 6 | 1 | 0 | 1 | 13 | 1% | 2.2 |
| Bloody diarrhoea* | 1 | 0 | 1 | 1.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% | 0.2 |
| Malaria (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Skin infection | 7 | 3 | 10 | 10.6 | 2 | 2 | 4 | 6 | 5 | 11 | 0 | 1 | 1 | 26 | 2% | 4.3 |
| Eye infection | 2 | 3 | 5 | 5.3 | 8 | 2 | 10 | 7 | 7 | 14 | 0 | 2 | 2 | 31 | 3% | 5.1 |
| Ear infection | 0 | 2 | 2 | 2.1 | 7 | 2 | 9 | 7 | 6 | 13 | 0 | 2 | 2 | 26 | 2% | 4.3 |
| Dental Conditions | 0 | 0 | 0 | 0.0 | 4 | 1 | 5 | 19 | 16 | 35 | 2 | 1 | 3 | 43 | 4% | 7.1 |
| Intestinal Worms | 0 | 3 | 3 | 3.2 | 4 | 0 | 4 | 4 | 1 | 5 | 0 | 0 | 0 | 12 | 1% | 2.0 |
| Acute Jaundice Syndrome* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Measles (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Meningitis (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Fever of unknown origin | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Sexually Transmitted Infection (STI) | 0 | 0 | 0 | 0.0 | 0 | 2 | 2 | 0 | 114 | 114 | 0 | 0 | 0 | 116 | 10% | 19.2 |
| Urinary tract infection (UTI) | 0 | 0 | 0 | 0.0 | 1 | 1 | 2 | 2 | 1 | 3 | 16 | 0 | 16 | 21 | 2% | 3.5 |
| HIV/AIDS | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute malnutrition | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 3 | 3 | 0 | 1 | 1 | 4 | 0% | 0.7 |
| Leishmaniasis | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Other | 0 | 1 | 1 | 1.1 | 0 | 9 | 9 | 12 | 25 | 37 | 8 | 10 | 18 | 65 | 6% | 10.8 |
| Total | 152 | 127 | 279 | 294.6 | 100 | 96 | 196 | 185 | 374 | 559 | 45 | 37 | 82 | 1116 | 100% | 184.8 |
| % by age group | | | 25% | | | | 18% | | 50% | | | | 7% | 100% | | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **incidence is cases per 1000 per week

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause | |
|--|----------|----------|-----------|----------|----------|---------------|-----------|-----------|----------------|-----------|-----------|------------|-------------|-------------|--|
| | M | F | | M | F | | M | F | | M | F | | | | |
| Diabetes | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 3 | 7 | 5 | 8 | 13 | 20 | 16% | |
| Hypertension | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 8 | 16 | 9 | 12 | 21 | 37 | 29% | |
| Asthma | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 2 | 3 | 4 | 3% | |
| Ischaemic heart disease | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 3 | 4 | 5 | 2 | 7 | 11 | 9% | |
| Cardiovascular disease (other) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | |
| Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | |
| Liver disease | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | |
| Thyroid problems | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | |
| Cancer | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | |
| Other | 0 | 0 | 0 | 1 | 4 | 5 | 15 | 21 | 36 | 5 | 10 | 15 | 56 | 44% | |
| Total | 0 | 0 | 0 | 1 | 4 | 5 | 28 | 36 | 64 | 25 | 34 | 59 | 128 | 100% | |
| % by age group | | | 0% | | | 4% | | | 50% | | | 46% | 100% | | |

3.4 Mental health

| | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|---|----------|----------|-----------|----------|----------|---------------|----------|----------|----------------|-----------|----------|------------|-------------|------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Epilepsy / seizures | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 2 | 2% |
| 2. Alcohol and other substance abuse disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Psychotic disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4. Mental retardation / intellectual disability | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 5. Severe emotional disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 6. Medically unexplained somatic complaint | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 4 | 6 | 82 | 6 | 88 | 94 | 98% |
| Total | 1 | 0 | 1 | 0 | 0 | 0 | 2 | 4 | 6 | 82 | 6 | 88 | 94 | 98% |
| % by age group | | | 1% | | | 0% | | | 7% | | | 92% | 100% | |

3.5 Injuries

| | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|---------------------------|----------|----------|------------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Injury (intentional) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 2. Injury (unintentional) | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 5% |
| 3. Burns | 2 | 0 | 2 | 0 | 0 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 4 | 20% |
| 4. Accident | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 2 | 10% |
| Other | 0 | 0 | 0 | 0 | 1 | 1 | 5 | 5 | 10 | 2 | 0 | 2 | 13 | 65% |
| Total | 3 | 1 | 4 | 0 | 1 | 1 | 7 | 6 | 13 | 2 | 0 | 2 | 20 | 100% |
| % by age group | | | 20% | | | 5% | | | 65% | | | 10% | 100% | |

4. Referrals

4.1 Referrals - general

| | 0 to 4 | 5 to 14 | 15 to 49 | ≥ 50 | Grand total |
|-----------------------|----------|----------|----------|----------|-------------|
| Emergency | 0 | 0 | 0 | 0 | 0 |
| Elective | 0 | 0 | 0 | 0 | 0 |
| Other | 0 | 0 | 0 | 0 | 0 |
| Total | 0 | 0 | 0 | 0 | 0 |
| % by age group | | | | | 100% |

ND: no data submitted

5. Laboratory & other investigations

| | Total |
|-------------------------|------------|
| Haematology | 0 |
| Imaging (X-ray, MRI/CT) | 0 |
| Other | 0 |
| Total | 168 |

6. EPI and Vitamin A

| | <1 | 1 to <5 | Total |
|--|----|---------|-------|
| BCG | 4 | 0 | 4 |
| Hepatitis B birth dose | 4 | 0 | 4 |
| DPT I/HiB I/Hep B I* (Pentavalent I) | 2 | 0 | 2 |
| DPT II/HiB II* (Tetravalent) | 6 | 0 | 6 |
| DPT III/HiB III/Hep B II* (Pentavalent II) | 4 | 0 | 4 |
| Polio 0 | 2 | 0 | 2 |
| Polio I | 2 | 0 | 2 |
| Polio II | 6 | 0 | 6 |
| Polio III | 4 | 0 | 4 |
| Measles | 9 | 0 | 9 |
| MMR | 0 | 5 | 5 |
| Rotavirus 1 | 1 | 0 | 1 |
| Rotavirus 2 | 2 | 0 | 2 |
| Rotavirus 3 | 1 | 0 | 1 |
| Vitamin A | 15 | 0 | 15 |
| Additional vaccines | 0 | 12 | 12 |

| EPI Indicators (since December 2013) | | |
|--------------------------------------|------------------------|------|
| Total vaccinated this week | | 79 |
| a. Program dropout rate* | | -25% |
| b. Immunisation coverage | | |
| | BCG | 60% |
| | Hepatitis B birth dose | 60% |
| | DPT I | 60% |
| | DPT III | 75% |
| | Polio I | 78% |
| | Polio III | 82% |
| | Measles | 164% |
| | MMR** | 105% |
| | Rotavirus I | 37% |
| | Rotavirus III | 37% |
| c. Vitamin A coverage | | 110% |

*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%
**For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

7. Reproductive Health

7.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|--|----------------|------|-------|
| First antenatal visit < 1st trimester | | | |
| First antenatal visit > 1st trimester | | | |
| Number of pregnant women at time of delivery who received: | NO DATA | | |
| ≥ 4 ANC visits | | | |
| 2 TT doses | | | |
| anaemia screening | | | |

ND: no data submitted

| ANC Indicators | |
|--|--|
| Proportion of first time antenatal visits made < 1st trimester | |

7.2 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|----------------|
| Women who attended 3 postnatal visits within 6 wks of delivery | | | NO DATA |

ND: no data submitted

6.3 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|---|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | | | | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 0 | 0 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

| SGBV Indicators | | |
|--|--|------------|
| Incidence of reported rape | | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | | |
| Proportion of female rape survivors who receive ECP < 120 hrs | | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | | |

Data Sources

1. Health Information System

Publication date:

04 March 2014

Contact Information

Erbil
Dr Chimmen Akrawi, DOH Erbil

Email: chimmentya@yahoo.com

Erbil
Mohammed Marzooq, UNHCR

Email: MARZOOQ@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

