



### 3.4 Mental health

	0 to 4			5 to 14			15 to 49			≥ 50			Total	Grand total	% by cause
	M	F	Total < 5	M	F	Total 5 to 14	M	F	Total 15 to 49	M	F	Total ≥ 50			
1. Epilepsy / seizures	0	0	0	0	0	0	0	1	1	0	0	0	0	1	100%
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
3. Psychotic disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
4. Mental retardation / intellectual disability	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
5. Severe emotional disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
6. Medically unexplained somatic complaint	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
<b>Total</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>100%</b>
% by age group			0%			0%			100%					0%	100%

### 3.5 Injuries

	0 to 4			5 to 14			15 to 49			≥ 50			Total	Grand total	% by cause
	M	F	Total < 5	M	F	Total 5 to 14	M	F	Total 15 to 49	M	F	Total ≥ 50			
1. Injury (intentional)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
2. Injury (unintentional)	2	0	2	0	0	0	0	0	0	0	0	0	2	17%	
3. Burns	2	4	6	1	2	3	0	0	0	0	0	0	9	75%	
4. Accident	0	0	0	1	0	1	0	0	0	0	0	0	1	8%	
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0%	
<b>Total</b>	<b>4</b>	<b>4</b>	<b>8</b>	<b>2</b>	<b>2</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>12</b>	<b>100%</b>	
% by age group			67%			33%			0%				0%	100%	

## 4. Referrals

### 4.1 Referrals - general

	0 to 4	5 to 14	15 to 49	≥ 50	Grand total
Emergency	0	0	0	0	0
Elective	0	0	0	0	0
Other	0	0	0	0	0
<b>Total</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>
% by age group					100%

ND: no data submitted

## 5. Laboratory & other investigations

	Total
Haematology	0
Imaging (X-ray, MRI/CT)	0
Other	0
<b>Total</b>	<b>0</b>

## 6. EPI and Vitamin A

	<1	≥1 to <5	Total
BCG	0	0	0
Hepatitis B birth dose	0	0	0
DPT I/HiB I/Hep B I* (Pentavalent I)	1	0	1
DPT II/HiB II* (Tetavalent)	3	0	3
DPT III/HiB III/Hep B II* (Pentavalent II)	0	0	0
Polio 0	0	0	0
Polio I	1	0	1
Polio II	3	0	3
Polio III	0	0	0
Measles	2	0	2
MMR	0	3	3
Rotavirus 1	1	0	1
Rotavirus 2	3	0	3
Rotavirus 3	0	0	0
Vitamin A	5	0	5
Additional vaccines	0	8	8

EPI Indicators (since December 2013)		
Total vaccinated this week		30
a. Program dropout rate*		
Program dropout rate		-20%
b. Immunisation coverage		
BCG		116%
Hepatitis B birth dose		34%
DPT I		51%
DPT III		61%
Polio I		51%
Polio III		68%
Measles		116%
MMR**		82%
Rotavirus I		48%
Rotavirus III		61%
c. Vitamin A coverage		
Vitamin A coverage		108%

\*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines  
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%  
\*\*For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

## 7. Reproductive Health

### 7.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester	0	0	0
First antenatal visit > 1st trimester	0	0	0
Number of pregnant women at time of delivery who received:			
≥ 4 ANC visits	0	0	0
2 TT doses	0	0	0
anaemia screening	0	0	0

ND: no data submitted

ANC Indicators	
Proportion of first time antenatal visits made < 1st trimester	

### 7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery	0	0	0

ND: no data submitted

### 7.3 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No of rape survivors given PEP within 72 hours	0	0	0	0	0
No of female rape survivors given ECP within 120 hours					0
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators	
Incidence of reported rape	0.0
Proportion of rape survivors who receive PEP < 72 hrs	
Proportion of female rape survivors who receive ECP < 120 hrs	
Proportion of rape survivors who receive STI presumptive treatment < 2 wks	

Data Sources: 1. Health Information System

Publication date: 31 March 2014

Contact Information: Erbil, Dr Chimmen Akrawi, DOH Erbil

Email: chimmmentya@yahoo.com

Erbil, Sandra Harlass, UNHCR

Email: HARLASS@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

