



### 3.4 Mental health

	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
1. Epilepsy / seizures	0	0	0	0	0	0	2	2	4	0	0	0	4	15%
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
3. Psychotic disorder	0	0	0	0	0	0	4	4	8	0	0	0	8	31%
4. Mental retardation / intellectual disability	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
5. Severe emotional disorder	0	0	0	0	0	0	1	6	7	0	0	0	7	27%
6. Medically unexplained somatic complaint	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	1	0	1	1	1	2	2	2	4	0	0	0	7	27%
<b>Total</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>9</b>	<b>14</b>	<b>23</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>26</b>	<b>100%</b>
% by age group			4%			8%			88%			0%	100%	

### 3.5 Injuries

	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
1. Injury (intentional)	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
2. Injury (unintentional)	7	3	10	5	3	8	15	21	36	19	14	33	87	20%
3. Burns	4	3	7	2	3	5	9	17	26	14	7	21	59	14%
4. Accident	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	26	31	57	75	62	137	30	16	46	26	18	44	284	66%
<b>Total</b>	<b>37</b>	<b>37</b>	<b>74</b>	<b>82</b>	<b>68</b>	<b>150</b>	<b>54</b>	<b>54</b>	<b>108</b>	<b>59</b>	<b>39</b>	<b>98</b>	<b>430</b>	<b>100%</b>
% by age group			17%			35%			25%			23%	100%	

## 4. Referrals

### 4.1 Referrals - general

	0 to 4	5 to 14	15 to 49	≥ 50	Grand total
Emergency	1	0	7	1	9
Elective	9	7	44	14	74
Other	0	0	0	0	0
<b>Total</b>	<b>10</b>	<b>7</b>	<b>51</b>	<b>15</b>	<b>83</b>
% by age group	12%	8%	61%	18%	100%

## 5. Laboratory & other investigations

	Total
Haematology	9
Imaging (X-ray, MRI/CT)	74
Other	0
<b>Total</b>	<b>83</b>

## 6. EPI and Vitamin A

	<1	≥1 to <5	Total
BCG	0	0	0
Hepatitis B birth dose	0	0	0
DPT I/HiB I/Hep B I* (Pentavalent I)	1	0	1
DPT II/HiB II* (Tetravalent)	1	0	1
DPT III/HiB III/Hep B II* (Pentavalent II)	0	0	0
Polio 0	0	0	0
Polio I	1	0	1
Polio II	1	0	1
Polio III	1	0	1
Measles	2	0	2
MMR	0	0	0
Rotavirus 1	0	0	0
Rotavirus 2	0	0	0
Rotavirus 3	0	0	0
Vitamin A	1	0	1
Additional vaccines	0	8	8

EPI Indicators (since December 2013)		Total
Total vaccinated this week		16
a. Program dropout rate*		25%
b. Immunisation coverage		
BCG		94%
Hepatitis B birth dose		91%
DPT I		101%
DPT III		76%
Polio I		102%
Polio III		76%
Measles		53%
MMR**		88%
Rotavirus I		80%
Rotavirus III		48%
c. Vitamin A coverage		89%

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

\*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines  
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%  
\*\*For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.  
NAD=No adequate data; NR=Some or all camps did not consistently report

## 7. Reproductive Health

### 7.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester	5	95	100
First antenatal visit > 1st trimester	0	2	2
Number of pregnant women at time of delivery who received:			
≥ 4 ANC visits	0	0	0
2 TT doses	0	0	0
anaemia screening	0	0	0

#### ANC Indicators

Proportion of first time antenatal visits made < 1st trimester	98%
--	-----

### 7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery	0	38	38

## 6.3 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No of rape survivors given PEP within 72 hours	0	0	0	0	0
No of female rape survivors given ECP within 120 hours	0	0	0	0	0
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators		Total
Incidence of reported rape		0.0
Proportion of rape survivors who receive PEP < 72 hrs		
Proportion of female rape survivors who receive ECP < 120 hrs		
Proportion of rape survivors who receive STI presumptive treatment < 2 wks		

Data Sources: 1. Health Information System

Publication date:

15 April 2014

Contact Information

Duhok  
Dr Bakhtiyar AHMED, DOH Duhok,  
Email: BAHMED@duhokhealth.org

Erbil  
Sandra Harlass, UNHCR  
Email: HARLASS@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

