

| Population | 0 to < | :1 | Total <1 | 1 to |) 4 E | Total 1 to 4 | 5 t | o 17 | Total 5 to | 18 to | 59 E | Total 18 to 59 | > 6 M | 50 E | Total > | Grand total |
|--------------------|--------|-----|----------|-------|----------|-----------------|-------|-------|------------|-------|---------|-------------------|----------|---------|---------|----------------|
| Refugee population | 657 | 620 | 1,277 | 3,463 | 3,288 | | 7,159 | 6,721 | 13,880 | 6,684 | 7,185 | | 301 | 415 | 716 | |
| % by age group | | | 4% | | | 19% | | | 38% | | | 38% | | | 2% | 100% |

2.1 Mortality

| | Record ONE cause of mortality per case | 0 to | 4 | Total <5 | U5 % | 5 to | 17 | Total 5 | 18 to 5 | 59 | Total 18 | > (| 60 | Total > | Grand | Crude % |
|-----|--|------|---|----------|------|------|----|---------|---------|----|----------|-----|----|---------|-------|---------|
| | Record ONE cause of mortality per case | M | F | TOTAL <3 | mort | M | F | to 17 | М | F | to 59 | M | F | 60 | total | mort |
| 1. | Lower Respiratory Tract Infection (LRTI) | 1 | 1 | 2 | 40% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 3 | 21% |
| 2. | Watery diarrhoea | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. | Bloody diarrhoea | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4. | Ischaemic heart disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 7% |
| 5. | Cardiovascular disorders (other) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 1 | 0 | 1 | 2 | 1 | 3 | 4 | 29% |
| 6. | Cerebrovascular disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 7% |
| 7. | Chronic kidney disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 8. | Injury | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 9. | Cancer | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 7% |
| 10. | Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 7% |
| 11. | Tuberculosis (confirmed) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 12. | Neonatal death | 2 | 1 | 3 | 60% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 21% |
| 13. | Other | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| | Total | 3 | 2 | 5 | 100% | 0 | 0 | 0 | 3 | 0 | 3 | 4 | 2 | 6 | 14 | 100% |
| | % by age group | | | 36% | | | | 0% | | | 21% | | | 43% | 100% | |

| Mortality indicators | Male | Female | Total |
|---|------|--------|-------|
| Crude Mortality Rate (CMR) (/1000 pooulation/month) | 0.2 | 0.1 | 0.1 |
| Under 5 Mortality Rate (U5MR) (/1000 pooulation/month) | 0.2 | 0.2 | 0.2 |
| Neonatal Mortality Rate (NNMR) (/1000 pooulation/month) | | | 8.2 |

2.2 Maternal death

| Number of maternal deaths reported | . 0 |
|--|-----|
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3.1 Staffing and service availability

| Number of full-time clinicians | 22.3 |
|---|--------|
| Average number of full days on which each health facility was functioning | 5.7 |
| Total number of outpatient consultations | 73,558 |
| Health facility utilisation rate | 8.1 |
| Consultations per clinician per day | 45 |

3.2 Acute health conditions

| Bear of Marcon of Marcon only | 0 to | 4 | T-1-1 5 | U5 Incid | 5 to | 17 | Total 5 | 18 to | 59 | Total 18 | > 60 | 0 | Total > | Grand | % by | Crude |
|--|-------|-------|----------|----------|-------|-------|---------|-------|--------|----------|------|-----|---------|--------|-------|--------|
| Record NEW cases of disease only | M | F | Total <5 | U5 Incia | M | F | to 17 | M | F | to 59 | М | F | 60 | total | cause | Incide |
| Upper Respiratory Tract Infection (URTI) | 3,860 | 3,142 | 7,002 | 872.1 | 2,579 | 2,409 | 4,988 | 2,018 | 2,515 | 4,533 | 95 | 155 | 250 | 16,773 | 32% | 459.6 |
| Influenza-like illness (ILI) | 951 | 783 | 1,734 | 216.0 | 924 | 861 | 1,785 | 734 | 839 | 1,573 | 36 | 41 | 77 | 5,169 | 10% | 141.6 |
| Lower Respiratory Tract Infection (LRTI) | 829 | 609 | 1,438 | 179.1 | 165 | 130 | 295 | 129 | 117 | 246 | 14 | 14 | 28 | 2,007 | 4% | 55.0 |
| Watery diarrhoea* | 323 | 295 | 618 | 77.0 | 44 | 44 | 88 | 69 | 87 | 156 | 6 | 8 | 14 | 876 | 2% | 24.0 |
| Bloody diarrhoea* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 0% | 0.0 |
| Skin infection | 308 | 290 | 598 | 74.5 | 231 | 318 | 549 | 251 | 301 | 552 | 17 | 14 | 31 | 1,730 | 3% | 47.4 |
| Eye infection | 185 | 149 | 334 | 41.6 | 66 | 52 | 118 | 146 | 138 | 284 | 12 | 10 | 22 | 758 | 1% | 20.8 |
| Ear infection | 324 | 234 | 558 | 69.5 | 195 | 218 | 413 | 214 | 359 | 573 | 12 | 10 | 22 | 1,566 | 3% | 42.9 |
| Dental Conditions | 63 | 113 | 176 | 21.9 | 656 | 660 | 1,316 | 1,554 | 1,825 | 3,379 | 64 | 65 | 129 | 5,000 | 9% | 137.0 |
| Intestinal Worms | 44 | 45 | 89 | 11.1 | 82 | 62 | 144 | 20 | 18 | 38 | 0 | 2 | 2 | 273 | 1% | 7.5 |
| Acute Jaundice Syndrome* | 17 | 13 | 30 | 3.7 | 11 | 10 | 21 | 2 | 1 | 3 | 0 | 0 | 0 | 54 | 0% | 1.5 |
| Measles (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Meningitis (suspected)* | 2 | 2 | 4 | 0.5 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 0% | 0.1 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Leishmaniasis | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Fever of unknown origin | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Other | 1,563 | 1,700 | 3,263 | 406.4 | 1,730 | 2,198 | 3,928 | 4,093 | 6,654 | 10,747 | 372 | 515 | 887 | 18,825 | 35% | 515.8 |
| Total | 8,469 | 7,375 | 15,844 | 1973.4 | 6,684 | 6,962 | 13,646 | 9,230 | 12,855 | 22,085 | 628 | 834 | 1,462 | 53,037 | 100% | 1453.3 |
| % by age group | | | 30% | | | | 26% | | | 42% | | | 3% | 100% | | |

^{*} indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to | 4 | Total < | 5 to | 17 | Total 5 | 18 1 | to 59 | Total 18 | > 6 | 0 | Total | Grand | % by |
|--|------|-----|---------|------|-----|---------|------|-------|----------|-----|-----|-------|-------|-------|
| Record NEW and REVISITS | М | F | 5 | M | F | to 17 | M | F | to 59 | M | F | > 60 | total | cause |
| Diabetes | 1 | 1 | 2 | 4 | 10 | 14 | 191 | 257 | 448 | 174 | 232 | 406 | 870 | 20% |
| Hypertension | 0 | 1 | 1 | 1 | 5 | 6 | 345 | 446 | 791 | 300 | 423 | 723 | 1,521 | 35% |
| Asthma | 171 | 111 | 282 | 131 | 72 | 203 | 73 | 102 | 175 | 19 | 15 | 34 | 694 | 16% |
| Ischaemic heart disease | 0 | 0 | 0 | 2 | 0 | 2 | 55 | 18 | 73 | 52 | 29 | 81 | 156 | 4% |
| Cardiovascular disease (other) | 0 | 1 | 1 | 0 | 6 | 6 | 35 | 33 | 68 | 27 | 31 | 58 | 133 | 3% |
| Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 1 | 0 | 1 | 23 | 7 | 30 | 24 | 14 | 38 | 69 | 2% |
| Liver disease | 0 | 1 | 1 | 1 | 0 | 1 | 0 | 1 | 1 | 0 | 1 | 1 | 4 | 0% |
| Thyroid problems | 3 | 2 | 5 | 8 | 6 | 14 | 23 | 156 | 179 | 3 | 16 | 19 | 217 | 5% |
| Cancer | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 2 | 3 | 3 | 6 | 8 | 0% |
| Other | 40 | 23 | 63 | 20 | 17 | 37 | 178 | 184 | 362 | 99 | 74 | 173 | 635 | 15% |
| Total | 215 | 140 | 355 | 168 | 116 | 284 | 925 | 1,204 | 2,129 | 701 | 838 | 1,539 | 4,307 | 100% |
| % by age group | | | 8% | | | 7% | | | 49% | | | 36% | 100% | |

3.4 Mental health

| | 0 to | 4 | Total < | 5 to | 17 | Total 5 | 18 t | o 59 | Total 18 | > 6 | 0 | Total | Grand | % by |
|--|------|----|---------|------|-----|---------|------|------|----------|-----|----|-------|-------|-------|
| | M | F | 5 | M | F | to 17 | M | F | to 59 | M | F | > 60 | total | cause |
| Epilepsy / seizures | 18 | 25 | 43 | 131 | 70 | 201 | 160 | 132 | 292 | 5 | 11 | 16 | 552 | 28% |
| Alcohol and other substance use disorder | 0 | 0 | 0 | 0 | 0 | 0 | 13 | 1 | 14 | 0 | 0 | 0 | 14 | 1% |
| Psychotic disorder (including mania) | 0 | 0 | 0 | 8 | 7 | 15 | 65 | 69 | 134 | 0 | 8 | 8 | 157 | 8% |
| Intellecutal disability | 0 | 2 | 2 | 34 | 6 | 40 | 22 | 12 | 34 | 4 | 4 | 8 | 84 | 4% |
| 5. Severe emotional disorder, including moderate-severe depression | 0 | 0 | 0 | 34 | 18 | 52 | 252 | 339 | 591 | 18 | 13 | 31 | 674 | 34% |
| Medically unexplained somatic complaint | 0 | 0 | 0 | 0 | 0 | 0 | 11 | 8 | 19 | 0 | 4 | 4 | 23 | 1% |
| Other psychological complaints | 1 | 0 | 1 | 198 | 102 | 300 | 67 | 77 | 144 | 7 | 5 | 12 | 457 | 23% |
| Total | 19 | 27 | 46 | 405 | 203 | 608 | 590 | 638 | 1,228 | 34 | 45 | 79 | 1,961 | 100% |
| % by age group | | | 2% | | | 31% | | | 63% | | | 4% | 100% | i |

3.5 Injuries

| | 0 to | 4 | Total < | 5 to | 17 | Total 5 | 18 t | o 59 | Total 18 | > 6 | 0 | Total | Grand | % by |
|--|------|-----|---------|------|-----|---------|------|------|----------|-----|----|-------|-------|-------|
| | M | F | 5 | M | F | to 17 | M | F | to 59 | M | F | > 60 | total | cause |
| Bites (all causes) | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% |
| 2. Injury (war related) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Injury (Burns) | 45 | 69 | 114 | 31 | 35 | 66 | 10 | 24 | 34 | 0 | 1 | 1 | 215 | 11% |
| Assault (with weapon non-war related) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 0% |
| Other | 263 | 251 | 514 | 316 | 217 | 533 | 333 | 218 | 551 | 36 | 35 | 71 | 1,669 | 88% |
| Total | 308 | 320 | 628 | 348 | 252 | 600 | 343 | 243 | 586 | 36 | 36 | 72 | 1,886 | 100% |
| % by age group | | | 33% | | | 32% | | | 31% | | | 4% | 100% | i |

3.6 Reproductive Tract Infections

| | | < ' | 18 | Total < | ≥ ' | 18 | Total ≥ | Grand | % by |
|----------|-------------------|-----|----|---------|-----|------|---------|-------|-------|
| | | м | F | 18 | М | F | 18 | total | cause |
| 1. Vagir | nal discharge | 0 | 19 | 19 | 0 | 543 | 543 | 562 | 49% |
| 2. Ureth | nral discharge | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 0% |
| 3. Geni | tal ulcer | 0 | 0 | 0 | 1 | 3 | 4 | 4 | 0% |
| 3. Lowe | er abdominal pain | 0 | 14 | 14 | 0 | 141 | 141 | 155 | 14% |
| Othe | r | 0 | 0 | 0 | 0 | 416 | 416 | 416 | 37% |
| Tota | I | 0 | 33 | 33 | 2 | 1103 | 1105 | 1138 | 100% |
| % by | age group | | | 3% | | | 97% | 100% | |

4.1 In-patient Department Activities

| | | 0 to | 0 to 4 To | | 5 to | 17 | Total 5 | 181 | to 59 | Total 18 | > 6 | 0 | Total | Grand |
|-----------------------------|--------------|------|-----------|-----|------|----|---------|-----|-------|----------|-----|---|-------|-------|
| | | М | F | 5 | M | F | to 17 | М | F | to 59 | M | F | > 60 | total |
| Number at beginning of repo | rting period | 3 | 3 | 6 | 0 | 1 | 1 | 0 | 3 | 3 | 0 | 0 | 0 | 10 |
| Total admissions | | 299 | 146 | 445 | 22 | 36 | 58 | 26 | 406 | 432 | 0 | 1 | 1 | 936 |
| | authorized | 286 | 138 | 424 | 21 | 34 | 55 | 26 | 363 | 389 | 0 | 1 | 1 | 869 |
| Number of exits | unauthorized | 2 | 3 | 5 | 0 | 1 | 1 | 0 | 43 | 43 | 0 | 0 | 0 | 49 |
| Number of exits | death | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | referral | 10 | 7 | 17 | 1 | 2 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 20 |
| Total exits | | 298 | 148 | 446 | 22 | 37 | 59 | 26 | 406 | 432 | 0 | 1 | 1 | 938 |
| Deaths within 24 hours | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| Number of days in reporting period | 91 |
|---|-------|
| Number of beds | 26 |
| Sum number of days stay for authorised discharges | 1,995 |
| Proportion of < 5 IPD deaths within 24 hours | |
| Average length of stay (days) | 2.3 |
| Bed occupancy | 91% |
| Hospitalisation rate (/1000 population/month) | 8.5 |

4.2 In-patient Admissions and Deaths

| | 0 to | 4 | I US CER I | 5 to | 17 | 5-17 | 181 | to 59 | 18-59 | > (| | Total adm | Total | Crude |
|--------------------------|------|--------|------------|------|--------|------|-----|--------|-------|-----|--------|-------------|--------|-------|
| | Adm | Deaths | | Adm | Deaths | CFR | Adm | Deaths | CFR | Adm | Deaths | Total autil | deaths | CFR |
| Paediatric medicine | 312 | 0 | 0% | 44 | 0 | 0% | 0 | 0 | | 0 | 0 | 356 | 0 | 0% |
| 2. General Surgery | 133 | 0 | 0% | 7 | 0 | 0% | 92 | 0 | 0% | 1 | 0 | 233 | 0 | 0% |
| Injury (non-war related) | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | 0 | 0 | |
| 4. Internal medicine | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | 0 | 0 | |
| 5. Obstetrics | 0 | 0 | | 7 | 0 | 0% | 335 | 0 | 0% | 0 | 0 | 342 | 0 | 0% |
| 6. Gynaecology | 0 | 0 | | 0 | 0 | | 5 | 0 | 0% | 0 | 0 | 5 | 0 | 0% |
| 7. Other | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | 0 | 0 | |
| Total | 445 | 0 | 0% | 58 | 0 | 0% | 432 | 0 | 0% | 1 | 0 | 936 | 0 | 0% |

Adm. = Admission; CFR = Case Fatality Rate

5.1 Referrals by location

| | | 0 to 4 | Total < | 5 to 17 | Total 5 to 17 | 18 to 59 | Total 18 to 59 | > 60 | Total | Grand total |
|----|-----------------------------|--------|---------|---------|------------------|----------|-------------------|------|-------|----------------|
| 1. | MOH Hospital | 28 | 28 | 64 | 64 | 78 | 78 | 5 | 5 | 175 |
| 2. | Jordan University Hospital | 243 | 243 | 209 | 209 | 230 | 230 | 39 | 39 | 721 |
| 3. | Private affiliated Hospital | 378 | 378 | 352 | 352 | 505 | 505 | 82 | 82 | 1,317 |
| 4. | Royal Medical Services | 0 | 0 | 2 | 2 | 2 | 2 | 0 | 0 | 4 |
| 5. | Other | 0 | 0 | 1 | 1 | 1 | 1 | 0 | 0 | 2 |

| Total number of reffered cases out side the camp | 2.219 |
|--|-------|

5.2 Referrals by service

| | | 0 to 4 | Total < | 5 to 17 | Total 5 to 17 | 18 to 59 | Total 18 to 59 | > 60 | Total | Grand total |
|----|-------------------|--------|---------|---------|------------------|----------|-------------------|------|-------|----------------|
| 1. | Paediatrics | 476 | 476 | 50 | 50 | 0 | 0 | 0 | 0 | 526 |
| 2. | Internal medicine | 17 | 17 | 193 | 193 | 260 | 260 | 59 | 59 | 529 |
| 3. | Obstetric | 0 | 0 | 3 | 3 | 30 | 30 | 0 | 0 | 33 |
| 4. | Gynaecological | 0 | 0 | 7 | 7 | 42 | 42 | 1 | 1 | 50 |
| 5. | Surgical | 78 | 78 | 253 | 253 | 295 | 295 | 39 | 39 | 665 |
| 6 | Other | 78 | 78 | 122 | 122 | 189 | 189 | 27 | 27 | 416 |

| | Total |
|--------------------------------------|--------|
| Number of laboratory tests conducted | 21,964 |

8.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|---------------------------------------|------|------|-------|
| First antenatal visit < 1st trimester | 20 | 411 | 431 |
| First antenatal visit > 1st trimester | 15 | 316 | 331 |

| ANC Indicators | |
|--|-----|
| Proportion of first time antenatal visits made < 1st trimester | 57% |

8.2 Antenatal care in facility

| A CONTRACTOR OF THE CONTRACTOR | | | | |
|--|-------------------|---|-----|-----|
| Number of property was as | ≥ 4 ANC visits | 4 | 222 | 226 |
| Number of pregnant women at time of delivery who received: | 2 TT doses | 3 | 61 | 64 |
| | anaemia screening | 7 | 359 | 366 |

| Coverage of complete antenatal care | 61% |
|--|-----|
| Coverage of antenatal tetanus immunisation | 17% |
| Coverage of anaemia screening in pregnancy | 99% |

8.3 Delivery care

| | < 18 | ≥ 18 | Total |
|---|------|------|-------|
| Live births | 7 | 361 | 368 |
| Still births | 0 | 3 | 3 |
| Low birth weight (<2500 g) | 0 | 14 | 14 |
| Attended by skilled health worker | 7 | 352 | 359 |
| Number of obstetric complications treated | 1 | 18 | 19 |
| Number of caesarean sections performed | 0 | 75 | 75 |

| Delivery Indicators | |
|--|-----|
| Crude birth rate (/1000 population/ month) | 3.4 |
| Proportion of births attended by skilled health worker (excl. TBA) | 98% |
| Proportion of low birth weight deliveries | 4% |
| Proportion of all births performed by caesarean section | 20% |
| Proportion of obsetric complications treated among girls < 18 | 14% |

8.4 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|-------|
| At least three PNC visits within 6 weeks | 8 | 207 | 215 |

| Postnatal Indicators | |
|-------------------------------------|-----|
| Coverage of complete postnatal care | 58% |

8.5 Family planning

| | New | | New Repeat | | Total |
|-----------------------------|------|------|------------|------|-------|
| | < 18 | ≥ 18 | < 18 | ≥ 18 | Total |
| Oral contraceptives | 18 | 311 | 12 | 964 | 1,305 |
| Emergency Contaceptive Pill | 0 | 0 | 0 | 0 | 0 |
| Injectable (Depo-Provera) | 1 | 27 | 0 | 49 | 77 |
| Implantable (Norplant) | 0 | 0 | 0 | 0 | 0 |
| Intra-Uterine Device (IUD) | 0 | 70 | 0 | 12 | 82 |
| Condom (Male) | 25 | 503 | 15 | 585 | 1,128 |
| Other | 0 | 0 | 0 | 0 | 0 |
| | | | | | |

| | < 18 | ≥ 18 |
|-----------------------------|------|-------|
| Family Planning Counseling* | 124 | 3,202 |
| | | |

8.6 Sexual and gender-based violence

| | < 18 | | 8 ≥ 18 | | Total |
|---|------|---|--------|---|-------|
| | М | F | M | F | Total |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | 0 | | 0 | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 5 | 5 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

| SGBV Indicators | |
|--|-----|
| Incidence of reported rape (/10,000 population/year) | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

Note

This report for non-camp Syrian refugees in Jordan is compiled from health data obtained from reporting agencies and clinics. Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Published on

Contact Information UNHCR Adam Musa Khalifa Email: KHALIFAA@unhcr.org

We gratefully acknowledge the contributions of the following agencies:











