

Population	0 to < 1		Total < 1	1 to 4		Total 1 to 4	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total
	M	F		M	F		M	F		M	F		M	F		
Refugee population	680	642	1,322	3,169	3,098	6,267	7,349	6,721	14,070	7,034	8,253	15,287	333	504	837	37,784
% by age group			4%			17%			37%			40%			2%	100%

**2.1 Mortality**

Record ONE cause of mortality per case	0 to 4		Total < 5	US % mort	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total	Crude % mort
	M	F			M	F		M	F		M	F			
1. Lower Respiratory Tract Infection (LRTI)	2	1	3	38%	0	0	0	0	0	0	2	1	3	6	24%
2. Watery diarrhoea	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%
3. Bloody diarrhoea	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%
4. Ischaemic heart disease	0	0	0	0%	0	0	0	1	0	1	1	0	1	2	8%
5. Cardiovascular disorders (other)	0	0	0	0%	0	0	0	1	2	3	1	1	2	5	20%
6. Cerebrovascular disease	0	1	1	13%	0	0	0	0	0	0	0	0	0	1	4%
7. Chronic kidney disease	0	1	1	13%	0	0	0	0	0	0	0	0	0	1	4%
8. Injury	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%
9. Cancer	0	0	0	0%	0	1	1	0	0	0	0	0	0	1	4%
10. Chronic obstructive pulmonary disease (COPD)	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%
11. Tuberculosis (confirmed)	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%
12. Neonatal death	1	1	2	25%	0	0	0	0	0	0	0	0	0	2	8%
13. COVID-19	0	0	0	0%	0	0	0	4	0	4	1	1	2	6	24%
14. Other	1	0	1	13%	0	0	0	0	0	0	0	0	0	1	4%
<b>Total</b>	<b>4</b>	<b>4</b>	<b>8</b>	<b>100%</b>	<b>0</b>	<b>1</b>	<b>1</b>	<b>6</b>	<b>2</b>	<b>8</b>	<b>5</b>	<b>3</b>	<b>8</b>	<b>25</b>	<b>100%</b>
% by age group			32%			4%			32%			32%		100%	

Mortality indicators	Male	Female	Total
Crude Mortality Rate (CMR) (/1000 population/month)	0.3	0.2	0.2
Under 5 Mortality Rate (U5MR) (/1000 population/month)	0.4	0.4	0.4
Neonatal Mortality Rate (NNMR) (/1000 population/month)			7.5

**2.2 Maternal death**

Number of maternal deaths reported	0
Number of maternal deaths investigated	0
Proportion of maternal deaths investigated	

**3.1 Staffing and service availability**

Number of full-time clinicians	28.3
Average number of full days on which each health facility was functioning	5.6
Total number of outpatient consultations	64,194
Health facility utilisation rate	6.9
Consultations per clinician per day	31

**3.2 Acute health conditions**

Record NEW cases of disease only	0 to 4		Total < 5	US Incid	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total	% by cause	Crude Incide
	M	F			M	F		M	F		M	F				
Upper Respiratory Tract Infection (URTI)	2,307	1,976	4,283	564.3	1,070	1,094	2,164	1,054	1,527	2,581	59	72	131	9,159	24%	242.4
Influenza-like illness (ILI)	357	310	667	87.9	104	113	217	210	215	425	6	10	16	1,325	3%	35.1
Lower Respiratory Tract Infection (LRTI)	346	246	592	78.0	39	27	66	38	40	78	6	13	19	755	2%	20.0
Watery diarrhoea*	156	159	315	41.5	59	57	116	41	75	116	3	5	8	555	1%	14.7
Bloody diarrhoea*	0	2	2	0.3	0	0	0	0	0	0	0	0	0	2	0%	0.1
Skin infection	170	204	374	49.3	174	284	458	157	260	417	10	7	17	1,266	3%	33.5
Eye infection	102	80	182	24.0	62	69	131	170	168	338	21	5	26	677	2%	17.9
Ear infection	185	155	340	44.8	91	126	217	191	387	578	23	27	50	1,185	3%	31.4
Dental Conditions	51	40	91	12.0	943	1,003	1,946	2,062	2,953	5,015	47	51	98	7,150	18%	189.2
Intestinal Worms	83	90	173	22.8	150	145	295	73	73	146	1	0	1	615	2%	16.3
Acute Jaundice Syndrome*	6	5	11	1.4	4	2	6	3	0	3	0	0	0	20	0%	0.5
Measles (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Meningitis (suspected)*	0	0	0	0.0	0	1	1	0	0	0	0	0	0	1	0%	0.0
Acute Flaccid Paralysis*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Leishmaniasis	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Fever of unknown origin	4	1	5	0.7	0	0	0	1	1	2	0	1	1	8	0%	0.2
Tuberculosis (suspected)	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
COVID-19 (suspected)	10	12	22	2.9	57	67	124	248	302	550	14	41	55	751	2%	19.9
COVID-19 (confirmed)	4	4	8	1.1	17	22	39	98	151	249	11	16	27	323	1%	8.5
Other	1,691	1,673	3,364	443.2	1,494	2,075	3,569	2,872	4,559	7,431	374	406	780	15,144	39%	400.8
<b>Total</b>	<b>5,472</b>	<b>4,957</b>	<b>10,429</b>	<b>1374.1</b>	<b>4,264</b>	<b>5,085</b>	<b>9,349</b>	<b>7,218</b>	<b>10,711</b>	<b>17,929</b>	<b>575</b>	<b>654</b>	<b>1,229</b>	<b>38,936</b>	<b>100%</b>	<b>1030.5</b>
% by age group			27%			24%			46%			3%	100%			

\* Indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

**3.3 Chronic health conditions**

Record NEW and REVISITS	0 to 4		Total < 5	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total	% by cause
	M	F		M	F		M	F		M	F			
Diabetes	0	0	0	11	21	32	179	218	397	95	142	237	666	19%
Hypertension	0	2	2	1	5	6	344	460	804	172	222	394	1,206	34%
Asthma	46	14	60	40	27	67	58	65	123	21	15	36	286	8%
Ischaemic heart disease	0	0	0	1	1	2	78	40	118	39	36	75	194	5%
Cardiovascular disease (other)	0	2	2	1	2	3	22	13	35	10	18	28	68	2%
Chronic obstructive pulmonary disease (COPD)	0	0	0	0	0	0	11	5	16	15	7	22	38	1%
Liver disease	0	0	0	0	0	0	0	2	2	3	0	3	5	0%
Thyroid problems	0	0	0	15	7	22	23	190	213	1	14	15	250	7%
Cancer	0	0	0	2	0	2	2	4	6	1	0	1	9	0%
Other	35	19	54	22	42	64	290	302	592	73	54	127	837	24%
<b>Total</b>	<b>81</b>	<b>37</b>	<b>118</b>	<b>92</b>	<b>105</b>	<b>197</b>	<b>1,007</b>	<b>1,299</b>	<b>2,306</b>	<b>430</b>	<b>508</b>	<b>938</b>	<b>3,559</b>	<b>100%</b>
% by age group			3%			6%			65%			26%	100%	



## 5.2 Referrals by service

		0 to 4	Total < 5	5 to 17	Total 5 to 17	18 to 59	Total 18 to 59	> 60	Total	Grand total
1.	Paediatrics	104	104	13	13	0	0	0	0	117
2.	Internal medicine	11	11	76	76	340	340	56	56	483
3.	Obstetric	0	0	1	1	60	60	0	0	61
4.	Gynaecological	0	0	5	5	6	6	0	0	11
5.	Surgical	8	8	70	70	32	32	8	8	118
6.	Other	2	2	12	12	16	16	20	20	50

	Total
Number of laboratory tests conducted	25,179

## 8.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester	8	448	456
First antenatal visit > 1st trimester	8	127	135

ANC Indicators	
Proportion of first time antenatal visits made < 1st trimester	77%

## 8.2 Antenatal care in facility

	≥ 4 ANC visits	2 TT doses	anaemia screening
Number of pregnant women at time of delivery who received:	5	2	7
	200	44	256
	205	46	263

Coverage of complete antenatal care	77%
Coverage of antenatal tetanus immunisation	17%
Coverage of anaemia screening in pregnancy	99%

## 8.3 Delivery care

	< 18	≥ 18	Total
Live births	7	258	265
Still births	0	2	2
Low birth weight (<2500 g)	0	12	12
Attended by skilled health worker	7	252	259
Number of obstetric complications treated	0	2	2
Number of caesarean sections performed	1	61	62

Delivery Indicators	
Crude birth rate (/1000 population/ month)	2.4
Proportion of births attended by skilled health worker (excl. TBA)	98%
Proportion of low birth weight deliveries	5%
Proportion of all births performed by caesarean section	23%
Proportion of obstetric complications treated among girls < 18	0%

## 8.4 Postnatal care

	< 18	≥ 18	Total
At least three PNC visits within 6 weeks	3	155	158

Postnatal Indicators	
Coverage of complete postnatal care	60%

## 8.5 Family planning

	New		Repeat		Total
	< 18	≥ 18	< 18	≥ 18	
Oral contraceptives	4	195	1	365	565
Emergency Contraceptive Pill	0	0	0	0	0
Injectable (Depo-Provera)	0	39	0	37	76
Implantable (Norplant)	0	0	0	0	0
Intra-Uterine Device (IUD)	0	14	0	0	14
Condom (Male)	2	149	2	120	273
Other	0	0	0	0	0

	< 18	≥ 18
Family Planning Counseling*	27	1,563

## 8.6 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No of rape survivors given PEP within 72 hours	0	0	0	0	0
No of female rape survivors given ECP within 120 hours	0	0	0	0	0
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV indicators	
Incidence of reported rape (/10,000 population/year)	0.0
Proportion of rape survivors who receive PEP < 72 hrs	
Proportion of female rape survivors who receive ECP < 120 hrs	
Proportion of rape survivors who receive STI presumptive treatment < 2 wks	

## Note

*This report for non-camp Syrian refugees in Jordan is compiled from health data obtained from reporting agencies and clinics. Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.*

Published on

Contact  
Information

UNHCR  
Adam Musa Khalifa  
Email: KHALIFAA@unhcr.org



We gratefully  
acknowledge the contributions of  
the following agencies:

