




Refugee population

Total refugee population:

116,429 \*

Demographic breakdown:



Key facts

Camp opened:  
July 2012

\* Source of population data:  
WFP Distribution Figures

Indicators at a glance

|   |     |   |
|---|-----|---|
| 1 Health utilisation rate                           | 6.2 | ✗ |
| 2 Number of consultations per clinician per day     | 39  | ✓ |
| 3 Hospitalisation rate                              | 4.0 | ⚠ |
| 4 Referral rate                                     |     | ? |
| 5 % deliveries attended by skilled personnel        | 84% | ✗ |
| 6 Crude Mortality Rate (CMR) (/1000/month)          | 0.2 | ✓ |
| 7 Under-five Mortality Rate (U5MR) (/1000/month)    | 0.4 | ✓ |
| 8 Neonatal Mortality Rate (NNMR) (/1000 livebirths) | 0.0 | ✓ |

Notes


This public health profile for Za'atri refugee camp is compiled from health facility data from the following partners:

- French Field Hospital
- Gynécologie Sans Frontières
- IMC Clinic
- JHAS Clinic
- MDM Clinic
- MSF Hospital
- Moroccan Field Hospital
- RMS Field Hospital
- Saudi clinic

Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible. However, at this early stage of implementation, indicators should be interpreted with care whilst the quality of data continues to be improved.

Morbidity

Acute health conditions (%)




Top 5 acute health conditions


| Condition                                  | % morbidity | # of cases |
|--|-------------|------------|
| 1 Upper Respiratory Tract Infection (URTI) | 24%         | 2,513      |
| 2 Influenza-like illness (ILI)             | 12%         | 1,218      |
| 3 Lower Respiratory Tract Infection (LRTI) | 9%          | 975        |
| 4 Watery diarrhoea*                        | 6%          | 598        |
| 5 Skin infection                           | 3%          | 283        |
| - Other                                    | 47%         | 4,902      |

Breakdown of acute health conditions


by age



by gender



Chronic health conditions (%)




Top 5 chronic health conditions


| Condition                  | % morbidity | # of cases |
|----------------------------|-------------|------------|
| 1 Hypertension             | 13%         | 371        |
| 2 Diabetes                 | 12%         | 339        |
| 3 Cardiovascular disorders | 9%          | 257        |
| 4 Asthma                   | 9%          | 243        |
| 5 Ischaemic heart disease  | 3%          | 79         |
| - Other                    | 54%         | 1,539      |

Breakdown of chronic health conditions

by age



by gender



UNHCR gratefully acknowledges the support of the following partners:




## Public Health

| Health Impact                                      | Indicator | Standard  |                |
|--|-----------|-----------|----------------|
| Crude Mortality Rate (CMR) (/1000/month)           | 0.2       | < 1.5     | ✓              |
| Under-five Mortality Rate (U5MR) (/1000/month)     | 0.4       | < 3.0     | ✓              |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths)  | 0.0       | < 40      | ✓              |
| <b>Access and utilization</b>                      |           |           |                |
| No. of health facilities                           | 6         | 1 : 19405 | 1 : < 10,000 ✗ |
| Total number of consultations                      | 13,813    |           |                |
| Health utilisation rate                            | 6.2       | 1 - 4     | ✗              |
| No. of consultations per trained clinician per day | 39        | < 50      | ✓              |


| Outbreak Alert and Response                 | Indicator | Standard |
|---|-----------|----------|
| Number of outbreaks reported                | 0         |          |
| % of outbreaks investigated within 48 hours |           | 100%     |
| <b>Laboratory</b>                           |           |          |
| Number of laboratory samples                | 678       |          |
| <b>Immunization</b>                         |           |          |
| Measles vaccination rate                    |           | > 95% ?  |
| Full vaccination rate                       |           | > 95% ?  |
| Program drop-out rate                       |           | < 10% ?  |
| Vaccine wastage rate                        |           | < 8% ?   |

### Breakdown of mental health conditions


By cause



By age




By gender




### Breakdown of injuries


By cause



By age



By gender



## In-patient and referral

| In-patient department                                | Indicator | Standard |
|--|-----------|----------|
| Hospitalisation rate (/1000/month)                   | 4.0       | 50 - 150 |
| Bed occupancy rate                                   | 161%      | > 75%    |
| Mean length of stay for authorised discharges (days) | 3.1       |          |

| Overall case fatality rate                        | Indicator | Standard |
|---|-----------|----------|
| Overall case fatality rate                        | 0%        |          |
| Proportion of deaths within 24 hours of admission | 0%        |          |
| <b>Referral</b>                                   |           |          |
| Total referral rate                               |           | ?        |

## Reproductive Health

| Antenatal care  | Indicator | Standard  |
|---|-----------|-----------|
| Coverage of antenatal care (4 or more visits)           | 0%        | ≥ 90% ✗   |
| Coverage of TT vaccination                              | 0%        | ≥ 90% ✗   |
| Coverage of anaemia screening in pregnancy              | 74%       | ≥ 90% ✗   |
| <b>Delivery care</b>                                    |           |           |
| Proportion of deliveries attended by skilled personnel  | 84%       | ≥ 90% ✗   |
| Proportion of deliveries performed by caesarean section | 23%       | 5 - 15% ✗ |
| Proportion of low birth weight deliveries (<2500g)      | 1%        | < 15% ✓   |
| <b>Maternal mortality</b>                               |           |           |
| Number of maternal deaths reported                      | 0         |           |
| % of maternal deaths investigated within 48 hours       |           | 100%      |

| Postnatal care   | Indicator | Standard |
|--|-----------|----------|
| Coverage of postnatal care                                   | 6%        | ≥ 90%    |
| <b>Family Planning</b>                                       |           |          |
| Contraceptive prevalence rate                                | 0%        |          |
| <b>Sexual and Gender-based Violence</b>                      |           |          |
| Incidence of rape (reported cases/10,000/year)*              | 0         |          |
| % of rape survivors who received PEP** < 72 hours            |           | 100%     |
| % of rape survivors who received ECP*** < 120 hours          |           | 100%     |
| % of rape survivors who received STI**** prophylaxis < 2 wks |           | 100%     |

\*Only includes rapes reported at health facilities \*\*Post-exposure prophylaxis \*\*\*Emergency contraceptive pill \*\*\*\*Sexually transmitted infection

