

1. Population

Population	0 to <1		Total <1	1 to 4		Total 1 to 4	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total
	M	F		M	F		M	F		M	F		M	F		
Refugee population	49	50	99	197	198	395	374	298	672	728	552	1,280	202	162	364	2,810
% by age group			4%			14%			24%			46%			13%	100%

2. Mortality

2.1 Mortality

Record ONE cause of mortality per case	0 to 4		Total <5	U5 % mort	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total > 50	Grand total	Crude % mort
	M	F			M	F		M	F		M	F			
1. Lower Respiratory Tract Infection (LRTI)	0	0	0		0	0	0	0	0	0	0	0	0	0	0%
2. Watery diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	0%
3. Bloody diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	0%
4. Cardiovascular disorders	0	0	0		0	0	0	1	0	1	0	0	0	1	100%
5. Ischaemic heart disease	0	0	0		0	0	0	0	0	0	0	0	0	0	0%
6. Injury	0	0	0		0	0	0	0	0	0	0	0	0	0	0%
7. Cancer	0	0	0		0	0	0	0	0	0	0	0	0	0	0%
8. Chronic obstructive pulmonary disease (COPD)	0	0	0		0	0	0	0	0	0	0	0	0	0	0%
9. Tuberculosis (confirmed)	0	0	0		0	0	0	0	0	0	0	0	0	0	0%
10. Neonatal death	0	0	0		0	0	0	0	0	0	0	0	0	0	0%
Other	0	0	0		0	0	0	0	0	0	0	0	0	0	0%
Total	0	0	0	100%	0	0	0	1	0	1	0	0	0	1	100%
% by age group			0%				0%			100%			0%	100%	

Mortality indicators	Male	Female	Total
Crude Mortality Rate (CMR)	2.6	0.0	1.4
Under 5 Mortality Rate (USMR)	0.0	0.0	0.0
Neonatal Mortality Rate (NNMR)			0.0

2.2 Maternal death

Number of maternal deaths reported	0
Number of maternal deaths investigated	0
Proportion of maternal deaths investigated	

3. Morbidity

3.1 Staffing and service availability

Number of full-time clinicians	2
Average number of full days on which each health facility was	6.0
Total number of outpatient consultations	555
Health facility utilisation rate	10.3
Consultations per clinician per day	46

3.2 Acute health conditions

Record NEW cases of disease only	0 to 4		Total <5	U5 Incid**	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause	Crude Incid**
	M	F			M	F		M	F		M	F				
Upper Respiratory Tract Infection (URTI)	15	16	31	62.8	20	25	45	19	26	45	7	9	16	137	26%	48.8
Lower Respiratory Tract Infection (LRTI)	4	3	7	14.2	17	13	30	7	6	13	4	2	6	56	11%	19.9
Watery diarrhoea*	4	5	9	18.2	6	2	8	2	2	4	3	3	6	27	5%	9.6
Bloody diarrhoea*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Malaria (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Skin infection	0	0	0	0.0	4	6	10	1	3	4	0	1	1	15	3%	5.3
Eye infection	0	1	1	2.0	2	1	3	0	2	2	2	1	3	9	2%	3.2
Ear infection	1	2	3	6.1	2	2	4	3	3	6	2	1	3	16	3%	5.7
Dental Conditions	1	1	2	4.0	7	12	19	1	3	4	0	0	0	25	5%	8.9
Intestinal Worms	0	0	0	0.0	1	1	2	3	0	3	0	0	0	5	1%	1.8
Acute Jaundice Syndrome*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Measles (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Meningitis (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute Flaccid Paralysis*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Tuberculosis (suspected)	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Fever of unknown origin	0	0	0	0.0	1	0	1	4	9	13	4	7	11	25	5%	8.9
Sexually Transmitted Infection (STI)	0	0	0	0.0	0	1	1	0	0	0	0	0	0	1	0%	0.4
Urinary tract infection (UTI)	1	2	3	6.1	1	2	3	0	2	2	0	0	0	8	2%	2.8
HIV/AIDS	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute malnutrition	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Leishmaniasis	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Other	3	3	6	12.1	57	70	127	17	33	50	5	10	15	198	38%	70.5
Total	29	33	62	125.5	118	135	253	57	89	146	27	34	61	522	100%	185.8
% by age group			12%				48%			28%			12%	100%		

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **incidence is cases per 1000 per week

3.3 Chronic health conditions

Record NEW and REVISITS	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
Diabetes	0	1	1	0	0	0	0	4	4	1	1	2	7	33%
Hypertension	0	1	1	0	1	1	0	1	1	0	1	1	4	19%
Asthma	1	0	1	1	2	3	1	1	2	1	2	3	9	43%
Ischaemic heart disease	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Cardiovascular disease (other)	0	0	0	0	0	0	0	0	0	0	1	1	1	5%
Chronic obstructive pulmonary disease (COPD)	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Liver disease	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Thyroid problems	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Cancer	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Total	1	2	3	1	3	4	1	6	7	2	5	7	21	100%
% by age group			14%			19%			33%			33%	100%	

3.4 Mental health

	0 to 4			5 to 14			15 to 49			≥ 50			Total	Grand total	% by cause
	M	F	Total < 5	M	F	Total 5 to 14	M	F	Total 15 to 49	M	F	Total ≥ 50			
1. Epilepsy / seizures	0	0	0	0	1	1	1	0	1	0	0	0	0	2	50%
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
3. Psychotic disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
4. Mental retardation / intellectual disability	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
5. Severe emotional disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
6. Medically unexplained somatic complaint	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	0	0	0	0	2	2	0	0	0	0	0	0	2	50%	
Total	0	0	0	0	3	3	1	0	1	0	0	0	4	100%	
% by age group	0%			75%			25%			0%			100%		

3.5 Injuries

	0 to 4			5 to 14			15 to 49			≥ 50			Total	Grand total	% by cause
	M	F	Total < 5	M	F	Total 5 to 14	M	F	Total 15 to 49	M	F	Total ≥ 50			
1. Injury (intentional)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
2. Injury (unintentional)	2	0	2	0	2	2	0	0	0	0	0	0	4	50%	
3. Burns	0	0	0	0	0	0	0	0	0	0	0	0	0	0%	
4. Accident	0	0	0	0	0	0	0	0	0	0	0	0	0	0%	
Other	0	0	0	0	3	3	0	0	0	1	0	1	4	50%	
Total	2	0	2	0	5	5	0	0	0	1	0	1	8	100%	
% by age group	25%			63%			0%			13%			100%		

4. Referrals

4.1 Referrals - general

	0 to 4	5 to 14	15 to 49	≥ 50	Grand total
Emergency	0	0	3	0	3
Elective	0	4	12	0	16
Other	0	0	0	0	0
Total	0	4	15	0	19
% by age group	0%	21%	79%	0%	100%

5. Laboratory & other investigations

	Total
Haematology	0
Imaging (X-ray, MRI/CT)	1
Other	18
Total	21

6. EPI and Vitamin A

	<1	≥1 to <5	Total
BCG	2	0	2
Hepatitis B birth dose	1	0	1
DPT I/HiB I/Hep B I* (Pentavalent I)	3	3	6
DPT II/HiB II* (Tetavalent)	0	0	0
DPT III/HiB III/Hep B II* (Pentavalent II)	2	0	2
Polio 0	1	0	1
Polio I	3	3	6
Polio II	0	0	0
Polio III	2	0	2
Measles	1	0	1
MMR	0	9	9
Rotavirus 1	2	0	2
Rotavirus 2	0	0	0
Rotavirus 3	0	0	0
Vitamin A	2	3	5
Additional vaccines	0	43	43

EPI Indicators (since December 2013)		
Total vaccinated this week		80
a. Program dropout rate*		14%
b. Immunisation coverage		
	BCG	63%
	Hepatitis B birth dose	47%
	DPT I	37%
	DPT III	32%
	Polio I	74%
	Polio III	26%
	Measles	69%
	MMR**	200%
	Rotavirus I	NR
	Rotavirus III	NR
c. Vitamin A coverage		NR

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%
**For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.
NAD=No adequate data; NR=Not consistently reported

7. Reproductive Health

7.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester	3	4	7
First antenatal visit > 1st trimester	0	5	5
Number of pregnant women at time of delivery who received:			
≥ 4 ANC visits	0	0	0
2 TT doses	0	0	0
anaemia screening	0	0	0

ANC Indicators	
Proportion of first time antenatal visits made < 1st trimester	58%

7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery	0	0	0

7.3 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No of rape survivors given PEP within 72 hours	0	0	0	0	0
No of female rape survivors given ECP within 120 hours	0	0	0	0	0
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators		
Incidence of reported rape		0.0
Proportion of rape survivors who receive PEP < 72 hrs		
Proportion of female rape survivors who receive ECP < 120 hrs		
Proportion of rape survivors who receive STI presumptive treatment < 2 wks		

Data Sources

1. Health Information System

Publication date:

14 March 2014

Contact Information

Dohuk
Dr Bakhtiyar AHMED, DOH Duhok

Email: BAHMED@duhokhealth.org

Erbil
Mohammed Marzooq, UNHCR

Email: MARZOOG@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

