

1. Population

Population	0 to <1		Total <1	1 to 4		Total 1 to 4	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total
	M	F		M	F		M	F		M	F		M	F		
Refugee population	254	242	496	1,015	968	1,983	1,596	1,571	3,167	3,058	2,890	5,948	890	848	1,738	13,332
% by age group			4%			15%			24%			45%			13%	100%

* estimates from registered population

2. Mortality

2.1 Mortality

Record ONE cause of mortality per case	0 to 4		Total <5	U5 % mort	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total > 50	Grand total	Crude % mort
	M	F			M	F		M	F		M	F			
1. Lower Respiratory Tract Infection (LRTI)	0	0	0		0	0	0	0	0	0	0	0	0	0	
2. Watery diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	
3. Bloody diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	
4. Cardiovascular disorders	0	0	0		0	0	0	0	0	0	0	0	0	0	
5. Ischaemic heart disease	0	0	0		0	0	0	0	0	0	0	0	0	0	
6. Injury	0	0	0		0	0	0	0	0	0	0	0	0	0	
7. Cancer	0	0	0		0	0	0	0	0	0	0	0	0	0	
8. Chronic obstructive pulmonary disease (COPD)	0	0	0		0	0	0	0	0	0	0	0	0	0	
9. Tuberculosis (confirmed)	0	0	0		0	0	0	0	0	0	0	0	0	0	
10. Neonatal death	0	0	0		0	0	0	0	0	0	0	0	0	0	
Other	0	0	0		0	0	0	0	0	0	0	0	0	0	
Total	0	0	0	100%	0	0	0	0	0	0	0	0	0	0	100%
% by age group															

Mortality indicators

	Male	Female	Total
Crude Mortality Rate (CMR)	0.0	0.0	0.0
Under 5 Mortality Rate (USMR)	0.0	0.0	0.0
Neonatal Mortality Rate (NNMR)			0.0

2.2 Maternal death

Number of maternal deaths reported	0
Number of maternal deaths investigated	0
Proportion of maternal deaths investigated	

3. Morbidity

3.1 Staffing and service availability

Number of full-time clinicians	4
Average number of full days on which each health facility was	6.0
Total number of outpatient consultations	1,207
Health facility utilisation rate	4.7
Consultations per clinician per day	50

3.2 Acute health conditions

Record NEW cases of disease only	0 to 4		Total <5	U5 Incid**	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause	Crude Incid**
	M	F			M	F		M	F		M	F				
Upper Respiratory Tract Infection (URTI)	71	74	145	58.5	46	37	83	63	64	127	4	10	14	369	35%	27.7
Lower Respiratory Tract Infection (LRTI)	29	20	49	19.8	1	2	3	4	4	8	0	2	2	62	6%	4.7
Watery diarrhoea*	12	7	19	7.7	3	3	6	5	2	7	0	0	0	32	3%	2.4
Bloody diarrhoea*	1	1	2	0.8	0	1	1	0	0	0	1	0	1	4	0%	0.3
Malaria (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Skin infection	15	4	19	7.7	10	10	20	17	22	39	2	0	2	80	8%	6.0
Eye infection	9	4	13	5.2	5	4	9	8	15	23	1	1	2	47	4%	3.5
Ear infection	14	11	25	10.1	6	3	9	13	16	29	0	1	1	64	6%	4.8
Dental Conditions	0	1	1	0.4	3	7	10	3	1	4	1	0	1	16	2%	1.2
Intestinal Worms	1	3	4	1.6	7	3	10	1	1	2	0	0	0	16	2%	1.2
Acute Jaundice Syndrome*	2	1	3	1.2	3	2	5	0	0	0	0	0	0	8	1%	0.6
Measles (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Meningitis (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute Flaccid Paralysis*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Tuberculosis (suspected)	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Fever of unknown origin	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Sexually Transmitted Infection (STI)	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Urinary tract infection (UTI)	2	12	14	5.6	1	6	7	17	25	42	1	4	5	68	6%	5.1
HIV/AIDS	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute malnutrition	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Leishmaniasis	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Other	23	12	35	14.1	20	9	29	70	152	222	6	8	14	300	28%	22.5
Total	179	150	329	132.7	105	87	192	201	302	503	16	26	42	1066	100%	80.0
% by age group			31%				18%			47%			4%	100%		

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **incidence is cases per 1000 per week

3.3 Chronic health conditions

Record NEW and REVISITS	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause	
	M	F		M	F		M	F		M	F				
Diabetes	0	0	0	0	0	0	9	4	13	6	2	8	21	24%	
Hypertension	0	0	0	0	0	0	7	12	19	2	8	10	29	33%	
Asthma	0	1	1	0	1	1	0	1	1	0	1	1	4	5%	
Ischaemic heart disease	0	0	0	0	0	0	3	3	6	2	0	2	8	9%	
Cardiovascular disease (other)	1	2	3	1	1	2	1	5	6	1	2	3	14	16%	
Chronic obstructive pulmonary disease (COPD)	0	0	0	0	0	0	0	0	0	0	0	0	0	0%	
Liver disease	0	0	0	0	0	0	0	0	0	0	0	0	0	0%	
Thyroid problems	0	2	2	0	2	2	2	1	3	0	2	2	9	10%	
Cancer	0	0	0	0	0	0	1	2	3	0	0	0	3	3%	
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0%	
Total	1	5	6	1	4	5	23	28	51	11	15	26	88	100%	
% by age group			7%			6%			58%			30%	100%		

3.4 Mental health

	0 to 4			5 to 14			15 to 49			≥ 50			Total ≥ 50	Grand total	% by cause
	M	F	Total < 5	M	F	Total 5 to 14	M	F	Total 15 to 49	M	F	Total ≥ 50			
1. Epilepsy / seizures	0	0	0	0	0	0	1	1	2	0	0	0	2	2	10%
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
3. Psychotic disorder	0	0	0	0	0	0	1	1	2	0	0	0	2	2	10%
4. Mental retardation / intellectual disability	0	0	0	1	0	1	0	0	0	0	0	0	1	1	5%
5. Severe emotional disorder	0	0	0	0	0	0	4	4	8	0	0	0	8	8	38%
6. Medically unexplained somatic complaint	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	0	1	1	0	1	1	5	1	6	0	0	0	6	6	27%
Total	0	1	1	1	1	2	11	7	18	0	0	0	21	21	100%
% by age group			5%			10%			86%				0%	100%	

3.5 Injuries

	0 to 4			5 to 14			15 to 49			≥ 50			Total ≥ 50	Grand total	% by cause
	M	F	Total < 5	M	F	Total 5 to 14	M	F	Total 15 to 49	M	F	Total ≥ 50			
1. Injury (intentional)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
2. Injury (unintentional)	0	0	0	5	3	8	10	3	13	0	0	0	21	21	66%
3. Burns	0	0	0	0	0	0	0	2	2	0	0	0	2	2	6%
4. Accident	2	1	3	2	0	2	0	0	0	0	0	0	5	5	16%
Other	0	0	0	1	0	1	2	1	3	0	0	0	4	4	13%
Total	2	1	3	8	3	11	12	6	18	0	0	0	32	32	100%
% by age group			9%			34%			56%				0%	100%	

4. Referrals

4.1 Referrals - general

	0 to 4	5 to 14	15 to 49	≥ 50	Grand total
Emergency	2	2	7	1	12
Elective	3	1	19	3	26
Other	0	0	0	0	0
Total	5	3	26	4	38
% by age group	13%	8%	68%	11%	100%

ND: no data submitted

5. Laboratory & other investigations

	Total
Haematology	0
Imaging (X-ray, MRI/CT)	0
Other	0
Total	0

6. EPI and Vitamin A

	<1	1 to <5	Total
BCG	0	0	0
Hepatitis B birth dose	0	0	0
DPT I/HiB I/Hep B I* (Pentavalent I)	2	0	2
DPT II/HiB II* (Tetravalent)	2	0	2
DPT III/HiB III/Hep B II* (Pentavalent II)	7	0	7
Polio 0	0	0	0
Polio I	2	0	2
Polio II	2	0	2
Polio III	7	0	7
Measles	5	0	5
MMR	0	5	5
Rotavirus 1	2	0	2
Rotavirus 2	2	0	2
Rotavirus 3	7	0	7
Vitamin A	9	0	9
Additional vaccines	0	8	8

EPI Indicators (since December 2013)		
Total vaccinated this week		60
a. Program dropout rate*		-22%
b. Immunisation coverage	BCG	52%
	Hepatitis B birth dose	52%
	DPT I	86%
	DPT III	105%
	Polio I	86%
	Polio III	105%
c. Vitamin A coverage	Measles	107%
	MMR**	129%
	Rotavirus I	86%
	Rotavirus III	105%

*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%
**For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

7. Reproductive Health

7.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester			
First antenatal visit > 1st trimester			
Number of pregnant women at time of delivery who received:	NO DATA		
≥ 4 ANC visits			
2 TT doses			
anaemia screening			

ND: no data submitted

ANC Indicators	
Proportion of first time antenatal visits made < 1st trimester	

7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery			NO DATA

ND: no data submitted

7.3 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No of rape survivors given PEP within 72 hours	0	0	0	0	0
No of female rape survivors given ECP within 120 hours					0
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators		
Incidence of reported rape		0.0
Proportion of rape survivors who receive PEP < 72 hrs		
Proportion of female rape survivors who receive ECP < 120 hrs		
Proportion of rape survivors who receive STI presumptive treatment < 2 wks		

Data Sources: 1. Health Information System

Publication date: 24 March 2014

Contact Information: Erbil, Dr Chimmen Akrawi, DOH Erbil

Email: chimmentya@yahoo.com

Erbil, Mohammed Marzooq, UNHCR

Email: MARZOOQ@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

