



1.0 Population

This section was not collected

2.0 Mortality

2.1 Mortality

This section was not collected

2.2 Maternal Death

This section was not collected

3.0 Out-patient Department

3.1 Staffing and service availability

This section was not collected

3.2 Acute health conditions

| Record NEW cases of disease only | 0 - 4 | | Total 0 - 4 | U5% by cause | 5 - 17 | | Total 5 - 17 | 18 - 59 | | Total 18 - 59 | > 60 | | Total > 60 | Grand Total | % by cause |
|--|------------|------------|--------------|--------------|------------|------------|--------------|------------|------------|---------------|-----------|-----------|------------|--------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| Ear infection | 26 | 24 | 50 | 4 % | 6 | 16 | 22 | 13 | 22 | 35 | 0 | 1 | 1 | 108 | 3 % |
| Acute Jaundice Syndrome* | 0 | 0 | 0 | 0 % | 2 | 2 | 4 | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 0 % |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0 % | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 % |
| Measles* | 1 | 0 | 1 | 0 % | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 % |
| Prematurity | 0 | 0 | 0 | 0 % | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 0 % |
| Intestinal Worms | 6 | 2 | 8 | 1 % | 6 | 2 | 8 | 0 | 3 | 3 | 0 | 0 | 0 | 19 | 1 % |
| Meningitis* | 0 | 0 | 0 | 0 % | 1 | 3 | 4 | 1 | 3 | 4 | 0 | 1 | 1 | 9 | 0 % |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0 % | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 % |
| Eye infection | 3 | 3 | 6 | 1 % | 7 | 7 | 14 | 12 | 23 | 35 | 0 | 3 | 3 | 58 | 2 % |
| Fever of unknown origin | 15 | 7 | 22 | 2 % | 4 | 1 | 5 | 0 | 5 | 5 | 0 | 0 | 0 | 32 | 1 % |
| Neonatal sepsis | 0 | 0 | 0 | 0 % | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 % |
| Skin infection | 1 | 1 | 2 | 0 % | 2 | 1 | 3 | 0 | 3 | 3 | 0 | 0 | 0 | 8 | 0 % |
| Urinary tract infection (UTI) | 2 | 3 | 5 | 0 % | 3 | 7 | 10 | 9 | 39 | 48 | 0 | 4 | 4 | 67 | 2 % |
| Sexually Transmitted Infection (STI) | 0 | 0 | 0 | 0 % | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 % |
| Upper Respiratory Tract Infection (URTI) | 160 | 136 | 296 | 26 % | 123 | 108 | 231 | 45 | 62 | 107 | 2 | 6 | 8 | 642 | 19 % |
| Watery diarrhoea* | 4 | 5 | 9 | 1 % | 2 | 3 | 5 | 0 | 3 | 3 | 0 | 0 | 0 | 17 | 1 % |
| Lower Respiratory Tract Infection (LRTI) | 226 | 176 | 402 | 36 % | 146 | 126 | 272 | 62 | 110 | 172 | 7 | 13 | 20 | 866 | 26 % |
| Scabies | 6 | 2 | 8 | 1 % | 16 | 20 | 36 | 8 | 8 | 16 | 1 | 1 | 2 | 62 | 2 % |
| Other | 161 | 156 | 317 | 28 % | 96 | 141 | 237 | 108 | 712 | 820 | 20 | 41 | 61 | 1,435 | 43 % |
| Bloody diarrhoea* | 0 | 0 | 0 | 0 % | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 0 % |
| Total | 611 | 515 | 1,126 | 100 % | 415 | 437 | 852 | 258 | 995 | 1,253 | 30 | 70 | 100 | 3,331 | 100 % |
| % by age group | | | 34 % | | | | 26 % | | | 38 % | | | 3 % | 100 % | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 - 4 | | Total < 5 | 5 - 17 | | Total 5 - 17 | 18 - 59 | | Total 18 - 59 | > 60 | | Total > 60 | Grand Total | % by cause |
|----------------------------------|-------|----|-----------|--------|------|--------------|---------|-----|---------------|------|-------|------------|-------------|------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| Asthma/COPD | 64 | 66 | 130 | 39 | 32 | 71 | 18 | 39 | 57 | 6 | 4 | 10 | 268 | 50 % |
| Ischaemic heart disease | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 2 | 0 | 0 | 0 | 2 | 0 % |
| Renal disease | 1 | 0 | 1 | 3 | 1 | 4 | 0 | 4 | 4 | 0 | 1 | 1 | 10 | 2 % |
| Other | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 6 | 6 | 2 | 0 | 2 | 10 | 2 % |
| Hypertension | 0 | 0 | 0 | 0 | 0 | 0 | 27 | 34 | 61 | 26 | 33 | 59 | 120 | 22 % |
| Liver disease | 0 | 0 | 0 | 2 | 0 | 2 | 0 | 1 | 1 | 0 | 0 | 0 | 3 | 1 % |
| Cardiovascular disorders (other) | 0 | 0 | 0 | 0 | 0 | 0 | 6 | 4 | 10 | 1 | 6 | 7 | 17 | 3 % |
| Cancer and other neoplasms | 1 | 1 | 2 | 0 | 2 | 2 | 3 | 4 | 7 | 0 | 1 | 1 | 12 | 2 % |
| Diabetes | 0 | 0 | 0 | 2 | 8 | 10 | 17 | 33 | 50 | 9 | 24 | 33 | 93 | 17 % |
| Total | 66 | 67 | 133 | 46 | 45 | 91 | 72 | 126 | 198 | 44 | 69 | 113 | 535 | 100 % |
| % by age group | | | 25 % | | 17 % | | 37 % | | 21 % | | 100 % | | | |

3.4 Mental Health

| | 0 - 4 | | Total < 5 | 5 - 17 | | Total 5 - 17 | 18 - 59 | | Total 18 - 59 | > 60 | | Total > 60 | Grand Total | % by cause |
|--|-------|---|-----------|--------|------|--------------|---------|----|---------------|------|-------|------------|-------------|------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| Severe emotional disorder | 1 | 2 | 3 | 2 | 8 | 10 | 5 | 9 | 14 | 0 | 0 | 0 | 27 | 42 % |
| Psychotic disorder | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 2 | 3 % |
| Epilepsy / seizures | 1 | 0 | 1 | 3 | 2 | 5 | 6 | 0 | 6 | 0 | 0 | 0 | 12 | 18 % |
| Other | 5 | 1 | 6 | 4 | 1 | 5 | 3 | 0 | 3 | 0 | 0 | 0 | 14 | 22 % |
| Alcohol and other substance abuse disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 % |
| Mental retardation / intellectual disability | 4 | 1 | 5 | 2 | 0 | 2 | 1 | 1 | 2 | 0 | 0 | 0 | 9 | 14 % |
| Medically unexplained somatic complaint | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2 % |
| Total | 12 | 4 | 16 | 12 | 11 | 23 | 16 | 10 | 26 | 0 | 0 | 0 | 65 | 100 % |
| % by age group | | | 25 % | | 35 % | | 40 % | | 0 % | | 100 % | | | |

3.5 Injuries

| | 0 - 4 | | Total < 5 | 5 - 17 | | Total 5 - 17 | 18 - 59 | | Total 18 - 59 | > 60 | | Total > 60 | Grand Total | % by cause |
|------------------------|-------|---|-----------|--------|------|--------------|---------|---|---------------|------|-------|------------|-------------|------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| Injury (burns) | 1 | 3 | 4 | 1 | 1 | 2 | 0 | 1 | 1 | 0 | 0 | 0 | 7 | 23 % |
| Injury (unintentional) | 0 | 1 | 1 | 0 | 1 | 1 | 3 | 1 | 4 | 0 | 1 | 1 | 7 | 23 % |
| Accident | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 % |
| Other | 1 | 0 | 1 | 1 | 0 | 1 | 9 | 6 | 15 | 0 | 0 | 0 | 17 | 55 % |
| Injury (intentional) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 % |
| Total | 2 | 4 | 6 | 2 | 2 | 4 | 12 | 8 | 20 | 0 | 1 | 1 | 31 | 100 % |
| % by age group | | | 19 % | | 13 % | | 65 % | | 3 % | | 100 % | | | |

4.0 In-patient Department

4.1 In-patient department activities

This section was not collected

4.2 In-patient admissions and deaths

This section was not collected

5.0 Referral

5.1 Referrals by location

This section was not collected

5.2 Referrals by service

This section was not collected

6.0 Laboratory

This section was not collected

7.0 EPI and Vitamin A

7.1 Vaccination

This section was not collected

7.2 Vaccine wastage

This section was not collected

8.0 Reproductive Health

8.1 Antenatal care

This section was not collected

8.2 Delivery care

This section was not collected

8.3 Postnatal care

| | < 18 | > 18 | Total |
|--|------|------|-------|
| Women who attended 3 postnatal visits within 6 wks of delivery | | | 3 |

Postnatal Indicators

| | |
|-------------------------------------|-----|
| Coverage of complete postnatal care | 0 % |
|-------------------------------------|-----|

8.4 Family planning

This section was not collected

8.5 Sexual and gender-based violence

This section was not collected

9.0 HIV/AIDS

9.1 Voluntary counselling and testing (VCT)

This section was not collected

9.2 Prevention of mother-to-child transmission (PMTCT)

This section was not collected

