Camps reporting this week: Kawergosk, Qushtapa, Basirma and Darashakran

1. Population

| Population | 0 to |) < 1 F | Total <1 | 1 to | o 4 F | Total 1 to 4 | 5 to | 14 F | Total 5 to 14 | 15 to | o 49 F | Total 15 to 49 | ≥ 5 M | 60 F | Total ≥ 50 | Grand total |
|---------------------------------------|------|------------|-------------|-------|----------|-----------------|-------|---------|------------------|-------|-----------|----------------------|----------|---------|---------------|----------------|
| Refugee population of reporting camps | 481 | 474 | 955 | 1,925 | 1,893 | 3,818 | 3,512 | 3,126 | 6,638 | 6,249 | 5,807 | 12,056 | 1,808 | 1,753 | 3,561 | 27,028 |
| % by age group | | | 4% | , | | 14% | | | 25% | | | 45% | | | 13% | 100% |

^{*} estimates from registered population in Kawergosk, Qushtapa, Basirma and Darashakran camps

2. Mortality

2.1 Mortality

| | December 1 ONE and a few stall for many | | o 4 | Total | U5 % | 5 to | 14 | Total 5 | 15 to | o 49 | Total 15 | ≥ ! | 50 | Total > | Grand | Crude % |
|-----|--|---|-----|-------|------|------|----|---------|-------|------|----------|-----|----|---------|-------|---------|
| | Record ONE cause of mortality per case | М | F | <5 | mort | М | F | to 14 | М | F | to 49 | М | F | 50 | total | mort |
| 1. | Lower Respiratory Tract Infection (LRTI) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 2. | Watery diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 3. | Bloody diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 4. | Cardiovascular disorders | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 5. | Ischaemic heart disease | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 6. | Injury | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 7. | Cancer | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 8. | Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 9. | Tuberculosis (confirmed) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 10. | Neonatal death | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | Other | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | Total | 0 | 0 | 0 | 100% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100% |
| | % by age group | | | | | | | | | | | | | | 100% | |

| Mortality indicators | Male | Female | Total |
|--------------------------------|------|--------|-------|
| Crude Mortality Rate (CMR) | 0.0 | 0.0 | 0.0 |
| Under 5 Mortality Rate (U5MR) | 0.0 | 0.0 | 0.0 |
| Neonatal Mortality Rate (NNMR) | | | 0.0 |

2.2 Maternal death

| Number of maternal deaths reported | 0 |
|--|---|
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3. Morbidity

3.1 Staffing and service availability

| Number of full-time clinicians | 11 |
|---|-------|
| Average number of full days on which each health facility was | 6.3 |
| Total number of outpatient consultations | 3,705 |
| Health facility utilisation rate | 7.1 |
| Consultations per clinician per day | 53 |

3.2 Acute health conditions

| | 0 to | o 4 | Total | U5 | 5 to | 14 | Total 5 | 15 to | 49 | Total 15 | ≥ 5 | 0 | Total ≥ | Grand | % bv | Crude |
|--|------|-----|-------|---------|------|-----|---------|-------|------|----------|-----|-----|---------|-------|-------|---------|
| Record NEW cases of disease only | M | F | <5 | Incid** | M | F | to 14 | M | F | to 49 | M | F | 50 | total | cause | Incid** |
| Upper Respiratory Tract Infection (URTI) | 199 | 158 | 357 | 74.8 | 131 | 107 | 238 | 244 | 232 | 476 | 21 | 23 | 44 | 1115 | 34% | 41.3 |
| Lower Respiratory Tract Infection (LRTI) | 133 | 114 | 247 | 51.7 | 47 | 61 | 108 | 36 | 53 | 89 | 9 | 8 | 17 | 461 | 14% | 17.1 |
| Watery diarrhoea* | 21 | 13 | 34 | 7.1 | 5 | 3 | 8 | 10 | 6 | 16 | 0 | 1 | 1 | 59 | 2% | 2.2 |
| Bloody diarrhoea* | 4 | 3 | 7 | 1.5 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 8 | 0% | 0.3 |
| Malaria (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Skin infection | 30 | 22 | 52 | 10.9 | 15 | 28 | 43 | 30 | 38 | 68 | 5 | 3 | 8 | 171 | 5% | 6.3 |
| Eye infection | 9 | 11 | 20 | 4.2 | 4 | 7 | 11 | 13 | 18 | 31 | 3 | 4 | 7 | 69 | 2% | 2.6 |
| Ear infection | 16 | 7 | 23 | 4.8 | 8 | 5 | 13 | 11 | 23 | 34 | 1 | 2 | 3 | 73 | 2% | 2.7 |
| Dental Conditions | 3 | 5 | 8 | 1.7 | 28 | 27 | 55 | 72 | 85 | 157 | 14 | 10 | 24 | 244 | 7% | 9.0 |
| Intestinal Worms | 1 | 4 | 5 | 1.0 | 3 | 4 | 7 | 0 | 1 | 1 | 0 | 0 | 0 | 13 | 0% | 0.5 |
| Acute Jaundice Syndrome* | 1 | 1 | 2 | 0.4 | 3 | 1 | 4 | 0 | 1 | 1 | 0 | 0 | 0 | 7 | 0% | 0.3 |
| Measles (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Meningitis (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Fever of unknown origin | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Sexually Transmitted Infection (STI) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 190 | 190 | 0 | 1 | 1 | 191 | 6% | 7.1 |
| Urinary tract infection (UTI) | 6 | 11 | 17 | 3.6 | 3 | 18 | 21 | 24 | 74 | 98 | 2 | 4 | 6 | 142 | 4% | 5.3 |
| HIV/AIDS | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute malnutrition | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Leishmaniasis | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Other | 30 | 17 | 47 | 9.8 | 33 | 38 | 71 | 171 | 350 | 521 | 25 | 55 | 80 | 719 | 22% | 26.6 |
| Total | 453 | 366 | 819 | 171.6 | 280 | 299 | 579 | 611 | 1072 | 1683 | 80 | 111 | 191 | 3272 | 100% | 121.1 |
| % by age group | | | 25% | | | | 18% | | | 51% | | | 6% | 100% | | |

^{*} indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **incidence is cases per 1000 per week

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 1 | to 4 | Total < | 5 to | 14 | Total 5 | 15 to | 49 | Total 15 | Total 15 ≥ 50 | | Total | Grand | % by |
|--|-----|------|---------|------|----|---------|-------|-----|----------|---------------|----|-------|-------|-------|
| Record NEW and REVISITS | M | F | 5 | М | F | to 14 | M | F | to 49 | М | F | ≥ 50 | total | cause |
| Diabetes | 0 | 0 | 0 | 3 | 0 | 3 | 15 | 14 | 29 | 9 | 19 | 28 | 60 | 18% |
| Hypertension | 0 | 0 | 0 | 0 | 0 | 0 | 12 | 20 | 32 | 20 | 34 | | | 26% |
| Asthma | 2 | 3 | 5 | 0 | 0 | 0 | 1 | 4 | 5 | 3 | 5 | 8 | 18 | 5% |
| Ischaemic heart disease | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 4 | 5 | 0 | 6 | 6 | 11 | 3% |
| Cardiovascular disease (other) | 0 | 1 | 1 | 0 | 0 | 0 | 6 | 4 | 10 | 5 | 7 | 12 | 23 | 7% |
| Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Liver disease | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Thyroid problems | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 1 | 1 | 2 | 1% |
| Cancer | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 3 | 3 | 1 | 0 | 1 | 5 | 2% |
| Other | 0 | 1 | 1 | 3 | 1 | 4 | 30 | 73 | 103 | 11 | 9 | 20 | 128 | 38% |
| Total | 2 | 5 | 7 | 6 | 2 | 8 | 65 | 123 | 188 | 49 | 81 | 130 | 333 | 100% |
| % by age group | | | 2% | | | 2% | | | 56% | | | 39% | 100% | |

3.4 Mental health

| | 0 t | o 4 | Total < | 5 to | 14 | Total 5 | 15 to | o 49 | Total | ≥ (| 50 | Total | Grand | % by |
|--|-----|-----|---------|------|----|---------|-------|------|-------------|-----|----|-------|-------|-------|
| | М | F | 5 | M | F | to 14 | M | F | 15 to 49 | M | F | ≥ 50 | total | cause |
| Epilepsy / seizures | 0 | 0 | 0 | 1 | 0 | 1 | 5 | 1 | 6 | 0 | 0 | 0 | 7 | 21% |
| Alcohol and other substance abuse disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Psychotic disorder | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 3% |
| Mental retardation / intellectual disability | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 3% |
| Severe emotional disorder | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 3 | 4 | 0 | 0 | 0 | 4 | 12% |
| Medically unexplained somatic complaint | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 0 | 0 | 2 | 6% |
| Other | 0 | 1 | 1 | 0 | 3 | 3 | 3 | 11 | 14 | 0 | 0 | 0 | 18 | 55% |
| Total | 0 | 1 | 1 | 1 | 3 | 4 | 10 | 18 | 28 | 0 | 0 | 0 | 33 | 100% |
| % by age group | | | 3% | | | 12% | | | 85% | | | 0% | 100% | |

3.5 Injuries

| | | 0 to | o 4 | Total < | 5 to | 14 | Total 5 | 15 t | o 49 | Total | ≥ | 50 | Total | Grand | % by |
|-------|-----------------------|------|-----|---------|------|----|---------|------|------|-------------|---|----|-------|-------|-------|
| | | M | F | 5 | М | F | to 14 | M | F | 15 to 49 | M | F | ≥ 50 | total | cause |
| 1. lt | njury (intentional) | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 2 | 3% |
| 2. lt | njury (unintentional) | 6 | 4 | 10 | 5 | 4 | 9 | 13 | 5 | 18 | 1 | 0 | 1 | 38 | 57% |
| 3. E | Burns | 7 | 4 | 11 | 1 | 2 | 3 | 1 | 6 | 7 | 0 | 0 | 0 | 21 | 31% |
| 4. A | Accident | 2 | 0 | 2 | 1 | 0 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 4 | 6% |
| C | Other | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 2 | 3% |
| Т | l Total | 16 | 8 | 24 | 8 | 6 | 14 | 15 | 13 | 28 | 1 | 0 | 1 | 67 | 100% |
| 9 | % by age group | | | 36% | | | 21% | | | 42% | | | 1% | 100% | |

4. Referrals

5. Laboratory & other investigations


4.1 Referrals - general

| | 0 to 4 | 5 to 14 | 15 to 49 | 1250 | Grand total |
|----------------|--------|---------|----------|------|-------------|
| Emergency | 2 | 3 | 5 | 3 | 13 |
| Elective | 7 | 5 | 24 | 4 | 40 |
| Other | 0 | 0 | 0 | 0 | 0 |
| Total | 9 | 8 | 29 | 7 | 53 |
| % by age group | 17% | 15% | 55% | 13% | 100% |

| | Total |
|-------------------------|-------|
| Haematology | 0 |
| Imaging (X-ray, MRI/CT) | 0 |
| Other | 0 |
| Total | 2 |

6. EPI and Vitamin A

| | <1 | ≥1 to | Total |
|--|----|-------|-------------|
| BCG | 0 | 0 | 0 |
| Hepatitis B birth dose | 0 | 0 | 0 |
| DPT I/HiB I/Hep B I* (Pentavalent I) | 2 | 0 | 2 |
| DPT II/HiB II* (Tetravalent) | 5 | 0 | 2 5 3 |
| DPT III/HiB III/Hep B II* (Pentavalent II) | 3 | 0 | 3 |
| Polio 0 | 0 | 0 | 0 |
| Polio I | 2 | 0 | 2 |
| Polio II | 5 | 0 | 2 5 3 |
| Polio III | 3 | 0 | 3 |
| Measles | 2 | 0 | 2 |
| MMR | 0 | 5 | 2 5 2 |
| Rotavirus 1 | 2 | 0 | 2 |
| Rotavirus 2 | 5 | 0 | 5 |
| Rotavirus 3 | 3 | 0 | 5 3 |
| Vitamin A | 7 | 0 | 7 |
| Additional vaccines | 0 | 12 | 12 |



Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

7. Reproductive Health

7.1 Antenatal care

| | | < 18 | ≥ 18 | Total |
|---|-------------------|------|-------|-------|
| First antenatal visit < 1st trimester | | | | |
| First antenatal visit > 1st trimester | | | | |
| Number of pregnant | ≥ 4 ANC visits | N N | O DA1 | ГА |
| women at time of delivery who received: | 2 TT doses | | | |
| | anaemia screening | | | |

7.2 Postnatal care

| | < 18 | ≥ 18 | Total | |
|--|------|---------|-------|--|
| Women who attended 3 postnatal visits within 6 wks of delivery | | NO DATA | | |

7.3 Sexual and gender-based violence

Proportion of first time antenatal visits made < 1st trimester

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|---|-------|
| | М | F | M | F | rotai |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | 0 | | 0 | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 0 | 0 |
| | | | | | |

| SGBV Indicators | |
|--|-----|
| Incidence of reported rape | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

1. Health Information System Data

Publication

02 April 2014

Sources

Contact

Dr Chimmen Akawi, DOH Erbil, Email: CHIMMENTYA@yahoo.com Information









We gratefully acknowledge the contributions of the following agencies: