

1. Population

| Population | 0 to < 1 | | Total <1 | 1 to 4 | | Total 1 to 4 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total |
|--------------------|----------|-------|----------|--------|-------|--------------|---------|--------|---------------|----------|--------|----------------|-------|-------|------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | M | F | | |
| Refugee population | 2,058 | 2,058 | 4,116 | 6,273 | 7,253 | 13,526 | 18,134 | 17,644 | 35,778 | 18,624 | 23,133 | 41,757 | 1,078 | 1,764 | 2,842 | 98,019 |
| % by age group | | | 4% | | | 14% | | | 37% | | | 43% | | | 3% | 100% |

2. Mortality
2.1 Mortality

| Record ONE cause of mortality per case | 0 to 4 | | Total <5 | US % mort | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | Crude % mort |
|---|--------|---|----------|-----------|---------|---|---------------|----------|---|----------------|------|---|------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| 1. Lower Respiratory Tract Infection (LRTI) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 2. Watery diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Bloody diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4. Cardiovascular disorders | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 5. Ischaemic heart disease | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 6. Injury | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 7. Cancer | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 33% |
| 8. Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 9. Tuberculosis (confirmed) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 10. Neonatal death | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 0 | 0 | 0 | | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 1 | 2 | 2 | 67% |
| Total | 0 | 0 | 0 | 100% | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 1 | 2 | 3 | 100% |
| % by age group | | | 0% | | | | 33% | | | 0% | | | 67% | 100% | |

| Mortality indicators | Male | Female | Total |
|--------------------------------|------|--------|-------|
| Crude Mortality Rate (CMR) | 0.1 | 0.2 | 0.1 |
| Under 5 Mortality Rate (USMR) | 0.0 | 0.0 | 0.0 |
| Neonatal Mortality Rate (NNMR) | | | 0.0 |

2.2 Maternal death

| | |
|--|---|
| Number of maternal deaths reported | 0 |
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3. Morbidity
3.1 Staffing and service availability

| | |
|---|--------|
| Number of full-time clinicians | 73.0 |
| Average number of full days on which each health facility was functioning | 6.4 |
| Total number of outpatient consultations | 15,423 |
| Health facility utilisation rate | 8.2 |
| Consultations per clinician per day | 33 |

3.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total <5 | US Incid | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause | Crude Incide |
|--|--------|-----|----------|----------|---------|------|---------------|----------|------|----------------|------|-----|------------|-------------|------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | | |
| Upper Respiratory Tract Infection (URTI) | 548 | 403 | 951 | 53.9 | 377 | 397 | 774 | 471 | 526 | 997 | 94 | 88 | 182 | 2904 | 32% | 29.6 |
| Influenza-like illness (ILI) | 102 | 92 | 194 | 11.0 | 101 | 91 | 192 | 235 | 190 | 425 | 77 | 76 | 153 | 964 | 11% | 9.8 |
| Lower Respiratory Tract Infection (LRTI) | 91 | 80 | 171 | 9.7 | 58 | 33 | 91 | 53 | 52 | 105 | 42 | 53 | 95 | 462 | 5% | 4.7 |
| Watery diarrhoea* | 39 | 39 | 78 | 4.4 | 9 | 11 | 20 | 9 | 8 | 17 | 0 | 0 | 0 | 115 | 1% | 1.2 |
| Bloody diarrhoea* | 0 | 2 | 2 | 0.1 | 6 | 3 | 9 | 1 | 1 | 2 | 0 | 0 | 0 | 13 | 0% | 0.1 |
| Skin infection | 138 | 97 | 235 | 13.3 | 160 | 162 | 322 | 175 | 87 | 262 | 21 | 34 | 55 | 874 | 10% | 8.9 |
| Eye infection | 50 | 19 | 69 | 3.9 | 62 | 60 | 122 | 73 | 89 | 162 | 39 | 52 | 91 | 444 | 5% | 4.5 |
| Dental Conditions | 32 | 20 | 52 | 2.9 | 51 | 60 | 111 | 105 | 91 | 196 | 23 | 26 | 49 | 408 | 5% | 4.2 |
| Intestinal Worms | 19 | 18 | 37 | 2.1 | 20 | 27 | 47 | 2 | 8 | 10 | 1 | 3 | 4 | 98 | 1% | 1.0 |
| Acute Jaundice Syndrome* | 4 | 2 | 6 | 0.3 | 1 | 2 | 3 | 1 | 3 | 4 | 0 | 0 | 0 | 13 | 0% | 0.1 |
| Measles* | 1 | 0 | 1 | 0.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% | 0.0 |
| Meningitis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Fever of unknown origin | 18 | 10 | 28 | 1.6 | 4 | 3 | 7 | 1 | 0 | 1 | 0 | 0 | 0 | 36 | 0% | 0.4 |
| Sexually Transmitted Infection (STI) | 5 | 2 | 7 | 0.4 | 0 | 21 | 21 | 1 | 272 | 273 | 0 | 1 | 1 | 302 | 3% | 3.1 |
| Leishmaniasis | 2 | 0 | 2 | 0.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% | 0.0 |
| Other | 235 | 210 | 445 | 25.2 | 218 | 278 | 496 | 482 | 607 | 1089 | 153 | 154 | 307 | 2337 | 26% | 23.8 |
| Total | 1286 | 994 | 2280 | 129.2 | 1067 | 1148 | 2215 | 1609 | 1934 | 3543 | 450 | 487 | 937 | 6975 | 100% | 91.6 |
| % by age group | | | 25% | | | | 25% | | | 39% | | | 10% | 100% | | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|--|--------|-----|-----------|---------|-----|---------------|----------|------|----------------|------|------|------------|-------------|------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| Diabetes | 0 | 0 | 0 | 4 | 4 | 8 | 98 | 114 | 212 | 106 | 145 | 251 | 471 | 10% |
| Hypertension | 0 | 0 | 0 | 1 | 1 | 2 | 97 | 128 | 225 | 128 | 179 | 307 | 534 | 11% |
| Asthma | 14 | 22 | 36 | 43 | 27 | 70 | 45 | 61 | 106 | 18 | 15 | 33 | 245 | 5% |
| Ischaemic heart disease | 0 | 0 | 0 | 0 | 0 | 0 | 61 | 51 | 112 | 72 | 56 | 128 | 240 | 5% |
| Cardiovascular disorders | 0 | 0 | 0 | 0 | 0 | 0 | 18 | 34 | 52 | 19 | 27 | 46 | 98 | 2% |
| Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0 | 0 | 0 | 19 | 29 | 48 | 29 | 35 | 64 | 112 | 2% |
| Liver disease | 0 | 0 | 0 | 0 | 0 | 0 | 15 | 14 | 29 | 9 | 6 | 15 | 44 | 1% |
| Thyroid problems | 0 | 0 | 0 | 1 | 17 | 18 | 25 | 27 | 52 | 13 | 31 | 44 | 114 | 2% |
| Cancer | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 47 | 48 | 95 | 68 | 63 | 131 | 114 | 118 | 232 | 84 | 78 | 162 | 620 | 13% |
| Total | 122 | 140 | 262 | 234 | 224 | 458 | 984 | 1152 | 2136 | 956 | 1144 | 2100 | 4956 | 100% |
| % by age group | | | 5% | | | 9% | | | 43% | | | 42% | 100% | |

6. Laboratory

| | Total |
|--------------------------------------|-------|
| Number of laboratory tests conducted | 554 |

8. Reproductive Health

8.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|-------|
| First antenatal visit < 1st trimester | 49 | 182 | 231 |
| First antenatal visit > 1st trimester | 55 | 136 | 191 |
| Number of pregnant women at time of delivery who received: | | | |
| ≥ 4 ANC visits | 0 | 31 | 31 |
| 2 TT doses | 0 | 9 | 9 |
| anaemia screening | 0 | 42 | 42 |

| ANC Indicators | |
|--|-----|
| Proportion of first time antenatal visits made < 1st trimester | 55% |
| Coverage of complete antenatal care | 46% |
| Coverage of antenatal tetanus immunisation | 13% |
| Coverage of anaemia screening in pregnancy | 62% |

8.2 Delivery care

| | < 18 | ≥ 18 | Total |
|---|------|------|-------|
| Live births | 3 | 65 | 68 |
| Still births | 0 | 0 | 0 |
| Low birth weight (<2500 g) | 0 | 0 | 0 |
| Attended by skilled health worker | 3 | 65 | 68 |
| Number of obstetric complications treated | 0 | 11 | 11 |
| Number of caesarean sections performed | 0 | 15 | 15 |

| Delivery Indicators | |
|--|------|
| Crude birth rate | 8.3 |
| Proportion of births attended by skilled health worker (excl. TBA) | 100% |
| Proportion of low birth weight deliveries | 0% |
| Proportion of all births performed by caesarean section | 22% |
| Proportion of obstetric complications treated among girls < 18 | 0% |

8.3 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|-------|
| Women who attended 3 postnatal visits within 6 wks of delivery | 24 | 165 | 189 |

| Postnatal Indicators | |
|-------------------------------------|-----|
| Coverage of complete postnatal care | 46% |

8.4 Family planning

| | New | | Repeat | | Total |
|------------------------------|------|------|--------|------|-------|
| | < 18 | ≥ 18 | < 18 | ≥ 18 | |
| Condom (Male) | 0 | 57 | 0 | 37 | 94 |
| Implantable (Norplant) | | | | | 0 |
| Emergency Contraceptive pill | | | | | 0 |
| Sterilisation (Female) | | | | | 0 |
| Injectable (Depo - Provera) | 0 | 24 | 0 | 1 | 25 |
| Intra-Uterine Device (IUD) | 0 | 4 | 0 | | 4 |
| Oral Contraceptive | 1 | 65 | 0 | 108 | 174 |
| Family Planning Counseling | 4 | 35 | 22 | 80 | 141 |

8.5 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|---|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | 0 | | 0 | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 0 | 0 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

| SGBV Indicators | |
|--|-----|
| Incidence of reported rape | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

| | | | |
|---------------------|------------------------------|--------------------------|--------------|
| Data Sources | 1. Health Information System | Publication date: | 21 June 2013 |
|---------------------|------------------------------|--------------------------|--------------|

| | | |
|----------------------------|--|--|
| Contact Information | Za'atri Ana Calvo, UNHCR Email: calvo@unhcr.org | Amman Ann Burton, UNHCR Email: burton@unhcr.org |
|----------------------------|--|--|

UNHCR gratefully acknowledges the support of the following partners:

