



### 3.4 Mental health

	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
1. Epilepsy / seizures	0	0	0	0	0	0	2	2	4	0	0	0	4	15%
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
3. Psychotic disorder	0	0	0	0	0	0	4	4	8	0	0	0	8	31%
4. Mental retardation / intellectual disability	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
5. Severe emotional disorder	0	0	0	0	0	0	1	6	7	0	0	0	7	27%
6. Medically unexplained somatic complaint	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	1	0	1	1	1	2	2	2	4	0	0	0	7	27%
<b>Total</b>	<b>1</b>	<b>0</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>9</b>	<b>14</b>	<b>23</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>26</b>	<b>100%</b>
<b>% by age group</b>			<b>4%</b>			<b>8%</b>			<b>88%</b>			<b>0%</b>	<b>100%</b>	

### 3.5 Injuries

	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
1. Injury (intentional)	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
2. Injury (unintentional)	7	3	10	5	3	8	15	21	36	19	14	33	87	21%
3. Burns	3	2	5	2	1	3	9	16	25	14	7	21	54	13%
4. Accident	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	26	31	57	74	62	136	29	16	45	26	18	44	282	67%
<b>Total</b>	<b>36</b>	<b>36</b>	<b>72</b>	<b>81</b>	<b>66</b>	<b>147</b>	<b>53</b>	<b>53</b>	<b>106</b>	<b>59</b>	<b>39</b>	<b>98</b>	<b>423</b>	<b>100%</b>
<b>% by age group</b>			<b>17%</b>			<b>35%</b>			<b>25%</b>			<b>23%</b>	<b>100%</b>	

## 4. Referrals

	0 to 4	5 to 14	15 to 49	≥ 50	Grand total
Emergency	1	0	2	1	4
Elective	6	2	18	10	36
Other	0	0	0	0	0
<b>Total</b>	<b>7</b>	<b>2</b>	<b>20</b>	<b>11</b>	<b>40</b>
<b>% by age group</b>	<b>18%</b>	<b>5%</b>	<b>50%</b>	<b>28%</b>	<b>100%</b>

## 5. Laboratory & other investigations

	Total
Haematology	151
Imaging (X-ray, MRI/CT)	2
Other	444
<b>Total</b>	<b>715</b>

## 6. EPI and Vitamin A

### 7.1 Vaccination

Total in previous month (Domiz EPI data is reported monthly)	<1	≥1 to <5	Total
BCG	0	0	0
Hepatitis B birth dose	0	0	0
DPT I/HiB I/Hep B I* (Pentavalent I)	0	0	0
DPT II/HiB II* (Tetravalent)	0	0	0
DPT III/HiB III/Hep B II* (Pentavalent II)	0	0	0
Polio 0	0	0	0
Polio I	0	0	0
Polio II	0	0	0
Polio III	0	0	0
Measles	0	0	0
MMR	0	0	0
Rotavirus 1	0	0	0
Rotavirus 2	0	0	0
Rotavirus 3	0	0	0
Vitamin A	0	0	0
Additional vaccines	0	0	0

EPI Indicators (since December 2013)		
Total reported since last month		0
a. Program dropout rate*		24%
b. Immunisation coverage		
	BCG	95%
	Hepatitis B birth dose	93%
	DPT I	101%
	DPT III	77%
	Polio I	101%
	Polio III	77%
	Measles	51%
	MMR**	82%
	Rotavirus I	83%
	Rotavirus III	51%
c. Vitamin A coverage		94%

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

\*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines  
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%  
\*\*For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

## 7. Reproductive Health

### 7.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester	5	95	100
First antenatal visit > 1st trimester	0	0	0
Number of pregnant women at time of delivery who received:			
≥ 4 ANC visits	0	0	0
2 TT doses	0	0	0
anaemia screening	0	0	0

ANC Indicators	
Proportion of first time antenatal visits made < 1st trimester	100%

### 7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery	0	36	36

### 7.3 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No. of rape survivors given PEP within 72 hours	0	0	0	0	0
No. of female rape survivors given ECP within 120 hours	0	0	0	0	0
No. of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators		
Incidence of reported rape		0.0
Proportion of rape survivors who receive PEP < 72 hrs		
Proportion of female rape survivors who receive ECP < 120 hrs		
Proportion of rape survivors who receive STI presumptive treatment < 2 wks		

Data Sources: 1. Health Information System

Publication date: 10 April 2014

Contact Information: Dohuk  
Dr Bakhtiyar AHMED, DOH Duhok

Email: BAHMED@duhokhealth.org

Contact Information: Erbil  
Sandra HARLASS, UNHCR

Email: HARLASS@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

