



### 3.4 Mental health

	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
1. Epilepsy / seizures	0	0	0	3	3	6	11	5	16	0	0	0	22	12%
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	4	0	4	0	0	0	4	2%
3. Psychotic disorder	0	0	0	0	0	0	11	12	23	4	1	5	28	15%
4. Mental retardation / intellectual disability	2	0	2	6	0	6	1	2	3	0	0	0	11	6%
5. Severe emotional disorder	5	4	9	14	11	25	14	30	44	0	3	3	81	44%
6. Medically unexplained somatic complaint	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	1	1	2	11	6	17	7	8	15	0	3	3	37	20%
<b>Total</b>	<b>8</b>	<b>5</b>	<b>13</b>	<b>34</b>	<b>20</b>	<b>54</b>	<b>48</b>	<b>57</b>	<b>105</b>	<b>4</b>	<b>7</b>	<b>11</b>	<b>183</b>	<b>100%</b>
% by age group			7%			30%			57%			6%	100%	

### 3.5 Injuries

	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
1. Injury (intentional)	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
2. Injury (unintentional)	23	14	37	28	19	47	86	106	192	100	111	211	487	21%
3. Burns	13	13	26	20	17	37	64	76	140	69	56	125	328	14%
4. Accident	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	83	132	215	418	458	876	105	99	204	111	105	216	1511	65%
<b>Total</b>	<b>119</b>	<b>159</b>	<b>278</b>	<b>466</b>	<b>494</b>	<b>960</b>	<b>255</b>	<b>281</b>	<b>536</b>	<b>280</b>	<b>272</b>	<b>552</b>	<b>2326</b>	<b>100%</b>
% by age group			12%			41%			23%			24%	100%	

## 4. Referrals

	0 to 4	5 to 14	15 to 49	≥ 50	Grand total
Emergency	12	14	15	9	50
Elective	31	29	91	32	183
Other	0	0	0	1	1
<b>Total</b>	<b>43</b>	<b>43</b>	<b>106</b>	<b>42</b>	<b>234</b>
% by age group	18%	18%	45%	18%	100%

## 5. Laboratory & other investigations

	Total
Haematology	772
Imaging (X-ray, MRI/CT)	7
Other	2503
<b>Total</b>	<b>3850</b>

## 6. EPI and Vitamin A

### 7.1 Vaccination

	<1	≥1 to <5	Total
BCG	0	0	0
Hepatitis B birth dose	0	0	0
DPT I/HiB I/Hep B I* (Pentavalent I)	0	0	0
DPT II/HiB II* (Tetravalent)	0	0	0
DPT III/HiB III/Hep B II* (Pentavalent II)	0	0	0
Polio 0	0	0	0
Polio I	0	0	0
Polio II	0	0	0
Polio III	0	0	0
Measles	0	0	0
MMR	0	0	0
Rotavirus 1	0	0	0
Rotavirus 2	0	0	0
Rotavirus 3	0	0	0
Vitamin A	0	0	0
Additional vaccines	0	0	0

EPI Indicators (since December 2013)		Total
Total reported		0
a. Program dropout rate*		24%
b. Immunisation coverage		
BCG		95%
Hepatitis B birth dose		93%
DPT I		101%
DPT III		77%
Polio I		101%
Polio III		77%
Measles		51%
MMR**		82%
Rotavirus I		83%
Rotavirus III		51%
c. Vitamin A coverage		94%

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

\*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines  
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%  
\*\*For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

## 7. Reproductive Health

### 7.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester	48	713	761
First antenatal visit > 1st trimester	0	0	0
Number of pregnant women at time of delivery who received:			
≥ 4 ANC visits	0	0	0
2 TT doses	0	0	0
anaemia screening	0	0	0

#### ANC Indicators

Proportion of first time antenatal visits made < 1st trimester	100%
--	------

### 7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery	6	188	194

### 7.3 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No of rape survivors given PEP within 72 hours	0	0	0	0	0
No of female rape survivors given ECP within 120 hours	0	0	0	0	0
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

#### SGBV Indicators

Incidence of reported rape	0.0
Proportion of rape survivors who receive PEP < 72 hrs	
Proportion of female rape survivors who receive ECP < 120 hrs	
Proportion of rape survivors who receive STI presumptive treatment < 2 wks	

Data Sources

1. Health Information System

Publication date:

30 April 2014

Contact Information

Dohuk  
Dr Bakhtiyar AHMED, DOH Duhok

Email: BAHMED@duhokhealth.org

Erbil  
Sandra HARLASS, UNHCR

Email: HARLASS@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

