


Refugee population

Total refugee population: **7,504** *

Demographic breakdown:



Key facts

Camp opened: **September 2013**

Planned capacity: **20,000**

Area: **1.15 Km sq**

Population: **Syrian refugees, majority from Aleppo & Qamishli**

* Source of population data: Estimates from UNHCR registration. For multi week reports, average population is presented

Indicators at a glance

Indicator	Indicator	Standard
1 Health utilisation rate	7.6	1 - 4
2 Number of consultations per clinician per day	59	< 50
3 Proportion of first time antenatal visits made < 1st trimester	N/A	≥ 90%
4 Measles immunisation coverage	170%	> 90%
5 Polio III immunisation coverage	89%	> 90%
6 Crude Mortality Rate (CMR) (/1000/month)	0.0	< 1.5
7 Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 3.0
8 Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0.0	< 40

Notes

HIGHLIGHTS

Health utilisation rate (HUR) and consultations per clinicians (CPC) elevated. A thorough investigation of possible explanations for elevated HUR and CPC needed.


This public health profile for Darashakran refugee camp is compiled from health data obtained from reporting agencies. Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Agencies reporting:


**DOH - Erbil
MSF**

Morbidity

Acute health conditions (%)



Chronic health conditions (%)



Top 5 acute health conditions


	% morbidity	# of cases
1 Upper Respiratory Tract Infection (URTI)	32%	1,457
2 Lower Respiratory Tract Infection (LRTI)	18%	816
3 Skin infection	5%	230
4 Dental Conditions	4%	160
5 Urinary tract infection (UTI)	3%	158
- Other	38%	1,750

Top 5 chronic health conditions


	% morbidity	# of cases
1 Hypertension	42%	307
2 Diabetes	10%	69
3 Asthma	3%	25
4 Thyroid problems	3%	24
5 Ischaemic heart disease	3%	20
- Other	39%	279

Breakdown of acute health conditions

by age




by gender




Breakdown of chronic health conditions


by age




by gender




Number of consultations




Weekly health utilisation rates




Watery diarrhoea




Bloody diarrhoea




Acute Jaundice Syndrome




Suspected measles




Suspected Meningitis




Acute flaccid paralysis



Upper respiratory tract infections



Lower respiratory tract infections



Calculation of disease alert thresholds:

Watery diarrhoea = 1.5 x baseline, Bloody diarrhoea = 1.5 x baseline, Acute Jaundice Syndrome = 1.5 x baseline, Upper respiratory tract infections = 1.5 x baseline,
 Meningitis = 5 cases, Acute Flaccid Paralysis = 1 case, Measles = 1 case,
 baseline = average incidence or number of cases over the previous 3 weeks

Public Health

Health Impact

Indicator	Value	Standard
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 1.5
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 3.0
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0.0	< 40

Outbreak Alert and Response

Indicator	Value	Standard
Number of outbreaks reported	0	
% of outbreaks investigated within 48 hours		100%

Routine immunization

Indicator	Value	Standard
BCG immunisation coverage	56%	> 90%
DPT 1 immunisation coverage	64%	> 90%
Measles immunisation coverage	170%	> 90%
Polio III immunisation coverage	89%	> 90%
MMR immunisation coverage	112%	> 90%
Vitamin A coverage	124%	> 90%

Access and utilization

Indicator	Value	Standard
No. of health facilities	1.0	1 : < 10,000
Total number of consultations	5,456	
Health utilisation rate	7.6	1 - 4
No. of consultations per trained clinician per day	59	< 50

Sexual and gender-based violence

Indicator	Value
Incidence of reported rape	0.0

Reproductive health

Indicator	Value	Standard
Proportion of first time antenatal visits made < 1st trimester	N/A	≥ 90%

Top 3 Mental health conditions


Rank	Condition	% morbidity	# of cases
1	Epilepsy / seizures	13%	12
2	Mental retardation / intellectual disability	1%	1
3	Severe emotional disorder	1%	1
-	Other	85%	78

Top 3 injuries


Rank	Injury	% morbidity	# of cases
1	Injury (unintentional)	33%	23
2	Burns	22%	15
3	Accident	6%	4
-	Other	39%	27

Breakdown of mental health conditions


By cause



By age




By gender




Breakdown of injuries


By cause



By age



By gender



We gratefully acknowledge the support of the following agencies:



View interactive maps and statistics online
<http://twine.unhcr.org>

