

1. Morbidity

1.1 Staffing and service availability

| | |
|---|---------|
| Number of full-time clinicians | 15.6 |
| Average number of full days on which each health facility was functioning | 5.5 |
| Total number of outpatient consultations | 104,245 |
| Consultations per clinician per day | 23 |

1.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total <5 | U5 Incid | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|--|--------------|--------------|--------------|----------|--------------|--------------|---------------|---------------|---------------|----------------|--------------|--------------|--------------|---------------|-------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| Upper Respiratory Tract Infection (URTI) | 2,553 | 2,259 | 4,812 | | 3,257 | 3,111 | 6,368 | 2,380 | 4,636 | 7,016 | 448 | 437 | 885 | 19,081 | 32% |
| Influenza-like illness (ILI) | 282 | 291 | 573 | | 480 | 509 | 989 | 462 | 902 | 1,364 | 70 | 81 | 151 | 3,077 | 5% |
| Lower Respiratory Tract Infection (LRTI) | 143 | 119 | 262 | | 187 | 128 | 315 | 353 | 427 | 780 | 103 | 75 | 178 | 1,535 | 3% |
| Watery diarrhoea* | 74 | 73 | 147 | | 55 | 37 | 92 | 48 | 49 | 97 | 11 | 8 | 19 | 355 | 1% |
| Bloody diarrhoea* | 0 | 0 | 0 | | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 | 2 | 0% |
| Skin infection | 301 | 315 | 616 | | 487 | 536 | 1,023 | 710 | 1,139 | 1,849 | 101 | 115 | 216 | 3,704 | 6% |
| Eye infection | 66 | 45 | 111 | | 104 | 87 | 191 | 160 | 200 | 360 | 41 | 37 | 78 | 740 | 1% |
| Ear infection | 120 | 92 | 212 | | 108 | 129 | 237 | 213 | 460 | 673 | 45 | 48 | 93 | 1,215 | 2% |
| Dental Conditions | 126 | 105 | 231 | | 1,435 | 1,696 | 3,131 | 2,768 | 5,098 | 7,866 | 314 | 266 | 580 | 11,808 | 20% |
| Intestinal Worms | 49 | 60 | 109 | | 161 | 144 | 305 | 42 | 89 | 131 | 4 | 5 | 9 | 554 | 1% |
| Acute Jaundice Syndrome* | 0 | 2 | 2 | | 0 | 2 | 2 | 1 | 1 | 2 | 0 | 0 | 0 | 6 | 0% |
| Measles (suspected)* | 1 | 1 | 2 | | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0% |
| Meningitis (suspected)* | 2 | 0 | 2 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Leishmaniasis | 1 | 3 | 4 | | 1 | 2 | 3 | 2 | 8 | 10 | 0 | 2 | 2 | 19 | 0% |
| Fever of unknown origin | 2 | 1 | 3 | | 2 | 0 | 2 | 1 | 1 | 2 | 0 | 0 | 0 | 7 | 0% |
| Reproductive tract infection | 0 | 0 | 0 | | 0 | 6 | 6 | 2 | 163 | 165 | 1 | 3 | 4 | 175 | 0% |
| Tuberculosis (suspected) | 0 | 0 | 0 | | 1 | 0 | 1 | 3 | 1 | 4 | 1 | 0 | 1 | 6 | 0% |
| Other | 623 | 663 | 1,286 | | 970 | 1,376 | 2,346 | 3,216 | 7,965 | 11,181 | 802 | 827 | 1,629 | 16,442 | 28% |
| Total | 4,343 | 4,029 | 8,372 | | 7,250 | 7,763 | 15,013 | 10,361 | 21,139 | 31,500 | 1,942 | 1,904 | 3,846 | 58,731 | 100% |
| % by age group | | | 14% | | | | 26% | | | 54% | | | 7% | 100% | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

1.3 Chronic health conditions

| Record NEW and REVISITS | | | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|--|--|--|--------|-----|-----------|---------|-----|---------------|----------|--------|----------------|-------|-------|------------|-------------|------------|
| | | | M | F | | M | F | | M | F | | M | F | | | |
| Diabetes | | | 4 | 1 | 5 | 65 | 67 | 132 | 1,938 | 2,399 | 4,337 | 1,856 | 1,853 | 3,709 | 8,183 | 21% |
| Hypertension | | | 0 | 0 | 0 | 7 | 11 | 18 | 2,406 | 3,012 | 5,418 | 2,642 | 2,453 | 5,095 | 10,531 | 27% |
| Asthma | | | 27 | 14 | 41 | 93 | 48 | 141 | 267 | 399 | 666 | 217 | 152 | 369 | 1,217 | 3% |
| Ischaemic heart disease | | | 0 | 0 | 0 | 4 | 10 | 14 | 1,009 | 426 | 1,435 | 960 | 557 | 1,517 | 2,966 | 8% |
| Cardiovascular disease (other) | | | 14 | 10 | 24 | 20 | 13 | 33 | 472 | 409 | 881 | 610 | 381 | 991 | 1,929 | 5% |
| Chronic obstructive pulmonary disease (COPD) | | | 0 | 0 | 0 | 0 | 0 | 0 | 11 | 12 | 23 | 49 | 20 | 69 | 92 | 0% |
| Liver disease | | | 1 | 0 | 1 | 5 | 1 | 6 | 34 | 38 | 72 | 4 | 5 | 9 | 88 | 0% |
| Thyroid problems | | | 5 | 8 | 13 | 17 | 49 | 66 | 113 | 836 | 949 | 71 | 148 | 219 | 1,247 | 3% |
| Cancer | | | 19 | 4 | 23 | 20 | 10 | 30 | 148 | 255 | 403 | 94 | 68 | 162 | 618 | 2% |
| Other | | | 267 | 169 | 436 | 603 | 615 | 1,218 | 3,166 | 4,346 | 7,512 | 1,742 | 1,150 | 2,892 | 12,058 | 31% |
| Total | | | 337 | 206 | 543 | 834 | 824 | 1,658 | 9,564 | 12,132 | 21,696 | 8,245 | 6,787 | 15,032 | 38,929 | 100% |
| % by age group | | | | | 1% | | | 4% | | | 56% | | | 39% | 100% | |

1.4 Mental health

| | | | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|----|---|--|--------|----|-----------|---------|-----|---------------|----------|-------|----------------|------|-----|------------|-------------|------------|
| | | | M | F | | M | F | | M | F | | M | F | | | |
| 1. | Epilepsy / seizures | | 30 | 11 | 41 | 290 | 217 | 507 | 465 | 382 | 847 | 31 | 10 | 41 | 1,436 | 21% |
| 2. | Alcohol and other substance use disorder | | 0 | 0 | 0 | 2 | 0 | 2 | 5 | 1 | 6 | 0 | 0 | 0 | 8 | 0% |
| 3. | Psychotic disorder (including mania) | | 0 | 0 | 0 | 57 | 51 | 108 | 1,251 | 916 | 2,167 | 46 | 25 | 71 | 2,346 | 34% |
| 4. | Intellectual disability | | 13 | 8 | 21 | 47 | 23 | 70 | 56 | 24 | 80 | 0 | 0 | 0 | 171 | 2% |
| 5. | Severe emotional disorder, including moderate-severe depression | | 1 | 2 | 3 | 31 | 28 | 59 | 884 | 896 | 1,780 | 65 | 80 | 145 | 1,987 | 29% |
| 6. | Medically unexplained somatic complaint | | 0 | 0 | 0 | 8 | 5 | 13 | 16 | 17 | 33 | 0 | 0 | 0 | 46 | 1% |
| | Other psychological complaints | | 5 | 5 | 10 | 89 | 65 | 154 | 330 | 350 | 680 | 19 | 24 | 43 | 887 | 13% |
| | Total | | 49 | 26 | 75 | 524 | 389 | 913 | 3,007 | 2,586 | 5,593 | 161 | 139 | 300 | 6,881 | 100% |
| | % by age group | | | | 1% | | | 13% | | | 81% | | | 4% | 100% | |

1.5 Injuries

| | | | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|---|--|--|--------|----|-----------|---------|----|---------------|----------|-----|----------------|------|----|------------|-------------|------------|
| | | | M | F | | M | F | | M | F | | M | F | | | |
| 1. Bites (all causes) | | | 2 | 1 | 3 | 5 | 0 | 5 | 6 | 6 | 12 | 0 | 0 | 0 | 20 | 3% |
| 2. Injury (war related) | | | 2 | 2 | 4 | 4 | 5 | 9 | 22 | 6 | 28 | 1 | 1 | 2 | 43 | 7% |
| 3. Injury (Burns) | | | 18 | 16 | 34 | 13 | 15 | 28 | 16 | 14 | 30 | 1 | 2 | 3 | 95 | 15% |
| 4. Assault (with weapon non-war related) | | | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 3 | 0 | 0 | 0 | 3 | 0% |
| Other | | | 34 | 26 | 60 | 113 | 49 | 162 | 162 | 80 | 242 | 13 | 7 | 20 | 484 | 75% |
| Total | | | 56 | 45 | 101 | 135 | 69 | 204 | 208 | 107 | 315 | 15 | 10 | 25 | 645 | 100% |
| % by age group | | | | | 16% | | | 32% | | | 49% | | | 4% | 100% | |

1.6 Reproductive Tract Infections

| | < 18 | | Total < 18 | ≥ 18 | | Total ≥ 18 | Grand total | % by cause |
|-------------------------|----------|-----------|------------|-----------|------------|------------|-------------|-------------|
| | M | F | | M | F | | | |
| 1. Vaginal discharge | 0 | 10 | 10 | 0 | 776 | 776 | 786 | 77% |
| 2. Urethral discharge | 1 | 0 | 1 | 9 | 0 | 9 | 10 | 1% |
| 3. Genital ulcer | 0 | 1 | 1 | 1 | 5 | 6 | 7 | 1% |
| 3. Lower abdominal pain | 0 | 8 | 8 | 0 | 153 | 153 | 161 | 16% |
| Other | 0 | 8 | 8 | 3 | 44 | 47 | 55 | 5% |
| Total | 1 | 27 | 28 | 13 | 978 | 991 | 1019 | 100% |
| % by age group | | | 3% | | | 97% | 100% | |

2. Referrals

2.1 Referrals by location

| | | 0 to 4 | Total < 5 | 5 to 17 | Total 5 to 17 | 18 to 59 | Total 18 to 59 | > 60 | Total | Grand total |
|----------------------------|-----------------------------|--------|-----------|---------|---------------|----------|----------------|-------|-------|-------------|
| 1. | MOH Hospital | 1,647 | 1,647 | 2,301 | 2,301 | 6,403 | 6,403 | 1,175 | 1,175 | 11,526 |
| 2. | Royal Medical Services | 86 | 86 | 60 | 60 | 223 | 223 | 24 | 24 | 393 |
| 3. | Private affiliated Hospital | 205 | 205 | 151 | 151 | 1,717 | 1,717 | 223 | 223 | 2,296 |
| 4. | Jordan University Hospital | 157 | 157 | 180 | 180 | 573 | 573 | 219 | 219 | 1,129 |
| 5. | Other | 9 | 9 | 3 | 3 | 5 | 5 | 1 | 1 | 18 |
| Total No of referred cases | | 15,362 | | | | | | | | |

2.2 Referrals by service

| | | 0 to 4 | Total < 5 | 5 to 17 | Total 5 to 17 | 18 to 59 | Total 18 to 59 | > 60 | Total | Grand total |
|----|-------------------|--------|-----------|---------|---------------|----------|----------------|------|-------|-------------|
| 1. | Paediatrics | 1,489 | 1,489 | 529 | 529 | 1 | 1 | 0 | 0 | 2,019 |
| 2. | Internal medicine | 279 | 279 | 640 | 640 | 2,488 | 2,488 | 790 | 790 | 4,197 |
| 3. | Obstetric | 3 | 3 | 529 | 529 | 2,503 | 2,503 | 245 | 245 | 3,280 |
| 4. | Gynaecological | 0 | 0 | 41 | 41 | 1,649 | 1,649 | 23 | 23 | 1,713 |
| 5. | Surgical | 67 | 67 | 151 | 151 | 578 | 578 | 101 | 101 | 897 |
| 6. | Other | 266 | 266 | 805 | 805 | 1,702 | 1,702 | 483 | 483 | 3,256 |

3. Reproductive Health

3.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|---------------------------------------|------|------|-------|
| First antenatal visit < 1st trimester | 8 | 321 | 329 |
| First antenatal visit > 1st trimester | 9 | 728 | 737 |

ANC Indicators

| | |
|--|-----|
| Proportion of first time antenatal visits made < 1st trimester | 31% |
|--|-----|

3.2 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|-------|
| At least three PNC visits within 6 weeks | 0 | 27 | 27 |

3.3 Family planning

| | New | | Repeat | | Total |
|------------------------------|------|-------|--------|------|-------|
| | < 18 | ≥ 18 | < 18 | ≥ 18 | |
| Oral contraceptives | 0 | 123 | 3 | 74 | 200 |
| Emergency Contraceptive Pill | 0 | 0 | 0 | 0 | 0 |
| Injectable (Depo-Provera) | 0 | 24 | 0 | 5 | 29 |
| Implantable (Norplant) | 0 | 0 | 0 | 0 | 0 |
| Intra-Uterine Device (IUD) | 0 | 21 | 0 | 12 | 33 |
| Condom (Male) | 0 | 82 | 1 | 48 | 131 |
| Other | 0 | 0 | 0 | 0 | 0 |
| | | | | | |
| | < 18 | ≥ 18 | | | |
| Family Planning Counseling* | 3 | 1,269 | | | |

3.4 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|---|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | 0 | | 0 | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 5 | 5 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

Note

This report for non-camp Syrian refugees in Jordan is compiled from health data obtained from reporting agencies and clinics. Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Clinics and agencies reporting:

JHAS - Amman governorate (Madina clinic); Irbid governorate (Cyber City clinic, Irbid clinic, Ramtha clinic); Mafraq governorate (Mafraq clinic); Zarqa governorate (Zarqa clinic); South governorates (Mobile medical unit)

Reproductive tract infections were collected differently in the first quarter of 2015 thus on the report it appears twice under Acute health conditions and as a separate component under Morbidity

Published on

10/03/2016

Contact
Information

UNHCR
Ann Burton

JHAS
Samah Al Ouran