Azraq | Jordan

Refugee population

Total refugee population: **36,860**

Demographic breakdown:

Age	Male	Female
> 60	1%	1%
18 - 59	19%	22%
5 - 17	19%	18%
0 - 4	10%	10%

Male % **Female**

Key facts

Camp opened:
May 2014

Indicators at a glance

|			
1	Health utilisation rate	7.7	X
2	Number of consultations per clinician per day	44	√
3	Crude Mortality Rate (CMR) (/1000/month)	0.2	√
4	Under-five Mortality Rate (U5MR) (/1000/month)	0.5	√
5	Neonatal Mortality Rate (NNMR) (/1000 livebirths)	16.4	X

Notes

This public health profile for Azraq refugee camp is compiled from health facility data from the following partners:
- **IMC Clinics in Villages 3,5, and 6**
- **IMC Hospital**
- **Arab Medical Relief Clinics in Villages 2 and 6**
- **IRC RH Services Villages 3,5, and 6**

Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Morbidity

Acute health conditions (%)

- Upper Respiratory Tract Infection (URTI) — 24%
- Dental Conditions — 10%
- Influenza-like illness (ILI) — 7%
- Skin infection — 4%
- Watery diarrhoea* — 4%
- Other — 51%

Chronic health conditions (%)

- Hypertension — 34%
- Diabetes — 19%
- Asthma — 17%
- Thyroid problems — 5%
- Ischaemic heart disease — 4%
- Other — 21%

Top 5 acute health conditions

		% morbidity	# of cases
1	Upper Respiratory Tract Infection (URTI)	24%	11,091
2	Dental Conditions	10%	4,536
3	Influenza-like illness (ILI)	7%	3,365
4	Skin infection	4%	1,926
5	Watery diarrhoea*	4%	1,842
-	Other	50%	23,211

Top 5 chronic health conditions

		% morbidity	# of cases
1	Hypertension	34%	1,380
2	Diabetes	19%	794
3	Asthma	17%	687
4	Thyroid problems	5%	199
5	Ischaemic heart disease	5%	186
-	Other	21%	871

Breakdown of acute health conditions

by age

- 0-4 — 33%
- 5 to 17 — 24%
- 18 to 59 — 40%
- > 60 — 3%

by gender

- Male — 46%
- Female — 54%

Breakdown of chronic health conditions

by age

- 0-4 — 12%
- 5 to 17 — 7%
- 18 to 59 — 52%
- > 60 — 29%

by gender

- Male — 43%
- Female — 57%

UNHCR gratefully acknowledges the support of the following partners:

Public Health

Health Impact

|	Indicator	Standard	
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.5	√
Under-five Mortality Rate (U5MR) (/1000/month)	0.5	< 1.5	√
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	16.4	< 14.9	X

Access and utilization

	Indicator	Standard	
No. of health facilities	1 : 7372	1 : < 10,000	√
Total number of consultations	71,231		
Health utilisation rate	7.7	1 - 4	X
No. of consultations per trained clinician per day	44	< 50	√

Breakdown of mental health conditions

By cause

- Severe emotional disorder, including moderate-severe depression
- Epilepsy / seizures
- Psychotic disorder (including mania)
- Intellecutal disability
- Other

37%, 23%, 7%, 5%, 28%

By age

- 0-4
- 5 to 17
- 18 to 59
- > 60

3% 3%, 31%, 63%

By gender

- Male
- Female

44%, 56%

Breakdown of injuries

By cause

- Injury (burns)
- Bites (all causes)
- Injury (war-related)
- Injury (war-related)
- Other

12%, 2%, 86%

By age

- 0-4
- 5 to 17
- 18 to 59
- > 60

4%, 33%, 34%, 29%

By gender

- Male
- Female

37%, 63%

In-patient and referral

In-patient department

	Indicator	Standard	
Hospitalisation rate (/1000/year)	96.5	50 - 150	^
Bed occupancy rate	77%	> 75%	√
Mean length of stay for authorised discharges (days)	2.0		

	Indicator	Standard
Overall case fatality rate	0%	
Proportion of deaths within 24 hours of admission	0%	

Reproductive Health

Antenatal care

	Indicator	Standard	
Coverage of antenatal care (4 or more visits)	67%	≥ 90%	X
Coverage of TT vaccination	30%	≥ 90%	X
Coverage of anaemia screening in pregnancy	98%	≥ 90%	√

Delivery care

	Indicator	Standard	
Proportion of deliveries attended by skilled personnel	95%	≥ 90%	√
Proportion of deliveries performed by caesarean section	16.4%	5 - 15%	X
Proportion of low birth weight deliveries (<2500g)	8%	< 15%	√

Maternal mortality

	Indicator	Standard
Number of maternal deaths reported	0	
% of maternal deaths investigated	0%	100%

Postnatal care

	Indicator	Standard	
Coverage of postnatal care	121%	≥ 90%	√

Sexual and Gender-based Violence

	Indicator	Standard		
Incidence of rape (reported cases/10,000/year)*	1.1		^	⚑
% of rape survivors who received PEP** < 72 hours		100%	^	⚑
% of rape survivors who received ECP*** < 120 hours		100%	^	⚑
% of rape survivors who received STI**** prophylaxis < 2 wks		100%	^	⚑

*Only includes rapes reported at health facilities **Post-exposure prophylaxis ***Emergency contraceptive pill ****Sexually transmitted infection

Legend:
- ✓ Standard met
- ⚠ Standard borderline
- ✗ Standard not met
- ⚑ Interpret data with care
- ? Data missing