




Refugee population^a

25,616

 at end of period

Figure 1 Population breakdown by age-group



U5: Under 5 WCBA: Women of child-bearing age



Photo: Kobe refugee camp © UNHCR / Aug 2011

Key dates:

Camp opened: 24 June 2011

Last nutrition survey: None conducted



Indicators at a glance:

1. Crude Mortality Rate ^c	1.1	⚠️
2. Under 5 Mortality Rate ^c	3.2	❌
3. Infant Mortality Rate	n/a	ℹ️
4. Severe Acute Malnutrition (SAM) rate ^d	16%	❌
5. Global Acute Malnutrition (GAM) rate ^d	37%	❌
6. Measles coverage ^e	85%	⚠️
7. Skilled attendance at delivery	100%	✅
8. Water (litres / refugee / day) ^f	16	✅

Table 1: Top causes of mortality

Not available

Table 2: Top causes of morbidity^b

1. URTI	31%
2. Intestinal worms	16%
3. LRTI	13%
4. Watery diarrhoea	11%
5. Skin disease	5%

LRTI: Lower respiratory tract infection

URT: Upper respiratory tract infection

Sources of data

- ^a Source: UNHCR registration database
- ^b Source: MSF weekly reports
- ^c Source: Grave counting
- ^d Source: Household MUAC screening (September 2011)
- ^e Source: MSF mass measles campaign and MUAC screening (August 2011)
- ^f Source: UNHCR WASH Monitoring Reports

Reporting period

- All indicators are for week 38, with the exception of GAM and SAM rates.
- GAM and SAM rates are based on a house-to-house MUAC screening conducted between 4 and 7 September 2011.
- WASH indicators are based on monitoring reports from 4th September 2011

Summary:

- There has been a steady decline in CMR in Kobe over the last few weeks. Efforts are ongoing to reduce mortality.
- The cause-specific deaths are taken from what is reported in HIS. There was limited cause-specific data on mortality this week
- Deliveries attended by skilled health worker have improved.
- UNICEF has provided 1 mobile health team to support basic treatment, community outreach, active case finding and referral. They are prioritizing zones on the camp periphery.
- IMC has 2 satellite MAM/BF sites operational.

Public Health Priorities:

- Immediate priority remains reduction of excess mortality in Kobe.
- Move from grave counting to reporting of deaths by community and other health workers. Closely monitor and strengthen community-based mortality surveillance.
- Strengthen active case finding, defaulter tracing and referral as a key factor in reduction of CMR.
- Coordination of community health programme including hygiene promotion.
- Hasten construction of additional latrines and laying down of water pipe from the water treatment plant.
- Reporting on ongoing community health activities
- HIS training for all health and nutrition partners planned for mid-October 2011 and Nutrition survey planned for mid-October 2011

Legend: ✅ Standard reached ⚠️ Standard borderline ❌ Standard not reached ? Data unreliable ℹ️ Indicator cannot be calculated n/a Data not available - Data not applicable

View interactive maps and statistics online: <http://his.unhcr.org/main.locsis>

Public Health						
Health Impact	No	Indicator	Emergency Standard	Outbreak Alert and Response	Indicator	Emergency Standard
Crude Mortality Rate (/10,000/day)		1.1	< 1	Number of outbreaks reported	0	
Under-five Mortality Rate (/10,000/day)		3.2	< 2	% of outbreaks investigated < 48 hours	-	100%
Infant Mortality Rate (IMR) (/1000 livebirths)		n/a	< 60	EPI and Vitamin A		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)		n/a	< 40	Measles vaccination coverage	85%	> 95%
Global Acute Malnutrition Rate (%)		37%	< 10%	Supervision		
Severe Acute Malnutrition Rate (%)		16%	< 2%	Do regular camp coordination meetings take place?	Yes	Yes
Access and Utilisation				Were any drug shortages reported during the period?	No	No
No. of health facilities	1	1 : 25,616	1 : <10,000			
No. of consultations per trained clinician per day		62	< 50			
Health Utilization Rate (new visits/person/year)		1.5	1 - 4			

Nutrition	Moderate Acute Malnutrition (MAM)*		Severe Acute Malnutrition (SAM)						
	Emergency Standard		Outpatient therapeutic program (OTP)	Emergency Standard	Stabilisation Centre (SC)**	Emergency Standard	CMAM** (SC and OTP combined)	Emergency Standard	
Number of new admissions		50	160		n/a		160		
Average length of stay		n/a	< 8 weeks	n/a	< 30 days	n/a	< 10 days	1.9	< 30 days
Average weight gain (g/kg/day)		-		n/a	> 5	n/a		0.1	> 5
Discharge rate		n/a	> 75%	64%	> 75%	n/a		67%	> 75%
Death rate		n/a	< 3%	0%	< 10%	n/a		0%	< 10%
Default rate		n/a	< 15%	32%	< 15%	n/a		33%	< 15%
Referral rate		n/a				n/a		0%	
Non-cured rate		-		0%		-		0%	

* no exits were recorded from MAM during the reporting period

** there is currently no SC in Kobe camp. SC and overall CMAM indicators represent data for Kobe and Melkadida combined.

Reproductive Health and HIV							
Maternal and Newborn Health	No	Indicator	Emergency Standard	Sexual and Gender-based Violence	No	Indicator	Emergency Standard
No. of basic EmOC facilities	1	1 : 25,616	1 : <500,000	Incidence of reported rape (/10,000/year)	0	0.0	
No. of comprehensive EmOC facilities	0	0	1 : <500,000	% rape survivors who received PEP < 72h	-	100%	
Number of maternal deaths		0		% rape survivors who received ECP < 120h	-	100%	
Number of maternal deaths investigated <48 hrs		-	100%	% rape survivors who received STI < 2 wks	-	100%	
Crude Birth Rate (CBR) (/1000/month)		0.5		HIV/AIDS			
Coverage complete antenatal care (> 4 visits)		n/a	> 90%	Condom distribution rate	n/a	> 0.5	
% deliveries performed by caesarean section		0%	5 - 15%	% of blood units screened for HIV	n/a	100%	
% deliveries attended by skilled personnel		100%	≥ 50%				
% low birth weight deliveries		0%	< 15%				

PEP: Post-exposure prophylaxis; ECP: Emergency Contraceptive Pill; STI: Sexually Transmitted Infection

Community Health Activities			Water, Sanitation and Hygiene		
	Indicator	Emergency Standard		Indicator	Emergency Standard
No. of health education and hygiene sessions conducted	-		Avg quantity of potable water / person / day (litres)	16	> 10
No. of sanitation campaigns conducted	-		No. of persons per usable water tap	170	< 250
No. pregnant women who received clean delivery kit	-		No. of persons per drop-hole in communal latrine	57	≤ 50
No. of complicated medical cases identified	-		% of population living within 200m from water point	146%	100%
% of complicated medical cases referred	-	≥ 90%	% of families with latrines	54%	> 50%

Legend: Standard reached Standard borderline Standard not reached Data unreliable Indicator cannot be calculated n/a Data not available - Data not applicable

View interactive maps and statistics online: <http://his.unhcr.org/main.locsis>

UNHCR gratefully acknowledges the support of the following partners:

