



Refugee population^a


20,774

 at end of period

Figure 1 Population breakdown by age-group



Photo: Hilaweyn refugee camp © UNHCR / Aug 2011



U5: Under 5 WCBA: Women of child-bearing age

Table 1: Top causes of mortality^b

1. Acute malnutrition	38%
2. LRTI	38%
3. Meningitis	25%

Table 2: Top causes of morbidity^b

1. Watery diarrhoea	24%
2. LRTI	17%
3. Other	13%
4. URTI	11%
5. Eye disease	7%

LRTI: Lower respiratory tract infection

Sources of data

^a Source: UNHCR registration database

^b Source: MSF-H weekly reports

^c Source: Grave count (3 – 9 September)

^d Source: ACF MUAC and W/H Screening (Aug/Sep 2011)

^e Source: MSF. All children are vaccinated prior to relocation.

^f Source: UNHCR WASH Monitoring Reports

URT: Upper respiratory tract infection

Reporting period

• All indicators are for week 39, with the exception of GAM and SAM rates.

• GAM and SAM rates are based on ACF MUAC and W/H Screening of all children relocated from transit to Hilaweyn between August 5th and September 9th. The next nutrition survey will take place in October 2011.

• WASH indicators are based on monitoring reports from 3rd October 2011.



Indicators at a glance:

1. Crude Mortality Rate ^c	1.2	✗
2. Under 5 Mortality Rate ^c	3.1	✗
3. Infant Mortality Rate	n/a	i
4. Severe Acute Malnutrition (SAM) rate ^d	26%	✗
5. Global Acute Malnutrition (GAM) rate ^d	44%	✗
6. Measles coverage ^e	>95%	✓
7. Skilled attendance at delivery	100%	✓
8. Water (litres / refugee / day) ^f	16	✓

Summary:

- The nutrition screening in Hilaweyn represent children who are not yet included in the Transit Camp nutrition programmes, so the rates do not show the real status of the new refugees coming to Hilaweyn, However these children are thereafter included in the nutrition programmes in Hilaweyn.
- In Hilaweyn the Crude Mortality level is borderline but Under Five Mortality level is above emergency threshold.
- The cause-specific deaths are taken from what is reported in HIS. There were 7 deaths reported this week.
- Unreported home deliveries continue to be of concern though clean delivery kits have been provided to all visibly pregnant women.
- ACF has a third satellite MAM/BF site operational.
- ACF has 2 satellite MAM/BF sites operational.

Public Health Priorities:

- Immediate priority remains reduction of excess mortality in Hilaweyn.
- Nutrition survey planned for 15th October 2011
- Closely monitor and strengthen community-based mortality surveillance.
- Strengthen active case finding, defaulter tracing and referral as a key factor in reduction of CMR.
- Active mobilization and awareness raising on the importance of health facility delivery.
- HIS meeting and on-job training for all health and nutrition partners planned for 12th October 2011

Legend: ✓ Standard reached ⚠ Standard borderline ✗ Standard not reached ? Data unreliable i Indicator cannot be calculated n/a Data not available - Data not applicable

View interactive maps and statistics online: <http://his.unhcr.org/main.locsis>

Public Health						
Health Impact	No	Indicator	Emergency Standard	Outbreak Alert and Response	Indicator	Emergency Standard
Crude Mortality Rate (/10,000/day)		1.2	< 1 ✘	Number of outbreaks reported	0	
Under-five Mortality Rate (/10,000/day)		3.1	< 2 ✘	% of outbreaks investigated < 48 hours	-	100%
Infant Mortality Rate (IMR) (/1000 livebirths)		n/a	< 60 i	EPI and Vitamin A		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)		n/a	< 40 i	Measles vaccination coverage	>95%	> 95% ✔
Global Acute Malnutrition Rate (%)		44%	< 10% ✘	Supervision		
Severe Acute Malnutrition Rate (%)		26%	< 2% ✘	Do regular camp coordination meetings take place?	Yes	Yes ✔
Access and Utilisation				Were any drug shortages reported during the period?	No	No ✔
No. of health facilities	1	1 : 20,774	1 : <10,000 ✘			
No. of consultations per trained clinician per day		30	< 50 ✔			
Health Utilization Rate (new visits/person/year)		2.1	1 - 4 ✔			

Nutrition	Moderate Acute Malnutrition (MAM)	Emergency Standard	Severe Acute Malnutrition (SAM)					
			Outpatient therapeutic program (OTP)*	Emergency Standard	Stabilisation Centre (SC)*	Emergency Standard	CMAM (SC and OTP combined)*	Emergency Standard
Number of new admissions	590		165		34		198	
Average length of stay	n/a	< 8 weeks i	n/a	< 30 days i	n/a	< 10 days i	n/a	< 30 days i
Average weight gain (g/kg/day)	-		n/a	> 5 i	n/a		n/a	> 5 i
Discharge rate	46%	> 75% ✘	34%	> 75% ✘	81%		33%	> 75% ✘
Death rate	0%	< 3% ✔	n/a	< 10% i	6%		1%	< 10% ✔
Default rate	18%	< 15% ✘	65%	< 15% ✘	13%		66%	< 15% ✘
Referral rate	18%		0%		0%		0%	
Non-cured rate	-		0%		-		0%	

* no marasmus or kwashiorkor exits were recorded in SC or OTP during the reporting period

Reproductive Health and HIV							
Maternal and Newborn Health	No	Indicator	Emergency Standard	Sexual and Gender-based Violence	No	Indicator	Emergency Standard
No. of basic EmOC facilities	1	1 : 20,774	1 : <500,000 ✔	Incidence of reported rape (/10,000/year)	0	0.0	?
No. of comprehensive EmOC facilities	0	0	1 : <500,000 ✘	% rape survivors who received PEP < 72h	-	100%	
Number of maternal deaths		0		% rape survivors who received ECP < 120h	-	100%	
Number of maternal deaths investigated <48 hrs		-	100%	% rape survivors who received STI < 2 wks	-	100%	
Crude Birth Rate (CBR) (/1000/month)		0.4		HIV/AIDS			
Coverage complete antenatal care (> 4 visits)		n/a	> 90% i	Condom distribution rate	n/a	> 0.5 i	
% deliveries performed by caesarean section		0%	5 - 15% ✘	% of blood units screened for HIV	n/a	100% i	
% deliveries attended by skilled personnel		100%	≥ 50% ✔				
% low birth weight deliveries		0%	< 15% ✔				

PEP: Post-exposure prophylaxis; ECP: Emergency Contraceptive Pill; STI: Sexually Transmitted Infection

Community Health Activities			Water, Sanitation and Hygiene		
	Indicator	Emergency Standard		Indicator	Emergency Standard
No. of health education and hygiene sessions conducted	-		Avg quantity of potable water / person / day (litres)	16	> 10 ✔
No. of sanitation campaigns conducted	-		No. of persons per usable water tap	329	< 250 ✘
No. pregnant women who received clean delivery kit	-		No. of persons per drop-hole in communal latrine	127	≤ 50 ✘
No. of complicated medical cases identified	-		% of population living within 200m from water point	76%	100% ✘
% of complicated medical cases referred	-	≥ 90%	% of families with latrines	39%	> 50% ✘

Legend: ✔ Standard reached ⚠ Standard borderline ✘ Standard not reached ? Data unreliable i Indicator cannot be calculated n/a Data not available - Data not applicable

View interactive maps and statistics online: <http://his.unhcr.org/main.locsis>

UNHCR gratefully acknowledges the support of the following partners:

