

1. Population

| Population | 0 to < 1 | | Total <1 | 1 to 4 | | Total 1 to 4 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total |
|--------------------|----------|-------|----------|--------|-------|--------------|---------|--------|---------------|----------|--------|----------------|-------|-------|------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | | | |
| Refugee population | 2,251 | 2,251 | 4,502 | 6,859 | 7,931 | 14,790 | 19,827 | 19,291 | 39,118 | 20,363 | 25,293 | 45,656 | 1,179 | 1,929 | 3,108 | 107,174 |
| % by age group | | | 4% | | | 14% | | | 36% | | | 43% | | | 3% | 100% |

2. Mortality

2.1 Mortality

| Record ONE cause of mortality per case | 0 to 4 | | Total <5 | US % mort | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | Crude % mort |
|---|----------|----------|----------|-------------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| 1. Lower Respiratory Tract Infection (LRTI) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 2. Watery diarrhoea | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Bloody diarrhoea | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4. Cardiovascular disorders | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 1 | 1 | 2 | 0 | 0 | 0 | 2 | 33% |
| 5. Ischaemic heart disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 6. Injury | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 7. Cancer | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 8. Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 9. Tuberculosis (confirmed) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 10. Neonatal death | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 0 | 1 | 1 | 100% | 1 | 0 | 1 | 0 | 0 | 0 | 2 | 2 | 2 | 4 | 67% |
| Total | 0 | 1 | 1 | 100% | 1 | 0 | 1 | 1 | 1 | 2 | 0 | 2 | 2 | 6 | 100% |
| % by age group | | | 17% | | | | 17% | | | 33% | | 33% | | 100% | |

| Mortality indicators | Male | Female | Total |
|--------------------------------|------|--------|-------|
| Crude Mortality Rate (CMR) | 0.2 | 0.3 | 0.2 |
| Under 5 Mortality Rate (USMR) | 0.0 | 0.4 | 0.2 |
| Neonatal Mortality Rate (NNMR) | | | 0.0 |

2.2 Maternal death

| | |
|--|---|
| Number of maternal deaths reported | 0 |
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3. Morbidity

3.1 Staffing and service availability

| | |
|---|--------|
| Number of full-time clinicians | 66 |
| Average number of full days on which each health facility was functioning | 6.0 |
| Total number of outpatient consultations | 19,638 |
| Health facility utilisation rate | 9.5 |
| Consultations per clinician per day | 50 |

3.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total <5 | US Incid | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause | Crude Incide |
|--|-------------|-------------|-------------|--------------|-------------|-------------|---------------|-------------|-------------|----------------|-------------|------------|-------------|--------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | | |
| Upper Respiratory Tract Infection (URTI) | 510 | 502 | 1012 | 52.5 | 641 | 603 | 1244 | 557 | 505 | 1062 | 255 | 248 | 503 | 3821 | 27% | 35.7 |
| Influenza-like illness (ILI) | 170 | 149 | 319 | 16.5 | 155 | 177 | 332 | 280 | 264 | 544 | 108 | 111 | 219 | 1414 | 10% | 13.2 |
| Lower Respiratory Tract Infection (LRTI) | 153 | 153 | 306 | 15.9 | 144 | 138 | 282 | 144 | 188 | 332 | 138 | 129 | 267 | 1187 | 8% | 11.1 |
| Watery diarrhoea* | 306 | 276 | 582 | 30.2 | 118 | 106 | 224 | 62 | 75 | 137 | 48 | 30 | 78 | 1021 | 7% | 9.5 |
| Bloody diarrhoea* | 21 | 25 | 46 | 2.4 | 14 | 6 | 20 | 6 | 7 | 13 | 0 | 0 | 0 | 79 | 1% | 0.7 |
| Skin infection | 37 | 50 | 87 | 4.5 | 40 | 48 | 88 | 30 | 49 | 79 | 15 | 16 | 31 | 285 | 2% | 2.7 |
| Eye infection | 67 | 50 | 117 | 6.1 | 54 | 65 | 119 | 25 | 36 | 61 | 16 | 10 | 26 | 323 | 2% | 3.0 |
| Dental Conditions | 21 | 28 | 49 | 2.5 | 48 | 54 | 102 | 164 | 123 | 287 | 34 | 20 | 54 | 492 | 4% | 4.6 |
| Intestinal Worms | 19 | 30 | 49 | 2.5 | 14 | 15 | 29 | 22 | 16 | 38 | 9 | 5 | 14 | 130 | 1% | 1.2 |
| Acute Jaundice Syndrome* | 15 | 10 | 25 | 1.3 | 11 | 16 | 27 | 7 | 1 | 8 | 0 | 0 | 0 | 60 | 0% | 0.6 |
| Measles* | 0 | 5 | 5 | 0.3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 0% | 0.0 |
| Meningitis* | 1 | 0 | 1 | 0.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% | 0.0 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Fever of unknown origin | 3 | 2 | 5 | 0.3 | 2 | 1 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 0% | 0.1 |
| Sexually Transmitted Infection (STI) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Other | 650 | 686 | 1336 | 69.3 | 784 | 793 | 1577 | 724 | 746 | 1470 | 432 | 365 | 797 | 5180 | 37% | 48.3 |
| Total | 1973 | 1966 | 3939 | 204.2 | 2025 | 2022 | 4047 | 2021 | 2010 | 4031 | 1055 | 934 | 1989 | 14006 | 100% | 130.7 |
| % by age group | | | 28% | | | | 29% | | | 29% | | 14% | | 100% | | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|--|------------|------------|------------|------------|------------|---------------|------------|------------|----------------|------------|------------|-------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| Diabetes | 0 | 0 | 0 | 8 | 5 | 13 | 132 | 158 | 290 | 153 | 164 | 317 | 620 | 13% |
| Hypertension | 0 | 0 | 0 | 1 | 4 | 5 | 156 | 145 | 301 | 171 | 162 | 333 | 639 | 14% |
| Asthma | 68 | 89 | 157 | 93 | 105 | 198 | 121 | 89 | 210 | 108 | 87 | 195 | 760 | 16% |
| Ischaemic heart disease | 2 | 1 | 3 | 4 | 7 | 11 | 21 | 32 | 53 | 40 | 27 | 67 | 134 | 3% |
| Cardiovascular disorders | 0 | 0 | 0 | 0 | 5 | 5 | 14 | 18 | 32 | 29 | 24 | 53 | 90 | 2% |
| Chronic obstructive pulmonary disease (COPD) | 1 | 2 | 3 | 3 | 0 | 3 | 26 | 26 | 52 | 26 | 15 | 41 | 99 | 2% |
| Liver disease | 0 | 3 | 3 | 2 | 4 | 6 | 13 | 12 | 25 | 9 | 3 | 12 | 46 | 1% |
| Thyroid problems | 0 | 0 | 0 | 1 | 5 | 6 | 5 | 24 | 29 | 5 | 25 | 30 | 65 | 1% |
| Cancer | 7 | 3 | 10 | 6 | 10 | 16 | 23 | 16 | 39 | 17 | 4 | 21 | 86 | 2% |
| Other | 120 | 171 | 291 | 339 | 325 | 664 | 289 | 382 | 671 | 233 | 231 | 464 | 2090 | 45% |
| Total | 198 | 269 | 467 | 457 | 470 | 927 | 800 | 902 | 1702 | 791 | 742 | 1533 | 4629 | 100% |
| % by age group | | | 10% | | | 20% | | | 37% | | | 33% | | 100% |

3.4 Mental health

| | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|---|----------|----------|-----------|-----------|-----------|---------------|-----------|-----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Epilepsy / seizures | 0 | 0 | 0 | 12 | 10 | 22 | 6 | 1 | 7 | 2 | 0 | 2 | 31 | 18% |
| 2. Alcohol and other substance abuse disorder | 0 | 0 | 0 | 0 | 7 | 7 | 3 | 0 | 3 | 0 | 0 | 0 | 10 | 6% |
| 3. Mental retardation / intellectual disability | 0 | 0 | 0 | 1 | 4 | 5 | 4 | 3 | 7 | 0 | 0 | 0 | 12 | 7% |
| 4. Psychotic disorder | 0 | 0 | 0 | 0 | 6 | 6 | 21 | 6 | 27 | 0 | 0 | 0 | 33 | 19% |
| 5. Severe emotional disorder | 1 | 0 | 1 | 2 | 1 | 3 | 14 | 13 | 27 | 2 | 0 | 2 | 33 | 19% |
| 6. Other psychological complaint | 0 | 0 | 0 | 6 | 3 | 9 | 12 | 5 | 17 | 0 | 0 | 0 | 26 | 15% |
| Other | 1 | 0 | 1 | 11 | 5 | 16 | 9 | 3 | 12 | 0 | 0 | 0 | 29 | 17% |
| Total | 2 | 0 | 2 | 32 | 36 | 68 | 69 | 31 | 100 | 4 | 0 | 4 | 174 | 100% |
| % by age group | | | 1% | | | 39% | | | 57% | | | 2% | 100% | |

3.5 Injuries

| | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause |
|-----------------------------|-----------|-----------|------------|------------|------------|---------------|------------|-----------|----------------|-----------|-----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Bites (all causes) | 4 | 7 | 11 | 4 | 8 | 12 | 11 | 6 | 17 | 15 | 10 | 25 | 65 | 8% |
| 2. Injury (war related) | 11 | 9 | 20 | 18 | 39 | 57 | 72 | 4 | 76 | 17 | 3 | 20 | 173 | 21% |
| 3. Injury (non war related) | 40 | 37 | 77 | 48 | 27 | 75 | 99 | 33 | 132 | 30 | 28 | 58 | 342 | 41% |
| 4. Accident | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 9 | 15 | 24 | 32 | 29 | 61 | 75 | 46 | 121 | 21 | 22 | 43 | 249 | 30% |
| Total | 64 | 68 | 132 | 102 | 103 | 205 | 257 | 89 | 346 | 83 | 63 | 146 | 829 | 100% |
| % by age group | | | 16% | | | 25% | | | 42% | | | 18% | 100% | |

4. In-patient Department

4.1 In-patient Department Activities

| | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | |
|--|--------------|----------|-----------|-----------|-----------|---------------|-----------|-----------|----------------|----------|----------|------------|-------------|-----|
| | M | F | | M | F | | M | F | | M | F | | | |
| Number at beginning of reporting period | 0 | 2 | 2 | 3 | 2 | 5 | 1 | 1 | 2 | 0 | 0 | 0 | 9 | |
| Total admissions | 3 | 5 | 8 | 15 | 13 | 28 | 14 | 55 | 69 | 3 | 2 | 5 | 110 | |
| Number of exits | authorized | 3 | 6 | 9 | 16 | 13 | 29 | 14 | 46 | 60 | 2 | 1 | 3 | 101 |
| | unauthorized | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | death | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | referral | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 9 | 9 | 0 | 1 | 1 | 10 |
| Total exits | 3 | 6 | 9 | 16 | 13 | 29 | 14 | 55 | 69 | 2 | 2 | 4 | 111 | |
| Deaths within 24 hours | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| % at end of reporting period by age group | | | 13% | | | 50% | | | 25% | | | 13% | 100% | |

| | |
|---|-------|
| Number of days in reporting period | 7 |
| Number of beds | 62 |
| Sum number of days stay for authorised discharges | 203.2 |
| Proportion of < 5 IPD deaths within 24 hours | |
| Average length of stay (days) | 2.0 |
| Bed occupancy | 51% |
| Hospitalisation rate | 4.1 |

4.2 In-patient Admissions and Deaths

| | 0 to 4 | | U5 CFR | 5 to 17 | | 5-17 CFR | 18 to 59 | | 18-59 CFR | > 60 | | Total adm | Total deaths | Crude CFR |
|-----------------------------|----------|----------|-----------|-----------|----------|-----------|-----------|----------|-----------|-----------|----------|------------|--------------|-----------|
| | Adm | Deaths | | Adm | Deaths | | Adm | Deaths | | Adm | Deaths | | | |
| 1. Paediatric medicine | 5 | 0 | 0% | 6 | 0 | 0% | 0 | 0 | 0% | 0 | 0 | 11 | 0 | 0% |
| 2. General Surgery | 0 | 0 | | 3 | 0 | 0% | 6 | 0 | 0% | 1 | 0 | 10 | 0 | 0% |
| 3. Injury (war related) | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | 0 | 0 | |
| 4. Injury (non war related) | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | 0 | 0 | |
| 5. Internal medicine | 0 | 0 | | 6 | 0 | 0% | 4 | 0 | 0% | 8 | 0 | 18 | 0 | 0% |
| 6. Obstetrics | 0 | 0 | | 5 | 0 | 0% | 48 | 0 | 0% | 0 | 0 | 53 | 0 | 0% |
| 7. Gynaecology | 0 | 0 | | 0 | 0 | | 0 | 0 | | 0 | 0 | 0 | 0 | |
| Other | 0 | 0 | | 2 | 0 | 0% | 6 | 0 | 0% | 7 | 0 | 15 | 0 | 0% |
| Total | 5 | 0 | 0% | 22 | 0 | 0% | 64 | 0 | 0% | 16 | 0 | 107 | 0 | 0% |

Adm. = Admission; CFR = Case Fatality Rate

6. Laboratory

| | Total |
|--------------------------------------|-------|
| Number of laboratory tests conducted | 583 |

8. Reproductive Health

8.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|--|-------------------|------|-------|
| First antenatal visit < 1st trimester | 98 | 81 | 179 |
| First antenatal visit > 1st trimester | 69 | 137 | 206 |
| Number of pregnant women at time of delivery who received: | ≥ 4 ANC visits | 0 | 8 |
| | 2 TT doses | 0 | 0 |
| | anaemia screening | 0 | 7 |

| ANC Indicators | |
|--|-----|
| Proportion of first time antenatal visits made < 1st trimester | 46% |
| Coverage of complete antenatal care | 15% |
| Coverage of antenatal tetanus immunisation | 0% |
| Coverage of anaemia screening in pregnancy | 13% |

8.2 Delivery care

| | < 18 | ≥ 18 | Total |
|--------------|------|------|-------|
| Live births | 1 | 53 | 54 |
| Still births | 0 | 1 | 1 |

| | | | |
|---|---|----|----|
| Low birth weight (<2500 g) | 0 | 1 | 1 |
| Attended by skilled health worker | 1 | 53 | 54 |
| Number of obstetric complications treated | 0 | 7 | 7 |
| Number of caesarean sections performed | 0 | 6 | 6 |

| Delivery Indicators | |
|--|------|
| Crude birth rate | 6.0 |
| Proportion of births attended by skilled health worker (excl. TBA) | 100% |
| Proportion of low birth weight deliveries | 2% |
| Proportion of all births performed by caesarean section | 11% |
| Proportion of obstetric complications treated among girls < 18 | 0% |

8.3 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|-------|
| Women who attended 3 postnatal visits within 6 wks of delivery | 0 | 4 | 4 |

| Postnatal Indicators | |
|-------------------------------------|----|
| Coverage of complete postnatal care | 7% |

8.5 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|---|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 0 | 0 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

| SGBV Indicators | |
|--|-----|
| Incidence of reported rape | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

Data Sources: 1. Health Information System

Publication date: 14 May 2013

Contact Information: **Za'atri** Ana Calvo, UNHCR Email: calvo@unhcr.org

Amman Ann Burton, UNHCR Email: burton@unhcr.org

