

1. Population

Population	0 to <1		Total <1	1 to 4		Total 1 to 4	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total
	M	F		M	F		M	F		M	F		M	F		
Refugee population	95	95	190	378	379	757	838	678	1,516	1,505	1,257	2,762	427	388	815	6,040
% by age group			3%			13%			25%			46%			13%	100%

* estimates from registered population

2. Mortality

2.1 Mortality

Record ONE cause of mortality per case	0 to 4		Total <5	U5 % mort	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total > 50	Grand total	Crude % mort
	M	F			M	F		M	F		M	F			
1. Lower Respiratory Tract Infection (LRTI)	0	0	0		0	0	0	0	0	0	0	0	0	0	
2. Watery diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	
3. Bloody diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	
4. Cardiovascular disorders	0	0	0		0	0	0	0	0	0	0	0	0	0	
5. Ischaemic heart disease	0	0	0		0	0	0	0	0	0	0	0	0	0	
6. Injury	0	0	0		0	0	0	0	0	0	0	0	0	0	
7. Cancer	0	0	0		0	0	0	0	0	0	0	0	0	0	
8. Chronic obstructive pulmonary disease (COPD)	0	0	0		0	0	0	0	0	0	0	0	0	0	
9. Tuberculosis (confirmed)	0	0	0		0	0	0	0	0	0	0	0	0	0	
10. Neonatal death	0	0	0		0	0	0	0	0	0	0	0	0	0	
Other	0	0	0		0	0	0	0	0	0	0	0	0	0	
Total	0	0	0	100%	0	0	0	0	0	0	0	0	0	0	100%
% by age group															

Mortality indicators	Male	Female	Total
Crude Mortality Rate (CMR)	0.0	0.0	0.0
Under 5 Mortality Rate (USMR)	0.0	0.0	0.0
Neonatal Mortality Rate (NNMR)			0.0

2.2 Maternal death

Number of maternal deaths reported	0
Number of maternal deaths investigated	0
Proportion of maternal deaths investigated	

3. Morbidity

3.1 Staffing and service availability

Number of full-time clinicians	3
Average number of full days on which each health facility was	7.0
Total number of outpatient consultations	878
Health facility utilisation rate	7.6
Consultations per clinician per day	42

3.2 Acute health conditions

Record NEW cases of disease only	0 to 4		Total <5	U5 Incid**	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause	Crude Incid**
	M	F			M	F		M	F		M	F				
Upper Respiratory Tract Infection (URTI)	26	31	57	60.2	31	17	48	84	44	128	6	13	19	252	32%	41.7
Lower Respiratory Tract Infection (LRTI)	42	48	90	95.0	20	31	51	15	12	27	2	1	3	171	22%	28.3
Watery diarrhoea*	2	3	5	5.3	1	1	2	1	3	4	0	0	0	11	1%	1.8
Bloody diarrhoea*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Malaria (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Skin infection	8	6	14	14.8	4	8	12	12	9	21	0	1	1	48	6%	7.9
Eye infection	4	4	8	8.4	1	3	4	6	6	12	0	2	2	26	3%	4.3
Ear infection	3	1	4	4.2	2	1	3	3	7	10	0	1	1	18	2%	3.0
Dental Conditions	0	0	0	0.0	8	5	13	18	19	37	4	3	7	57	7%	9.4
Intestinal Worms	0	0	0	0.0	0	1	1	0	0	0	0	0	0	1	0%	0.2
Acute Jaundice Syndrome*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Measles (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Meningitis (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute Flaccid Paralysis*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Tuberculosis (suspected)	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Fever of unknown origin	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Sexually Transmitted Infection (STI)	0	0	0	0.0	0	0	0	0	86	86	0	0	0	86	11%	14.2
Urinary tract infection (UTI)	0	0	0	0.0	2	3	5	1	10	11	0	1	1	17	2%	2.8
HIV/AIDS	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute malnutrition	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Leishmaniasis	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Other	0	0	0	0.0	1	2	3	23	62	85	5	8	13	101	13%	16.7
Total	85	93	178	188.0	70	72	142	163	258	421	17	30	47	788	100%	130.5
% by age group				23%			18%			53%			6%	100%		

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **incidence is cases per 1000 per week

3.3 Chronic health conditions

Record NEW and REVISITS	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
Diabetes	0	0	0	0	0	0	0	1	1	1	4	5	6	8%
Hypertension	0	0	0	0	0	0	4	3	7	3	7	10	17	22%
Asthma	1	2	3	0	0	0	0	3	3	2	0	2	8	10%
Ischaemic heart disease	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Cardiovascular disease (other)	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Chronic obstructive pulmonary disease (COPD)	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Liver disease	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Thyroid problems	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Cancer	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	0	0	0	1	0	1	11	19	30	8	8	16	47	60%
Total	1	2	3	1	0	1	15	26	41	14	19	33	78	100%
% by age group				4%		1%			53%			42%	100%	

3.4 Mental health

	0 to 4			5 to 14			15 to 49			≥ 50			Total ≥ 50	Grand total	% by cause
	M	F	Total < 5	M	F	Total 5 to 14	M	F	Total 15 to 49	M	F	Total ≥ 50			
1. Epilepsy / seizures	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
3. Psychotic disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4. Mental retardation / intellectual disability	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5. Severe emotional disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6. Medically unexplained somatic complaint	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	0	0	0	0	0	0	0	0	0	0	0	0	0	0	100%
% by age group															100%

3.5 Injuries

	0 to 4			5 to 14			15 to 49			≥ 50			Total ≥ 50	Grand total	% by cause
	M	F	Total < 5	M	F	Total 5 to 14	M	F	Total 15 to 49	M	F	Total ≥ 50			
1. Injury (intentional)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
2. Injury (unintentional)	2	0	2	0	0	0	0	0	0	0	0	0	0	2	17%
3. Burns	2	4	6	1	2	3	0	0	0	0	0	0	0	9	75%
4. Accident	0	0	0	1	0	1	0	0	0	0	0	0	0	1	8%
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Total	4	4	8	2	2	4	0	0	0	0	0	0	0	12	100%
% by age group			67%			33%			0%					0%	100%

4. Referrals

4.1 Referrals - general

	0 to 4	5 to 14	15 to 49	≥ 50	Grand total
Emergency	0	0	0	0	0
Elective	0	0	0	0	0
Other	0	0	0	0	0
Total	0	0	0	0	0
% by age group					100%

ND: no data submitted

5. Laboratory & other investigations

	Total
Haematology	0
Imaging (X-ray, MRI/CT)	0
Other	0
Total	0

6. EPI and Vitamin A

	<1	1 to <5	Total
BCG	0	0	0
Hepatitis B birth dose	0	0	0
DPT I/HiB I/Hep B I* (Pentavalent I)	1	0	1
DPT II/HiB II* (Tetravalent)	9	0	9
DPT III/HiB III/Hep B II* (Pentavalent II)	4	0	4
Polio 0	0	0	0
Polio I	1	0	1
Polio II	9	0	9
Polio III	4	0	4
Measles	4	0	4
MMR	0	5	5
Rotavirus 1	1	0	1
Rotavirus 2	3	0	3
Rotavirus 3	2	0	2
Vitamin A	4	6	10
Additional vaccines	0	15	15

EPI Indicators (since December 2013)		
Total vaccinated this week		68
a. Program dropout rate*		-14%
b. Immunisation coverage		
BCG		52%
Hepatitis B birth dose		61%
DPT I		63%
DPT III		82%
Polio I		78%
Polio III		151%
Measles		99%
MMR**		39%
Rotavirus I		39%
Rotavirus III		95%
c. Vitamin A coverage		

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%
**For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

7. Reproductive Health

7.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester			
First antenatal visit > 1st trimester			
Number of pregnant women at time of delivery who received:	NO DATA		
≥ 4 ANC visits			
2 TT doses			
anaemia screening			

ND: no data submitted

ANC Indicators	
Proportion of first time antenatal visits made < 1st trimester	

7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery			NO DATA

ND: no data submitted

6.3 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No of rape survivors given PEP within 72 hours	0	0	0	0	0
No of female rape survivors given ECP within 120 hours					0
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators		
Incidence of reported rape		0.0
Proportion of rape survivors who receive PEP < 72 hrs		
Proportion of female rape survivors who receive ECP < 120 hrs		
Proportion of rape survivors who receive STI presumptive treatment < 2 wks		

Data Sources: 1. Health Information System

Publication date: 03 February 2014

Contact Information: Erbil, Dr Chimmen Akrawi, DOH Erbil

Email: chimmentya@yahoo.com

Erbil, Mohammed Marzooq, UNHCR

Email: MARZOOQ@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

