

1. Population

Population	0 to < 1		Total <1	1 to 4		Total 1 to 4	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total
	M	F		M	F		M	F		M	F		M	F		
Refugee population	57	50	107	228	202	430	435	380	815	646	637	1,283	181	188	369	3,004
% by age group			4%			14%			27%			43%			12%	100%

* estimates from registered population

2. Mortality

2.1 Mortality

Record ONE cause of mortality per case	0 to 4		Total <5	U5 % mort	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	Crude % mort
	M	F			M	F		M	F		M	F			
1. Lower Respiratory Tract Infection (LRTI)	0	0	0		0	0	0	0	0	0	0	0	0	0	
2. Watery diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	
3. Bloody diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	
4. Cardiovascular disorders	0	0	0		0	0	0	0	0	0	0	0	0	0	
5. Ischaemic heart disease	0	0	0		0	0	0	0	0	0	0	0	0	0	
6. Injury	0	0	0		0	0	0	0	0	0	0	0	0	0	
7. Cancer	0	0	0		0	0	0	0	0	0	0	0	0	0	
8. Chronic obstructive pulmonary disease (COPD)	0	0	0		0	0	0	0	0	0	0	0	0	0	
9. Tuberculosis (confirmed)	0	0	0		0	0	0	0	0	0	0	0	0	0	
10. Neonatal death	0	0	0		0	0	0	0	0	0	0	0	0	0	
Other	0	0	0		0	0	0	0	0	0	0	0	0	0	
Total	0	0	0	100%	0	0	0	0	0	0	0	0	0	0	100%
% by age group															

Mortality indicators	Male	Female	Total
Crude Mortality Rate (CMR)	0.0	0.0	0.0
Under 5 Mortality Rate (U5MR)	0.0	0.0	0.0
Neonatal Mortality Rate (NNMR)			0.0

2.2 Maternal death

Number of maternal deaths reported	0
Number of maternal deaths investigated	0
Proportion of maternal deaths investigated	

3. Morbidity

3.1 Staffing and service availability

Number of full-time clinicians	2
Average number of full days on which each health facility was	7.0
Total number of outpatient consultations	857
Health facility utilisation rate	14.8
Consultations per clinician per day	61

3.2 Acute health conditions

Record NEW cases of disease only	0 to 4		Total <5	U5 Incid**	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause	Crude Incid**
	M	F			M	F		M	F		M	F				
Upper Respiratory Tract Infection (URTI)	85	48	133	247.7	63	48	111	101	110	211	5	9	14	469	56%	156.1
Lower Respiratory Tract Infection (LRTI)	10	7	17	31.7	1	2	3	7	6	13	0	2	2	35	4%	11.7
Watery diarrhoea*	6	8	14	26.1	4	2	6	7	6	13	0	2	2	35	4%	11.7
Bloody diarrhoea*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Malaria (suspected)*	3	1	4	7.4	1	0	1	1	0	1	0	0	0	6	1%	2.0
Skin infection	11	6	17	31.7	2	4	6	12	9	21	1	2	3	47	6%	15.6
Eye infection	0	0	0	0.0	1	0	1	5	6	11	1	0	1	13	2%	4.3
Ear infection	0	0	0	0.0	3	3	6	5	13	18	0	1	1	25	3%	8.3
Dental Conditions	0	0	0	0.0	1	1	2	1	0	1	0	0	0	3	0%	1.0
Intestinal Worms	0	0	0	0.0	1	0	1	0	0	0	0	0	0	1	0%	0.3
Acute Jaundice Syndrome*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Measles (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Meningitis (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute Flaccid Paralysis*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Tuberculosis (suspected)	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Fever of unknown origin	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Sexually Transmitted Infection (STI)	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Urinary tract infection (UTI)	2	1	3	5.6	4	4	8	6	8	14	5	1	6	27	3%	9.0
HIV/AIDS	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute malnutrition	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Leishmaniasis	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Other	11	10	21	39.1	4	4	8	53	74	127	15	1	16	172	21%	57.3
Total	128	81	209	389.2	81	68	149	198	232	430	27	18	45	833	100%	277.3
% by age group			25%				18%			52%			5%	100%		

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **Incidence is cases per 1000 per week

3.3 Chronic health conditions

Record NEW and REVISITS	0 to 4		Total < 5	5 to 14		Total 5 to 14	15 to 49		Total 15 to 49	≥ 50		Total ≥ 50	Grand total	% by cause
	M	F		M	F		M	F		M	F			
Diabetes	0	0	0	0	0	0	0	2	2	1	5	6	8	42%
Hypertension	0	0	0	0	0	0	0	0	0	4	5	9	9	47%
Asthma	0	0	0	0	0	0	0	0	0	0	2	2	2	11%
Ischaemic heart disease	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Cardiovascular disease (other)	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Chronic obstructive pulmonary disease (COPD)	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Liver disease	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Thyroid problems	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Cancer	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Total	0	0	0	0	0	0	2	2	2	5	12	17	19	100%
% by age group			0%			0%			11%			89%	100%	

3.4 Mental health

	0 to 4			5 to 14			15 to 49			≥ 50			Total ≥ 50	Grand total	% by cause
	M	F	Total < 5	M	F	Total 5 to 14	M	F	Total 15 to 49	M	F	Total ≥ 50			
1. Epilepsy / seizures	0	0	0	0	1	1	0	0	0	0	0	0	0	1	50%
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
3. Psychotic disorder	0	0	0	0	0	0	0	0	0	1	0	1	1	1	50%
4. Mental retardation / intellectual disability	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
5. Severe emotional disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
6. Medically unexplained somatic complaint	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Total	0	0	0	0	1	1	0	0	0	1	0	1	2	2	100%
% by age group			0%			50%			0%			50%	100%		

3.5 Injuries

	0 to 4			5 to 14			15 to 49			≥ 50			Total ≥ 50	Grand total	% by cause
	M	F	Total < 5	M	F	Total 5 to 14	M	F	Total 15 to 49	M	F	Total ≥ 50			
1. Injury (intentional)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
2. Injury (unintentional)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
3. Burns	0	0	0	0	0	0	0	2	2	1	0	1	3	3	100%
4. Accident	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Other	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
Total	0	0	0	0	0	0	0	2	2	1	0	1	3	3	100%
% by age group			0%			0%			67%			33%	100%		

4. Referrals

4.1 Referrals - general

	0 to 4	5 to 14	15 to 49	≥ 50	Grand total
Emergency	4	1	0	1	6
Elective	0	0	4	0	4
Other	4	3	32	0	39
Total	8	4	36	1	49
% by age group	16%	8%	73%	2%	100%

ND: no data submitted

5. Laboratory & other investigations

	Total
Haematology	0
Imaging (X-ray, MRI/CT)	0
Other	0
Total	4

6. EPI and Vitamin A

	<1	≥1 to <5	Total
BCG	2	0	2
Hepatitis B birth dose	2	0	2
DPT I/HiB I/Hep B I* (Pentavalent I)	2	0	2
DPT II/HiB II* (Tetavalent)	3	0	3
DPT III/HiB III/Hep B II* (Pentavalent II)	2	0	2
Polio 0	2	0	2
Polio I	0	0	0
Polio II	3	0	3
Polio III	2	0	2
Measles	0	0	0
MMR	0	0	0
Rotavirus 1	0	0	0
Rotavirus 2	2	0	2
Rotavirus 3	1	0	1
Vitamin A	0	0	0
Additional vaccines	0	0	0

EPI Indicators (since December 2013)	
Total vaccinated this week	21
a. Program dropout rate*	-88%
b. Immunisation coverage	
BCG	57%
Hepatitis B birth dose	52%
DPT I	42%
DPT III	78%
Polio I	31%
Polio III	78%
Measles	26%
MMR**	63%
Rotavirus I	5%
Rotavirus III	42%
c. Vitamin A coverage	52%

*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines

Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%

**For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

7. Reproductive Health

7.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester	NO DATA		
First antenatal visit > 1st trimester			
Number of pregnant women at time of delivery who received:			
≥ 4 ANC visits			
2 TT doses			
anaemia screening			

ND: no data submitted

ANC Indicators	
Proportion of first time antenatal visits made < 1st trimester	

7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery	NO DATA		

ND: no data submitted

7.3 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	0	0
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 - 120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	0	0
No. of rape survivors given PEP within 72 hours	0	0	0	0	0
No. of female rape survivors given ECP within 120 hours		0		0	0
No. of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	0	0

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators	
Incidence of reported rape	0.0
Proportion of rape survivors who receive PEP < 72 hrs	
Proportion of female rape survivors who receive ECP < 120 hrs	
Proportion of rape survivors who receive STI presumptive treatment < 2 wks	