




Refugee population

Total refugee population:

98,973 *

Demographic breakdown:



Key facts

Camp opened:
July 2012

* Source of population data:
WFP Distribution Figures

Indicators at a glance

1 Health utilisation rate	5.5	✓
2 Number of consultations per clinician per day	23	✓
3 Hospitalisation rate	3.4	⚠
4 Referral rate		?
5 % deliveries attended by skilled personnel	100%	⚠
6 Crude Mortality Rate (CMR) (/1000/month)	0.2	✓
7 Under-five Mortality Rate (U5MR) (/1000/month)	0.0	✓
8 Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0.0	✗

Notes


This public health profile for Za'atri refugee camp is compiled from health facility data from the following partners:

- IMC Clinic
- JHAS Clinic
- Jordan Italian Field Hospital
- MDM Clinics 1 and 2
- Moroccan Field Hospital
- Saudi Clinic

Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Morbidity

Acute health conditions (%)





Top 5 acute health conditions

	% morbidity	# of cases
1 Upper Respiratory Tract Infection (URTI)	31%	2,210
2 Influenza-like illness (ILLI)	14%	1,022
3 Skin infection	9%	645
4 Lower Respiratory Tract Infection (LRTI)	5%	368
5 Dental Conditions	5%	376
- Other	35%	3,203


Breakdown of acute health conditions

by age

by gender



Chronic health conditions (%)





Top 5 chronic health conditions

	% morbidity	# of cases
1 Diabetes	12%	335
2 Hypertension	10%	296
3 Asthma	9%	259
4 Ischaemic heart disease	2%	66
5 Thyroid problems	1%	34
- Other	65%	1,846

Breakdown of chronic health conditions

by age

by gender



UNHCR gratefully acknowledges the support of the following partners:



Public Health

Health Impact

	Indicator	Standard	
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 1.5	✓
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 3.0	✓
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0.0	< 40	✓

Access and utilization

No. of health facilities	7	1 : 16496	1 : < 10,000	✗
Total number of consultations	10,543			
Health utilisation rate	5.5	1 - 4		✗
No. of consultations per trained clinician per day	23	< 50		✓

Outbreak Alert and Response

	Indicator	Standard
Number of outbreaks reported	0	
% of outbreaks investigated within 48 hours		100%

Laboratory


Number of laboratory samples	444
------------------------------	-----

Immunization


Measles vaccination rate	> 95%	?
Full vaccination rate	> 95%	?
Program drop-out rate	< 10%	?
Vaccine wastage rate	< 8%	?

Breakdown of mental health conditions


By cause



By age




By gender




Breakdown of injuries


By cause



By age



By gender



In-patient and referral

In-patient department

	Indicator	Standard	
Hospitalisation rate (/1000/month)	3.4	50 - 150	✗
Bed occupancy rate	58%	> 75%	✗
Mean length of stay for authorised discharges (days)	1.7		

	Indicator	Standard
Overall case fatality rate	0%	
Proportion of deaths within 24 hours of admission	0%	

Referral

Total referral rate	?
---------------------	---

Reproductive Health

Antenatal care

	Indicator	Standard	
Coverage of antenatal care (4 or more visits)	38%	≥ 90%	✓
Coverage of TT vaccination	15%	≥ 90%	✗
Coverage of anaemia screening in pregnancy	52%	≥ 90%	✓

Delivery care

Proportion of deliveries attended by skilled personnel	100%	≥ 90%	⚠
Proportion of deliveries performed by caesarean section	19%	5 - 15%	✓
Proportion of low birth weight deliveries (<2500g)	0%	< 15%	✓

Maternal mortality

Number of maternal deaths reported	0		
% of maternal deaths investigated within 48 hours		100%	

Postnatal care

	Indicator	Standard	
Coverage of postnatal care	53%	≥ 90%	⚠

Family Planning

Contraceptive prevalence rate	2%		
-------------------------------	----	--	--

Sexual and Gender-based Violence

Incidence of rape (reported cases/10,000/year)*	0		✗
% of rape survivors who received PEP** < 72 hours		100%	✗
% of rape survivors who received ECP*** < 120 hours		100%	✗
% of rape survivors who received ST**** prophylaxis < 2 wks		100%	✗

*Only includes rapes reported at health facilities **Post-exposure prophylaxis ***Emergency contraceptive pill ****Sexually transmitted infection