Kawergosk camp | Erbil, Kurdistan Region, Iraq

1. Population																
Banadatian	0 to	< 1	Total	1 te	o 4	Total 1	5 to	14	Total 5	15 t	o 49	Total	≥ 5	50	Total ≥	Grand
Population	м	F	<1	м	F	to 4	м	F	to 14	м	F	15 to 49	м	F	50	total
Refugee population	236	216	452	945	862	1,807	1,568	1,442	3,010	3,288	2,895	6,183	976	915	1,891	13,343
% by age group			3%			14%			23%			46%			14%	100%
* estimates from registered population. For multiweek reports, mean population is presented																

2. Mortality

	0 t	o 4	Total	U5 %	5 to	14	Total 5	15 t	o 49	Total 15	≥ 5	0	Total >	Grand	Crude %
Record ONE cause of mortality per case	м	F	<5	mort	м	F	to 14	м	F	to 49	м	F	50	total	mort
1. Lower Respiratory Tract Infection (LRTI)	0	0	0		0	0	0	0	0	0	0	0	0	0	
2. Watery diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	
3. Bloody diarrhoea	0	0	0		0	0	0	0	0	0	0	0	0	0	
4. Cardiovascular disorders	0	0	0		0	0	0	0	0	0	0	0	0	0	
5. Ischaemic heart disease	0	0	0		0	0	0	0	0	0	0	0	0	0	
6. Injury	0	0	0		0	0	0	0	0	0	0	0	0	0	
7. Cancer	0	0	0		0	0	0	0	0	0	0	0	0	0	
8. Chronic obstructive pulmonary disease (COPD)	0	0	0		0	0	0	0	0	0	0	0	0	0	
9. Tuberculosis (confirmed)	0	0	0		0	0	0	0	0	0	0	0	0	0	
0. Neonatal death	0	0	0		0	0	0	0	0	0	0	0	0	0	
Other	0	0	0		0	0	0	0	0	0	0	0	0	0	
Total	0	0	0	100%	0	0	0	0	0	0	0	0	0	0	100%
% by age group														100%	l
ality indicators	Male	Female	Total												
le Mortality Rate (CMR)	0.0	0.0	0.0												
er 5 Mortality Rate (U5MR)	0.0	0.0	0.0												
natal Mortality Rate (NNMR)			0.0												

2.2 Maternal death

Number of maternal deaths reported	0
Number of maternal deaths investigated	0
Proportion of maternal deaths investigated	

3. Morbidity

3.1 Staffing and service availability

Average number of full-time clinicians	6.2
Average number full days each health facility was functioning	16.0
Total number of outpatient consultations	3,667
Health facility utilisation rate	3.6
Consultations per clinician per day	37

3.2 Acute health conditions

	0 to	b 4	Total	U5	5 to	14	Total 5	15 to	49	Total 15	≥ 50	D	Total ≥	Grand	% by	Crude
Record NEW cases of disease only	м	F	<5	Incid**	м	F	to 14	м	F	to 49	м	F	50	total	cause	Incid**
Upper Respiratory Tract Infection (URTI)	284	247	531	58.8	184	137	321	135	201	336	5	7	12	1200	36%	22.5
Lower Respiratory Tract Infection (LRTI)	79	55	134	14.8	34	7	41	16	17	33	6	2	8	216	6%	4.0
Watery diarrhoea*	82	59	141	15.6	11	11	22	12	10	22	2	1	3	188	6%	3.5
Bloody diarrhoea*	5	7	12	1.3	0	0	0	1	0	1	0	0	0	13	0%	0.2
Malaria (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Skin infection	25	19	44	4.9	17	10	27	43	60	103	5	2	7	181	5%	3.4
Eye infection	12	12	24	2.7	22	7	29	27	26	53	0	2	2	108	3%	2.0
Ear infection	36	42	78	8.6	17	17	34	29	30	59	1	0	1	172	5%	3.2
Dental Conditions	3	6	9	1.0	16	5	21	42	43	85	3	2	5	120	4%	2.2
Intestinal Worms	3	8	11	1.2	7	8	15	5	5	10	0	0	0	36	1%	0.7
Acute Jaundice Syndrome*	0	0	0	0.0	0	0	0	1	0	1	0	0	0	1	0%	0.0
Measles (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Meningitis (suspected)*	0	1	1	0.1	0	0	0	0	0	0	0	0	0	1	0%	0.0
Acute Flaccid Paralysis*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Tuberculosis (suspected)	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Fever of unknown origin	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Sexually Transmitted Infection (STI)	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Urinary tract infection (UTI)	7	21	28	3.1	7	21	28	20	56	76	0	3	3	135	4%	2.5
HIV/AIDS	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Acute malnutrition	1	0	1	0.1	0	0	0	0	0	0	0	0	0	1	0%	0.0
Leishmaniasis	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Other	71	57	128	14.2	72	42	114	223	422	645	38	52	90	977	29%	18.3
Total	608	534	1142	126.4	387	265	652	554	870	1424	60	71	131	3349	100%	62.7
% by age group			34%				<mark>19%</mark>			<mark>43%</mark>			4%	<mark>100%</mark>		

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **incidence is cases per 1000 per week

3.3 Chronic health conditions

	0 t	o 4	Total <	5 to	o 14	Total 5	15 to	o 49	Total 15	≥ 5	0	Total	Grand	% by
Record NEW and REVISITS	м	F	5	м	F	to 14	м	F	to 49	м	F	≥ 50	total	cause
Diabetes	0	0	0	0	0	0	8	18	26	7	18	25	51	29%
Hypertension	0	0	0	0	0	0	8	30	38	7	19	26	64	36%
Asthma	4	2	6	0	0	0	3	3	6	1	0	1	13	7%
Ischaemic heart disease	0	0	0	0	0	0	3	4	7	1	1	2	9	5%
Cardiovascular disease (other)	0	0	0	0	1	1	5	10	15	3	3	6	22	13%
Chronic obstructive pulmonary disease (COPD)	0	0	0	0	0	0	1	0	1	1	0	1	2	1%
Liver disease	0	0	0	0	0	0	0	2	2	0	0	0	2	1%
Thyroid problems	0	0	0	0	0	0	0	4	4	0	2	2	6	3%
Cancer	0	0	0	0	0	0	0	6	6	0	0	0	6	3%
Other	0	1	1	0	0	0	0	0	0	0	0	0	1	1%
Total	4	3	7	0	1	1	28	77	105	20	43	63	176	100%
% by age group			4%			1%			<mark>60%</mark>			36%	100%	

3.4 Mental health

	0 t	o 4	Total <	5 to	o 14	Total 5	15 to	o 49	Total 15 to	≥ (50	Total	Grand	% by
	м	F	5	м	F	to 14	М	F	49	М	F	≥ 50	total	cause
1. Epilepsy / seizures	0	0	0	0	1	1	1	2	3	0	0	0	4	15%
2. Alcohol and other substance abuse disorder	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
3. Psychotic disorder	0	0	0	0	0	0	5	0	5	0	0	0	5	19%
4. Mental retardation / intellectual disability	0	0	0	0	0	0	0	0	0	0	0	0	0	0%
5. Severe emotional disorder	0	0	0	0	0	0	1	4	5	0	0	0	5	19%
6. Medically unexplained somatic complaint	0	0	0	0	0	0	0	1	1	0	0	0	1	4%
Other	3	0	3	1	0	1	1	5	6	0	2	2	12	44%
Total	3	0	3	1	1	2	8	12	20	0	2	2	27	100%
% by age group			11%			7%			74%			7%	100%	

3.5 Injuries

	0 te	o 4	Total <	5 to	o 14	Total 5	15 t	o 49	Total 15 to	≥ :	50	Total	Grand	% by
	м	F	5	м	F	to 14	м	F	49	м	F	≥ 50	total	cause
1. Injury (intentional)	0	0	0	2	1	3	4	0	4	0	0	0	7	6%
2. Injury (unintentional)	3	4	7	11	8	19	33	6	39	3	0	3	68	59%
3. Burns	3	1	4	2	1	3	3	2	5	0	1	1	13	11%
4. Accident	1	1	2	1	0	1	9	0	9	0	0	0	12	10%
Other	2	1	3	3	1	4	6	2	8	0	0	0	15	13%
Total	9	7	16	19	11	30	55	10	65	3	1	4	115	100%
% by age group			14%			26%			57%			3%	100%	

4. Referrals

5. Laboratory & other investigations

4.1 Referrals - general

	0 to 4	5 to 14	15 to 49	2 50	Grand total		Total
Emergency	2	2	9	1	14		
Elective	11	9	49	12	81	Haematology	0
Other	0	0	0	0	0	Imaging (X-ray, MRI/CT)	0
Total	13	11	58	13	95	Other	0
% by age group	14%	12%	61%	14%	100%	Total	6
ND: no data submitted	·			-			
EDL and Mitagenie A							

6. EPI and Vitamin A

	<1	1 to <5	Total
BCG	(0 0	0
Hepatitis B birth dose	(0 0	0
DPT I/HiB I/Hep B I* (Pentavalent I)	() 0	0
DPT II/HiB II* (Tetravalent)	() 0	0
DPT III/HiB III/Hep B II* (Pentavalent II)	(0 0	0
Polio 0	() 0	0
Polio I	() 0	0
Polio II	(0 0	0
Polio III	() 0	0
Measles	() 0	0
MMR	() 0	0
Rotavirus 1	(0 0	0
Rotavirus 2	(0 0	0
Rotavirus 3	(0 0	0
Vitamin A	(0 0	0
Additional vaccines	(0 0	0

Total vaccinated		0
a. Program dropout rate*		-22%
b. Immunisation coverage	BCG	52%
	Hepatitis B birth of	52%
	DPT I	86%
	DPT III	105%
	Polio I	86%
	Polio III	105%
	Measles	107%
	MMR**	129%
	Rotavirus I	86%
	Rotavirus III	105%
c. Vitamin A coverage		100%
*If program dropout <0%, please review dat	ta recording and reporting proce	dures for m

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

UNHCR

7. Reproductive Health

ANC Indicators

7.1 Antenatal care					
			< 18	≥ 18	Total
First antenatal visit < 1st trir	nester	[
First antenatal visit > 1st trir	nester				
Number of pregnant	≥ 4 ANC visits		N	D DA1	A
women at time of delivery	2 TT doses				
who received:	anaemia screening				
ND: no data submitted	5				

ime antenatal visits made < 1st trimeste

7.2 Postnatal care

	< 18	≥ 18	Total
Women who attended 3 postnatal visits within 6 wks of delivery	N	O DAT	ΓA Ι
ND: no data submitted			

7.3 Sexual and gender-based violence

	< 18		≥18		Total
	м	F	м	F	TOLAI
Total no rape survivors seen	0	0	0	0	(
No. of rape survivors examined within 72 hours	0	0	0	0	(
No. of rape survivors examined within 72 -120 hours	0	0	0	0	(
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	(
No. of rape survivors examined after 2 weeks	0	0	0	0	(
No of rape survivors given PEP within 72 hours	0	0	0	0	(
No of female rape survivors given ECP within 120 hours		0		0	(
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	(
No. cases of trauma in health centre due to domestic violence	0	0	0	0	(

Incidence of reported rape	0.
Proportion of rape survivors who receive PEP < 72 hrs	
Proportion of female rape survivors who receive ECP < 120 hrs	
Proportion of rape survivors who receive STI presumptive treatment < 2 wks	

Publication date: 1. Health Information System 15 May 2014 Data Sources Erbil Sandra Harlass, UNHCR Erbil Contact Information Dr Raad Najmaddin, DOH Erbil Email: RAAD976@yahoo.com Email: HARLASS@unhcr.org INFPA unicef



