


### Refugee population

Total refugee population:

**2720** \*

Demographic breakdown:



### Key facts

Camp opened:

**September 2013**

Planned capacity:

**10,000**

Area:

**1.15 Km sq**

Population:

**Syrian refugees, majority from Qamishli**

\* Source of population data: Estimates from UNHCR registration.

### Indicators at a glance

Indicator	Value	Standard
1 Health utilisation rate	7.4	1 - 4
2 Number of consultations per clinician per day	32	< 50
3 Proportion of first time antenatal visits made < 1st trimester	N/A	≥ 90%
4 Measles immunisation coverage	81%	> 90%
5 Polio III immunisation coverage	51%	> 90%
6 Crude Mortality Rate (CMR) (/1000/month)	0.0	< 1.5
7 Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 3.0
8 Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0.0	< 40

### Notes

#### HIGHLIGHTS

Health facility utilisation rate, above expected range.


This public health profile for Gawilan refugee camp is compiled from health data obtained from reporting agencies. Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Agencies reporting:


**PU-AMI  
DOH - Dohuk  
UPP**

### Morbidity

#### Acute health conditions (%)



#### Chronic health conditions (%)



#### Top 5 acute health conditions


Condition	% morbidity	# of cases
1 Upper Respiratory Tract Infection (URTI)	22%	317
2 Lower Respiratory Tract Infection (LRTI)	10%	148
3 Watery diarrhoea*	6%	84
4 Skin infection	6%	83
5 Dental Conditions	5%	66
- Other	52%	746

#### Top 5 chronic health conditions


Condition	% morbidity	# of cases
1 Hypertension	52%	16
2 Diabetes	32%	10
3 Asthma	10%	3
4 Cardiovascular disease (other)	6%	2
5 Ischaemic heart disease	0%	0
- Other	0%	0

#### Breakdown of acute health conditions

##### by age




##### by gender




#### Breakdown of chronic health conditions


##### by age




##### by gender




Number of consultations




Weekly health utilisation rates




Watery diarrhoea




Bloody diarrhoea




Acute Jaundice Syndrome




Suspected measles




Suspected Meningitis




Acute flaccid paralysis



Upper respiratory tract infections



Lower respiratory tract infections



Calculation of disease alert thresholds:

Watery diarrhoea = 1.5 x baseline,      Bloody diarrhoea = 1.5 x baseline,      Acute Jaundice Syndrome = 1.5 x baseline,      Upper respiratory tract infections = 1.5 x baseline,  
 Meningitis = 5 cases,      Acute Flaccid Paralysis = 1 case,      Measles = 1 case,  
 baseline = average incidence or number of cases over the previous 3 weeks

Health Impact	Indicator	Standard	Outbreak Alert and Response	Indicator	Standard
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 1.5	Number of outbreaks reported	0	
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 3.0	% of outbreaks investigated within 48 hours	N/A	100%
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0.0	< 40	<b>Routine immunization</b>		
			BCG immunisation coverage	73%	> 90%
			DPT 1 immunisation coverage	99%	> 90%
			Measles immunisation coverage	81%	> 90%
			Polio III immunisation coverage	51%	> 90%
			MMR immunisation coverage	195%	> 90%
			Vitamin A coverage	9%	> 90%
Access and utilization			Sexual and gender-based violence		
No. of health facilities	1.0	1 : < 10,000	Incidence of reported rape	0.0	
Total number of consultations	1,547				
Health utilisation rate	7.4	1 - 4			
No. of consultations per trained clinician per day	32	< 50			
Reproductive health					
Proportion of first time antenatal visits made < 1st trimester	N/A	≥ 90%			

Top 3 Mental health conditions


	% morbidity	# of cases
1 Severe emotional disorder	73%	30
2 Psychotic disorder	12%	5
3 Epilepsy / seizures	7%	3
- Other	7%	3

Top 3 injuries


	% morbidity	# of cases
1 Burns	45%	14
2 Injury (unintentional)	26%	8
3 Injury (intentional)	3%	1
- Other	26%	8

Breakdown of mental health conditions


By cause



By age




By gender




Breakdown of injuries


By cause



By age



By gender



We gratefully acknowledge the support of the following agencies:



View interactive maps and statistics online  
<http://twine.unhcr.org>

