





Refugee population

Total refugee population: **79,035**

Demographic breakdown:



Key facts

Camp opened: **July 2012**

Indicators at a glance

| | | |
|---|-----|---|
| 1 Health utilisation rate | 9.2 | ✗ |
| 2 Number of consultations per clinician per day | 36 | ✓ |
| 3 Hospitalisation rate (/1000/month) | 3.7 | |
| 4 % deliveries attended by skilled personnel | 98% | ✓ |
| 5 Crude Mortality Rate (CMR) (/1000/month) | 0.3 | ✓ |
| 6 Under-five Mortality Rate (U5MR) (/1000/month) | 0.4 | ✓ |
| 7 Neonatal Mortality Rate (NNMR) (/1000 livebirths) | 9.9 | ✓ |

Notes


This public health profile for Za'atri refugee camp is compiled from health facility data from the following partners:

- IMC Clinic
- JHAS Clinic
- Moroccan Field Hospital
- RAF Qatari Clinic
- Saudi Clinic
- IRC Clinic
- AMR Clinic 1 (Comprehensive Charitable Medical Center)
- AMR Clinic 2 (Maternity and Childhood center)

Data is collected using standardised Health Information System (HIS) data collection and

Morbidity


Acute health conditions (%)



Top 5 acute health conditions

| Rank | Condition | % morbidity | # of cases |
|------|--|-------------|------------|
| 1 | Upper Respiratory Tract Infection (URTI) | 32% | 31,611 |
| 2 | Influenza-like illness (ILI) | 12% | 11,688 |
| 3 | Dental Conditions | 10% | 9,339 |
| 4 | Lower Respiratory Tract Infection (LRTI) | 6% | 5,368 |
| 5 | Ear infection | 4% | 4,152 |
| - | Other | 36% | 35,364 |

Chronic health conditions (%)





Top 5 chronic health conditions

| Rank | Condition | % morbidity | # of cases |
|------|--------------------------|-------------|------------|
| 1 | Hypertension | 21% | 5,736 |
| 2 | Diabetes | 16% | 4,460 |
| 3 | Asthma | 11% | 3,131 |
| 4 | Thyroid problems | 8% | 2,204 |
| 5 | Cardiovascular disorders | 5% | 1,362 |
| - | Other | 38% | 10,465 |

Breakdown of acute health conditions

by age



by gender



Breakdown of chronic health conditions

by age

by gender



UNHCR gratefully acknowledges the support of the following partners:



Public Health

Health Impact

| | Indicator | Standard | |
|---|-----------|----------|---|
| Crude Mortality Rate (CMR) (/1000/month) | 0.3 | < 0.51 | ✓ |
| Under-five Mortality Rate (U5MR) (/1000/month) | 0.4 | < 1.5 | ✓ |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | 9.9 | < 14.9 | ✓ |

Access and utilization

| | | | |
|--|----------|--------------|---|
| No. of health facilities | 1 : 9880 | 1 : < 10,000 | ✗ |
| Total number of consultations | 181,762 | | |
| Health utilisation rate | 9.2 | 1 - 4 | ✗ |
| No. of consultations per trained clinician per day | 36 | < 50 | ✓ |

Outbreak Alert and Response


| | Indicator | Standard |
|---|-----------|----------|
| Number of outbreaks reported | 0 | |
| % of outbreaks investigated within 48 hours | | 100% |

Laboratory


| | |
|------------------------------|--------|
| Number of laboratory samples | 39,045 |
|------------------------------|--------|

Breakdown of mental health conditions


By cause



By age




By gender




Breakdown of injuries


By cause



By age



By gender



In-patient and referral

In-patient department

| | Indicator | Standard | |
|--|-----------|----------|---|
| Hospitalisation rate (/1000/year) | 44.6 | 50 - 150 | ✗ |
| Bed occupancy rate | 22% | > 75% | ✗ |
| Mean length of stay for authorised discharges (days) | 0.9 | | |

Overall case fatality rate

| | |
|----------------------------|----|
| Overall case fatality rate | 0% |
|----------------------------|----|

Proportion of deaths within 24 hours of admission

| | |
|---|----|
| Proportion of deaths within 24 hours of admission | 0% |
|---|----|

Reproductive Health

Antenatal care

| | Indicator | Standard | |
|---|-----------|----------|---|
| Coverage of antenatal care (4 or more visits) | 73% | ≥ 90% | ✗ |
| Coverage of TT vaccination | 71% | ≥ 90% | ✗ |
| Coverage of anaemia screening in pregnancy | 94% | ≥ 90% | ✓ |

Delivery care

| | | | |
|---|-----|---------|---|
| Proportion of deliveries attended by skilled personnel | 98% | ≥ 90% | ✓ |
| Proportion of deliveries performed by caesarean section | 42% | 5 - 15% | ✗ |
| Proportion of low birth weight deliveries (<2500g) | 0% | < 15% | ✓ |

Maternal mortality

| | | | |
|------------------------------------|---------|------|--|
| Number of maternal deaths reported | 1 | | |
| % of maternal deaths investigated | #DIV/0! | 100% | |

Postnatal care

| | Indicator | Standard | |
|----------------------------|-----------|----------|---|
| Coverage of postnatal care | 70% | ≥ 90% | ✗ |

Sexual and Gender-based Violence

| | | | | |
|--|-----|------|---|---|
| Incidence of rape (reported cases/10,000/year)* | 0.0 | | ^ | ✗ |
| % of rape survivors who received PEP** < 72 hours | | 100% | ^ | ✗ |
| % of rape survivors who received ECP*** < 120 hours | | 100% | ^ | ✗ |
| % of rape survivors who received STI**** prophylaxis < 2 wks | | 100% | ^ | ✗ |

*Only includes rapes reported at health facilities **Post-exposure prophylaxis ***Emergency contraceptive pill ****Sexually transmitted infection



View interactive maps and statistics online
<http://twine.unhcr.org>

Legend:

- Standard met
- Standard borderline
- Standard not met
- Interpret data with care
- Data missing

