

Population	0 to < 1		Total <1	1 to 4		Total 1 to 4	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total
	M	F		M	F		M	F		M	F		M	F		
Refugee population	655	619	1,274	3,453	3,279	6,732	7,138	6,702	13,840	6,664	7,164	13,829	300	414	714	36,389
% by age group			4%			19%			38%			38%			2%	100%

### 2.1 Mortality

Record ONE cause of mortality per case	0 to 4		Total <5	U5 % mort	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total	Crude % mort
	M	F			M	F		M	F		M	F			
1. Lower Respiratory Tract Infection (LRTI)	3	1	4	19%	0	0	0	0	0	0	0	0	0	4	12%
2. Watery diarrhoea	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%
3. Bloody diarrhoea	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%
4. Ischaemic heart disease	0	0	0	0%	0	0	0	1	0	1	1	0	1	2	6%
5. Cardiovascular disorders (other)	0	1	1	5%	0	0	0	0	0	0	1	0	1	2	6%
6. Cerebrovascular disease	0	1	1	5%	0	0	0	0	0	0	1	1	2	3	9%
7. Chronic kidney disease	1	0	1	5%	0	0	0	0	1	1	0	0	2	6%	
8. Injury	0	0	0	0%	0	0	0	0	1	1	0	0	1	3%	
9. Cancer	0	0	0	0%	0	0	0	1	4	5	1	0	1	6	18%
10. Chronic obstructive pulmonary disease (COPD)	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%
11. Tuberculosis (confirmed)	0	0	0	0%	0	0	0	0	0	0	0	0	0	0	0%
12. Neonatal death	8	5	13	62%	0	0	0	0	0	0	0	0	0	13	38%
13. Other	0	1	1	5%	0	0	0	0	0	0	0	0	0	1	3%
<b>Total</b>	<b>12</b>	<b>9</b>	<b>21</b>	<b>100%</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>2</b>	<b>6</b>	<b>8</b>	<b>4</b>	<b>1</b>	<b>5</b>	<b>34</b>	<b>100%</b>
% by age group			62%				0%			24%			15%	100%	

Mortality indicators	Male	Female	Total
Crude Mortality Rate (CMR) (/1000 population/month)	0.2	0.1	0.2
Under 5 Mortality Rate (U5MR) (/1000 population/month)	0.5	0.4	0.4
Neonatal Mortality Rate (NNMR) (/1000 population/month)			17.8

### 2.2 Maternal death

Number of maternal deaths reported	0
Number of maternal deaths investigated	0
Proportion of maternal deaths investigated	

### 3.1 Staffing and service availability

Number of full-time clinicians	23.1
Average number of full days on which each health facility was functioning	5.6
Total number of outpatient consultations	140,731
Health facility utilisation rate	7.7
Consultations per clinician per day	41

### 3.2 Acute health conditions

Record NEW cases of disease only	0 to 4		Total <5	U5 Incid	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total	% by cause	Crude Incide
	M	F			M	F		M	F		M	F				
Upper Respiratory Tract Infection (URTI)	6,272	5,129	11,401	1424.1	3,241	3,568	6,809	2,883	3,677	6,560	177	230	407	25,177	27%	691.9
Influenza-like illness (ILI)	1,513	1,157	2,670	333.5	1,168	1,124	2,292	1,066	1,288	2,354	91	86	177	7,493	8%	205.9
Lower Respiratory Tract Infection (LRTI)	1,231	947	2,178	272.1	126	115	241	140	177	317	16	28	44	2,780	3%	76.4
Watery diarrhoea*	1,096	985	2,081	259.9	184	210	394	211	321	532	17	18	35	3,042	3%	83.6
Bloody diarrhoea*	2	2	4	0.5	3	2	5	1	2	3	1	1	2	14	0%	0.4
Skin infection	598	591	1,189	148.5	557	697	1,254	560	627	1,187	48	34	82	3,712	4%	102.0
Eye infection	422	318	740	92.4	289	238	527	387	333	720	40	29	69	2,056	2%	56.5
Ear infection	535	440	975	121.8	222	308	530	335	602	937	42	39	81	2,523	3%	69.3
Dental Conditions	121	131	252	31.5	1,008	1,052	2,060	2,703	3,012	5,715	64	70	134	8,161	9%	224.3
Intestinal Worms	136	130	266	33.2	227	216	443	41	58	99	18	5	23	831	1%	22.8
Acute Jaundice Syndrome*	27	22	49	6.1	8	4	12	1	1	2	0	0	0	63	0%	1.7
Measles (suspected)*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Meningitis (suspected)*	0	0	0	0.0	1	1	2	1	0	1	0	0	0	3	0%	0.1
Acute Flaccid Paralysis*	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0%	0.0
Leishmaniasis	0	1	1	0.1	0	2	2	0	0	0	0	0	0	3	0%	0.1
Fever of unknown origin	1	0	1	0.1	1	1	2	0	0	0	0	0	0	3	0%	0.1
Tuberculosis (suspected)	0	0	0	0.0	0	2	2	1	0	1	0	0	0	3	0%	0.1
Other	3,972	4,053	8,025	1002.4	3,334	4,914	8,248	7,229	12,623	19,852	727	894	1,621	37,746	40%	1037.3
<b>Total</b>	<b>15,926</b>	<b>13,906</b>	<b>29,832</b>	<b>3726.4</b>	<b>10,369</b>	<b>12,454</b>	<b>22,823</b>	<b>15,559</b>	<b>22,721</b>	<b>38,280</b>	<b>1,241</b>	<b>1,434</b>	<b>2,675</b>	<b>93,610</b>	<b>100%</b>	<b>2572.5</b>
% by age group			32%				24%			41%			3%	100%		

\* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

### 3.3 Chronic health conditions

Record NEW and REVISITS	0 to 4		Total < 5	5 to 17		Total 5 to 17	18 to 59		Total 18 to 59	> 60		Total > 60	Grand total	% by cause
	M	F		M	F		M	F		M	F			
Diabetes	4	5	9	16	44	60	409	505	914	300	458	758	1,741	20%
Hypertension	0	0	0	1	0	1	646	1,010	1,656	516	809	1,325	2,982	35%
Asthma	338	192	530	173	115	288	132	201	333	48	43	91	1,242	14%
Ischaemic heart disease	0	0	0	0	1	1	140	82	222	192	82	274	497	6%
Cardiovascular disease (other)	4	1	5	4	3	7	55	58	113	35	22	57	182	2%
Chronic obstructive pulmonary disease (COPD)	0	0	0	0	0	0	16	10	26	44	11	55	81	1%
Liver disease	0	1	1	0	2	2	1	3	4	0	1	1	8	0%
Thyroid problems	8	6	14	12	9	21	61	291	352	7	34	41	428	5%
Cancer	0	0	0	0	0	0	6	10	16	4	1	5	21	0%
Other	124	107	231	74	71	145	321	463	784	151	117	268	1,428	17%
<b>Total</b>	<b>478</b>	<b>312</b>	<b>790</b>	<b>280</b>	<b>245</b>	<b>525</b>	<b>1,787</b>	<b>2,633</b>	<b>4,420</b>	<b>1,297</b>	<b>1,578</b>	<b>2,875</b>	<b>8,610</b>	<b>100%</b>
% by age group			9%			6%			51%			33%	100%	



## 5.2 Referrals by service

		0 to 4	Total < 5	5 to 17	Total 5 to 17	18 to 59	Total 18 to 59	> 60	Total	Grand total
1.	Paediatrics	827	827	147	147	1	1	0	0	975
2.	Internal medicine	60	60	304	304	526	526	92	92	982
3.	Obstetric	0	0	0	0	73	73	0	0	73
4.	Gynaecological	0	0	11	11	52	52	0	0	63
5.	Surgical	236	236	590	590	646	646	73	73	1545
6.	Other	79	79	180	180	232	232	52	52	543

	Total
Number of laboratory tests conducted	36,916

## 8.1 Antenatal care

	< 18	≥ 18	Total
First antenatal visit < 1st trimester	67	799	866
First antenatal visit > 1st trimester	36	627	663

### ANC Indicators

Proportion of first time antenatal visits made < 1st trimester	57%
--	-----

## 8.2 Antenatal care in facility

Number of pregnant women at time of delivery who received:	≥ 4 ANC visits	2 TT doses	anaemia screening	
	19	18	33	457
				191
				722

Coverage of complete antenatal care	63%
Coverage of antenatal tetanus immunisation	26%
Coverage of anaemia screening in pregnancy	99%

## 8.3 Delivery care

	< 18	≥ 18	Total
Live births	33	698	731
Still births	1	8	9
Low birth weight (<2500 g)	1	48	49
Attended by skilled health worker	33	666	699
Number of obstetric complications treated	2	36	38
Number of caesarean sections performed	3	134	137

### Delivery Indicators

Crude birth rate (/1000 population/ month)	3.3
Proportion of births attended by skilled health worker (excl. TBA)	96%
Proportion of low birth weight deliveries	7%
Proportion of all births performed by caesarean section	19%
Proportion of obstetric complications treated among girls < 18	6%

## 8.4 Postnatal care

	< 18	≥ 18	Total
At least three PNC visits within 6 weeks	42	716	758

### Postnatal Indicators

Coverage of complete postnatal care	104%
-------------------------------------	------

## 8.5 Family planning

	New		Repeat		Total
	< 18	≥ 18	< 18	≥ 18	
Oral contraceptives	27	581	541	845	1,994
Emergency Contraceptive Pill	0	2	0	14	16
Injectable (Depo-Provera)	0	77	27	42	146
Implantable (Norplant)	0	0	0	0	0
Intra-Uterine Device (IUD)	1	72	0	5	78
Condom (Male)	42	934	440	340	1,756
Other	7	44	0	59	110

	< 18	≥ 18
Family Planning Counseling*	319	4,851

## 8.6 Sexual and gender-based violence

	< 18		≥ 18		Total
	M	F	M	F	
Total no rape survivors seen	0	0	0	1	1
No. of rape survivors examined within 72 hours	0	0	0	0	0
No. of rape survivors examined within 72 -120 hours	0	0	0	0	0
No. of rape survivors examined within 120 hrs - 2 weeks	0	0	0	0	0
No. of rape survivors examined after 2 weeks	0	0	0	1	1
No of rape survivors given PEP within 72 hours	0	0	0	0	0
No of female rape survivors given ECP within 120 hours		0		0	0
No of rape survivors given STI presumptive treatment within 2 weeks	0	0	0	0	0
No. cases of trauma in health centre due to domestic violence	0	0	0	13	13

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators	
Incidence of reported rape (/10,000 population/year)	0.5
Proportion of rape survivors who receive PEP < 72 hrs	
Proportion of female rape survivors who receive ECP < 120 hrs	
Proportion of rape survivors who receive STI presumptive treatment < 2 wks	

### Note

*This report for non-camp Syrian refugees in Jordan is compiled from health data obtained from reporting agencies and clinics. Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.*

Published on

Contact  
Information

UNHCR  
Adam Musa Khalifa  
Email: KHALIFAA@unhcr.org

We gratefully  
acknowledge the contributions of  
the following agencies:

